


Development of a Core Outcome Set (COS) for treatment of depression during or after pregnancy (antenatal and postpartum depression, report 314 (2020))

Appendix 6 Results from Survey 1

Outcomes to measure depression




The result presented in percentage, both in total as well as by the different perspectives. The most common score for each group is highlighted.

Outcome	Perspective	1	2	3	4	5	6	7	8	9
Depressive symptoms	All results	1%	0%	3%	3%	5%	19%	23%	15%	32%
	Researcher within the field	3%	0%	0%	3%	3%	13%	13%	17%	47%
	HTA-agencies policy makers and others	0%	0%	0%	0%	18%	27%	36%	0%	18%
	Patients	0%	0%	2%	0%	9%	19%	12%	19%	40%
	Health professional	0%	0%	4%	4%	1%	19%	31%	15%	24%
Depression in partner	All results	3%	3%	15%	9%	9%	30%	13%	9%	7%
	Researcher within the field	10%	3%	17%	17%	7%	20%	17%	7%	3%
	HTA-agencies policy makers and others	0%	9%	27%	0%	9%	18%	27%	0%	9%
	Patients	0%	7%	7%	12%	7%	26%	7%	16%	12%
	Health professional	1%	0%	18%	4%	10%	40%	12%	6%	4%
Psychiatric diagnosis of depression	All results	2%	1%	7%	5%	10%	14%	21%	19%	21%
	Researcher within the field	7%	0%	3%	3%	3%	13%	23%	30%	17%
	HTA-agencies policy makers and others	0%	0%	9%	9%	0%	0%	0%	27%	55%
	Patients	0%	0%	7%	5%	12%	21%	16%	16%	21%
	Health professional	1%	1%	7%	6%	13%	12%	27%	13%	16%
Recovery from depression (remission)	All results	2%	1%	3%	8%	9%	20%	17%	15%	23%
	Researcher within the field	3%	7%	0%	7%	7%	17%	13%	13%	33%
	HTA-agencies policy makers and others	0%	0%	9%	0%	9%	9%	18%	9%	45%
	Patients	2%	0%	0%	12%	7%	21%	21%	14%	21%

	Health professional	1%	0%	6%	7%	12%	22%	16%	16%	15%
Depression relapse	All results	1%	0%	1%	7%	9%	23%	22%	13%	21%
	Researcher within the field	3%	0%	0%	10%	0%	23%	23%	20%	17%
	HTA-agencies policy makers and others	0%	0%	9%	9%	0%	18%	36%	0%	27%
	Patients	0%	0%	2%	7%	14%	14%	23%	14%	23%
	Health professional	0%	0%	0%	6%	12%	30%	18%	10%	19%
Time to remission from depression	All results	2%	2%	3%	9%	11%	25%	17%	13%	14%
	Researcher within the field	3%	3%	3%	10%	10%	20%	13%	13%	20%
	HTA-agencies policy makers and others	0%	0%	9%	18%	0%	0%	27%	18%	18%
	Patients	5%	2%	2%	2%	14%	26%	16%	9%	23%
	Health professional	0%	1%	3%	12%	10%	30%	18%	15%	4%
Was depression treatment initiated	All results	1%	0%	1%	6%	10%	20%	17%	14%	28%
	Researcher within the field	3%	0%	3%	17%	7%	7%	20%	13%	27%
	HTA-agencies policy makers and others	0%	0%	0%	0%	9%	18%	18%	27%	18%
	Patients	0%	0%	0%	0%	7%	19%	14%	12%	47%
	Health professional	1%	0%	1%	6%	13%	27%	18%	13%	18%
Change in antidepressant medication	All results	4%	1%	11%	11%	11%	23%	17%	6%	11%
	Researcher within the field	7%	3%	17%	13%	3%	17%	30%	7%	3%
	HTA-agencies policy makers and others	0%	0%	9%	18%	9%	9%	18%	0%	27%
	Patients	7%	0%	2%	9%	12%	19%	21%	9%	14%
	Health professional	1%	0%	13%	10%	15%	30%	9%	4%	10%

Expert rating of overall disease burden	All results	3%	1%	8%	15%	9%	21%	9%	12%	13%
	Researcher within the field	7%	0%	3%	20%	13%	30%	7%	10%	10%
	HTA-agencies policy makers and others	0%	9%	9%	18%	9%	9%	18%	0%	18%
	Patients	2%	0%	7%	12%	5%	12%	7%	21%	16%
	Health professional	3%	1%	10%	15%	9%	25%	10%	9%	12%

Other psychiatric outcomes



The result presented in percentage, both in total as well as by the different perspectives. The most common score for each group is highlighted.


Outcome	Perspective	1	2	3	4	5	6	7	8	9
Anxiety symptoms	All results	0%	1%	5%	7%	9%	19%	28%	9%	17%
	Researcher within the field	0%	0%	7%	7%	10%	17%	30%	3%	20%
	HTA-agencies policy makers and others	0%	0%	18%	0%	18%	18%	27%	9%	0%
	Patients	0%	0%	2%	0%	7%	21%	33%	12%	23%
	Health professional	0%	1%	3%	12%	7%	18%	25%	10%	15%
Fear of childbirth	All results	1%	1%	7%	15%	11%	17%	14%	11%	19%
	Researcher within the field	0%	3%	7%	20%	7%	23%	13%	13%	7%
	HTA-agencies policy makers and others	0%	0%	36%	18%	0%	9%	27%	0%	0%
	Patients	2%	2%	7%	7%	7%	16%	9%	12%	33%
	Health professional	0%	0%	3%	16%	16%	15%	15%	12%	18%
Psychiatric diagnosis of anxiety	All results	1%	1%	6%	11%	7%	19%	21%	12%	17%
	Researcher within the field	0%	0%	7%	13%	3%	27%	10%	13%	20%
	HTA-agencies policy makers and others	0%	0%	9%	0%	0%	18%	27%	9%	27%
	Patients	0%	2%	2%	12%	7%	14%	23%	14%	19%
	Health professional	1%	0%	7%	12%	9%	19%	22%	10%	13%
Other psychiatric or neuropsychi	All results	1%	1%	5%	9%	10%	24%	20%	11%	11%
	Researcher within the field	0%	0%	3%	10%	3%	27%	17%	13%	20%

atric diagnoses	HTA-agencies policy makers and others	0%	9%	0%	18%	9%	9%	18%	18%	0%
	Patients	2%	0%	5%	9%	19%	12%	23%	9%	9%
	Health professional	0%	1%	7%	7%	7%	33%	19%	9%	10%
Mood	All results	0%	1%	10%	8%	5%	23%	15%	16%	16%
	Researcher within the field	0%	0%	13%	0%	0%	20%	13%	13%	30%
	HTA-agencies policy makers and others	0%	0%	18%	0%	9%	27%	18%	0%	9%
	Patients	0%	0%	7%	5%	2%	33%	19%	19%	14%
	Health professional	0%	3%	9%	15%	7%	16%	12%	18%	12%
Self-harm	All results	1%	1%	4%	9%	6%	17%	21%	15%	23%
	Researcher within the field	0%	3%	7%	7%	3%	20%	13%	7%	33%
	HTA-agencies policy makers and others	0%	0%	0%	9%	9%	36%	9%	0%	27%
	Patients	2%	0%	2%	12%	7%	5%	21%	14%	30%
	Health professional	0%	0%	4%	7%	6%	19%	25%	21%	12%
Suicidal thoughts or attempts	All results	1%	1%	0%	5%	1%	5%	14%	15%	53%
	Researcher within the field	0%	0%	0%	10%	0%	7%	7%	17%	53%
	HTA-agencies policy makers and others	0%	0%	0%	0%	0%	0%	27%	0%	55%
	Patients	0%	0%	0%	5%	2%	2%	14%	12%	60%
	Health professional	1%	1%	0%	4%	0%	6%	15%	18%	48%
	All results	2%	3%	14%	13%	6%	19%	15%	11%	13%

Cognitive ability	Researcher within the field	3%	3%	20%	17%	3%	20%	10%	7%	10%
	HTA-agencies policy makers and others	0%	18%	18%	0%	18%	9%	9%	18%	0%
	Patients	0%	0%	12%	9%	2%	7%	30%	9%	28%
	Health professional	3%	1%	12%	16%	7%	28%	7%	13%	6%
Eating disorder symptoms	All results	3%	2%	13%	11%	11%	17%	16%	10%	9%
	Researcher within the field	0%	3%	13%	13%	10%	20%	13%	10%	10%
	HTA-agencies policy makers and others	0%	0%	18%	9%	9%	18%	27%	0%	0%
	Patients	9%	2%	14%	12%	9%	7%	12%	12%	14%
	Health professional	1%	1%	10%	10%	13%	21%	18%	10%	7%
Visit to clinic for fear of childbirth	All results	3%	2%	11%	9%	11%	20%	11%	12%	15%
	Researcher within the field	0%	7%	23%	7%	7%	23%	13%	10%	3%
	HTA-agencies policy makers and others	9%	0%	36%	0%	18%	9%	9%	9%	0%
	Patients	5%	0%	5%	0%	12%	16%	12%	12%	35%
	Health professional	1%	1%	6%	18%	12%	22%	9%	13%	10%
Post-traumatic stress symptoms	All results	0%	1%	5%	8%	8%	23%	17%	15%	17%
	Researcher within the field	0%	3%	7%	3%	10%	27%	23%	10%	10%
	HTA-agencies policy makers and others	0%	0%	27%	9%	9%	27%	9%	0%	9%
	Patients	0%	0%	0%	7%	9%	21%	12%	19%	23%
	Health professional	0%	0%	4%	10%	6%	22%	19%	16%	18%

Psychiatric diagnosis of postpartum stress syndrome in relation with childbirth	All results	1%	0%	5%	9%	9%	13%	15%	15%	27%
	Researcher within the field	3%	0%	3%	17%	10%	13%	13%	17%	17%
	HTA-agencies policy makers and others	0%	0%	18%	0%	27%	0%	27%	0%	18%
	Patients	0%	0%	0%	9%	7%	7%	7%	19%	42%
	Health professional	0%	0%	6%	7%	7%	19%	19%	13%	24%
Psychiatric diagnosis of postpartum psychosis	All results	1%	1%	3%	9%	6%	15%	16%	15%	26%
	Researcher within the field	0%	3%	3%	7%	10%	10%	10%	13%	37%
	HTA-agencies policy makers and others	0%	9%	9%	18%	0%	0%	9%	18%	27%
	Patients	2%	0%	2%	9%	2%	23%	19%	12%	19%
	Health professional	0%	0%	3%	7%	7%	13%	18%	18%	27%

Outcomes relating to family circumstances



The result presented in percentage, both in total as well as by the different perspectives. The most common score for each group is highlighted.


Outcome	Perspective	1	2	3	4	5	6	7	8	9
Domestic/ intimate partner violence	All results	1%	2%	4%	9%	7%	11%	8%	11%	37%
	Researcher within the field	0%	7%	3%	13%	7%	17%	7%	3%	33%
	HTA-agencies policy makers and others	0%	0%	9%	18%	18%	9%	9%	9%	18%

	Patients	5%	0%	7%	9%	2%	12%	7%	9%	35%
	Health professional	0%	1%	1%	4%	9%	9%	9%	15%	43%
Family functioning	All results	1%	2%	5%	11%	6%	25%	15%	16%	11%
	Researcher within the field	0%	3%	3%	20%	3%	23%	10%	10%	17%
	HTA-agencies policy makers and others	0%	0%	18%	9%	18%	18%	27%	0%	0%
	Patients	0%	2%	5%	12%	2%	16%	16%	16%	23%
	Health professional	1%	1%	3%	6%	7%	31%	15%	21%	3%
Quality of relationship	All results	1%	1%	16%	15%	15%	19%	12%	9%	5%
	Researcher within the field	3%	0%	17%	7%	7%	23%	13%	10%	10%
	HTA-agencies policy makers and others	0%	0%	27%	36%	18%	0%	9%	0%	0%
	Patients	0%	2%	16%	12%	12%	23%	14%	9%	7%
	Health professional	1%	0%	13%	18%	19%	18%	10%	9%	1%
Parent to infant bonding	All results	1%	0%	3%	8%	7%	20%	12%	14%	28%
	Researcher within the field	3%	0%	3%	13%	7%	23%	7%	13%	20%
	HTA-agencies policy makers and others	0%	0%	9%	18%	9%	9%	0%	9%	27%
	Patients	0%	0%	0%	7%	5%	19%	14%	14%	37%
	Health professional	1%	0%	3%	4%	9%	21%	15%	15%	25%
Attachment	All results	1%	1%	1%	7%	6%	15%	15%	19%	28%
	Researcher within the field	3%	0%	3%	7%	10%	17%	17%	17%	17%

	HTA-agencies policy makers and others	0%	0%	0%	27%	18%	9%	0%	18%	18%
	Patients	0%	2%	0%	7%	7%	9%	16%	21%	30%
	Health professional	0%	0%	1%	3%	1%	18%	15%	19%	34%
Parenting Sense of Competency	All results	0%	1%	5%	10%	11%	22%	14%	10%	19%
	Researcher within the field	0%	0%	7%	7%	13%	30%	20%	10%	3%
	HTA-agencies policy makers and others	0%	0%	18%	27%	18%	9%	9%	0%	9%
	Patients	0%	0%	5%	9%	12%	26%	9%	9%	26%
	Health professional	0%	1%	3%	9%	9%	18%	15%	12%	24%
Parental stress	All results	1%	0%	5%	9%	11%	29%	19%	9%	9%
	Researcher within the field	0%	0%	10%	3%	13%	33%	13%	3%	13%
	HTA-agencies policy makers and others	0%	0%	18%	18%	18%	18%	0%	18%	0%
	Patients	0%	0%	0%	9%	12%	21%	35%	9%	7%
	Health professional	1%	0%	4%	10%	9%	34%	13%	10%	9%
General stress	All results	1%	1%	11%	14%	15%	26%	13%	7%	5%
	Researcher within the field	0%	3%	10%	17%	17%	23%	7%	10%	3%
	HTA-agencies policy makers and others	0%	0%	18%	27%	18%	9%	18%	0%	0%
	Patients	0%	0%	5%	12%	12%	26%	21%	12%	9%
	Health professional	3%	0%	15%	12%	15%	30%	10%	3%	3%
	All results	1%	1%	2%	7%	6%	15%	17%	13%	30%

Parent-infant interaction	Researcher within the field	0%	3%	3%	7%	0%	20%	23%	13%	20%
	HTA-agencies policy makers and others	0%	9%	0%	18%	18%	9%	18%	0%	18%
	Patients	0%	0%	2%	14%	7%	14%	19%	5%	33%
	Health professional	1%	0%	1%	1%	6%	13%	12%	19%	36%
Attitudes about parenting	All results	2%	3%	15%	13%	7%	25%	14%	7%	6%
	Researcher within the field	3%	3%	20%	13%	7%	27%	10%	3%	3%
	HTA-agencies policy makers and others	0%	9%	45%	9%	0%	9%	9%	9%	0%
	Patients	2%	0%	19%	9%	7%	21%	12%	9%	14%
	Health professional	1%	4%	6%	15%	7%	28%	18%	6%	3%
Family planning	All results	4%	2%	17%	17%	11%	16%	6%	11%	10%
	Researcher within the field	7%	7%	20%	17%	13%	17%	3%	0%	7%
	HTA-agencies policy makers and others	0%	0%	36%	27%	0%	9%	0%	18%	0%
	Patients	2%	0%	7%	12%	12%	14%	7%	16%	26%
	Health professional	4%	1%	19%	18%	10%	18%	7%	12%	3%

Outcomes relating to function




The result presented in percentage, both in total as well as by the different perspectives. The most common score for each group is highlighted.

Outcome	Perspective	1	2	3	4	5	6	7	8	9
Daily functioning level/ activities of daily living	All results	0%	1%	5%	7%	12%	23%	19%	11%	13%
	Researcher within the field	0%	0%	10%	3%	3%	30%	17%	13%	10%
	HTA-agencies policy makers and others	0%	9%	9%	0%	18%	9%	36%	9%	0%
	Patients	0%	0%	0%	5%	7%	21%	23%	16%	21%
	Health professional	0%	1%	4%	10%	18%	24%	15%	7%	10%
Sleep	All results	1%	2%	4%	5%	10%	17%	19%	9%	25%
	Researcher within the field	0%	7%	3%	3%	3%	23%	13%	7%	30%

	HTA-agencies policy makers and others	0%	9%	9%	9%	27%	0%	18%	0%	18%
	Patients	0%	0%	0%	2%	12%	12%	33%	5%	33%
	Health professional	1%	0%	6%	7%	9%	21%	13%	13%	18%
Fatigue	All results	1%	1%	7%	6%	12%	23%	18%	10%	13%
	Researcher within the field	0%	3%	3%	7%	13%	40%	10%	7%	7%
	HTA-agencies policy makers and others	9%	0%	18%	9%	18%	0%	18%	9%	9%
	Patients	0%	0%	0%	0%	12%	23%	26%	12%	21%
	Health professional	0%	1%	10%	9%	10%	19%	16%	10%	12%
Quality of life	All results	1%	1%	5%	7%	13%	22%	18%	12%	13%
	Researcher within the field	0%	7%	0%	10%	13%	17%	17%	17%	10%
	HTA-agencies policy makers and others	0%	0%	18%	0%	0%	27%	18%	9%	18%
	Patients	2%	0%	7%	5%	9%	21%	9%	19%	23%
	Health professional	1%	0%	4%	9%	18%	24%	24%	6%	6%
Sexual desire/function	All results	3%	5%	17%	20%	14%	19%	5%	3%	4%
	Researcher within the field	7%	7%	17%	20%	7%	23%	7%	3%	0%
	HTA-agencies policy makers and others	0%	18%	9%	9%	36%	9%	0%	0%	0%
	Patients	5%	0%	21%	19%	19%	19%	2%	2%	9%
	Health professional	1%	4%	15%	22%	10%	19%	7%	3%	3%
Stigma	All results	1%	2%	9%	17%	10%	14%	13%	8%	17%

	Researcher within the field	7%	3%	17%	17%	7%	13%	17%	0%	10%
	HTA-agencies policy makers and others	0%	0%	36%	9%	9%	9%	0%	18%	0%
	Patients	0%	0%	7%	12%	16%	5%	16%	14%	26%
	Health professional	0%	3%	3%	22%	7%	21%	12%	6%	16%
Social relationships	All results	0%	1%	3%	11%	14%	24%	19%	13%	7%
	Researcher within the field	0%	3%	3%	17%	3%	17%	7%	23%	17%
	HTA-agencies policy makers and others	0%	0%	18%	0%	36%	18%	0%	9%	0%
	Patients	0%	0%	0%	7%	14%	30%	28%	7%	9%
	Health professional	0%	0%	3%	12%	15%	24%	22%	12%	3%
Perceived ability to cope with life challenges	All results	3%	2%	9%	9%	13%	15%	19%	9%	11%
	Researcher within the field	13%	3%	10%	0%	10%	13%	20%	10%	10%
	HTA-agencies policy makers and others	0%	0%	18%	18%	27%	0%	0%	9%	9%
	Patients	0%	0%	9%	9%	12%	16%	19%	7%	19%
	Health professional	0%	3%	6%	12%	13%	16%	22%	10%	7%
Effort-based decision-making	All results	3%	3%	9%	13%	17%	18%	14%	7%	6%
	Researcher within the field	3%	7%	7%	10%	13%	20%	20%	3%	3%
	HTA-agencies policy makers and others	0%	9%	18%	9%	27%	0%	18%	0%	0%
	Patients	5%	0%	2%	12%	19%	21%	9%	14%	12%
	Health professional	3%	1%	13%	15%	15%	18%	13%	6%	4%

Outcomes relating to characteristics




The result presented in percentage, both in total as well as by the different perspectives. The most common score for each group is highlighted.

Outcome	Perspective	1	2	3	4	5	6	7	8	9
Self-compassion	All results	1%	4%	9%	17%	15%	17%	12%	10%	5%
	Researcher within the field	3%	13%	17%	13%	10%	20%	13%	0%	0%
	HTA-agencies policy makers and others	0%	0%	36%	27%	18%	0%	0%	9%	0%
	Patients	0%	0%	2%	14%	19%	14%	16%	19%	12%
	Health professional	0%	3%	4%	19%	15%	21%	10%	9%	3%
Lifestyle	All results	2%	4%	4%	17%	15%	15%	17%	7%	7%
	Researcher within the field	0%	3%	13%	0%	13%	17%	30%	3%	10%

	HTA-agencies policy makers and others	0%	9%	0%	27%	9%	27%	18%	0%	0%
	Patients	5%	0%	2%	23%	16%	19%	9%	14%	5%
	Health professional	1%	6%	1%	18%	16%	10%	16%	6%	9%
Verbal fluency	All results	18%	10%	23%	10%	9%	8%	3%	0%	0%
	Researcher within the field	23%	10%	23%	3%	10%	17%	0%	0%	0%
	HTA-agencies policy makers and others	36%	18%	0%	18%	0%	9%	0%	0%	0%
	Patients	16%	7%	19%	16%	9%	12%	2%	0%	0%
	Health professional	13%	10%	30%	7%	9%	1%	6%	0%	0%
Personality	All results	9%	6%	21%	14%	12%	9%	7%	5%	3%
	Researcher within the field	3%	17%	23%	13%	10%	3%	13%	0%	7%
	HTA-agencies policy makers and others	9%	0%	27%	18%	27%	0%	9%	0%	0%
	Patients	9%	5%	12%	19%	7%	16%	7%	12%	2%
	Health professional	10%	3%	24%	10%	13%	9%	3%	3%	3%
Perceived social support / Help- Seeking	All results	0%	3%	5%	11%	7%	20%	17%	16%	13%
	Researcher within the field	0%	3%	13%	17%	7%	7%	13%	13%	17%
	HTA-agencies policy makers and others	0%	9%	0%	36%	9%	27%	0%	0%	9%
	Patients	0%	0%	0%	5%	5%	21%	26%	26%	14%
	Health professional	0%	3%	6%	7%	7%	24%	15%	13%	12%
	All results	0%	3%	11%	13%	16%	14%	17%	9%	4%

Socio-emotional competence	Researcher within the field	0%	7%	20%	13%	13%	10%	20%	7%	0%
	HTA-agencies policy makers and others	0%	18%	18%	9%	27%	9%	9%	0%	0%
	Patients	0%	0%	7%	16%	7%	19%	19%	16%	7%
	Health professional	0%	1%	9%	10%	21%	13%	16%	6%	4%
Confidence to self-help	All results	1%	4%	11%	12%	11%	19%	15%	9%	7%
	Researcher within the field	0%	3%	27%	13%	10%	17%	17%	3%	0%
	HTA-agencies policy makers and others	0%	9%	18%	27%	0%	18%	0%	9%	0%
	Patients	2%	2%	7%	7%	12%	23%	19%	12%	9%
	Health professional	0%	4%	6%	12%	12%	18%	13%	10%	9%
Mentalizing, reflective functioning and level of mindfulness	All results	1%	4%	13%	17%	13%	17%	14%	6%	3%
	Researcher within the field	3%	7%	17%	23%	7%	20%	10%	3%	0%
	HTA-agencies policy makers and others	0%	27%	18%	9%	9%	18%	9%	0%	0%
	Patients	2%	0%	7%	14%	16%	16%	19%	12%	5%
	Health professional	0%	1%	15%	16%	15%	15%	13%	4%	4%
Self-esteem	All results	1%	5%	9%	12%	11%	16%	13%	16%	7%
	Researcher within the field	0%	10%	17%	17%	7%	17%	7%	13%	3%
	HTA-agencies policy makers and others	0%	9%	27%	18%	0%	27%	0%	9%	0%
	Patients	2%	0%	0%	12%	7%	16%	21%	23%	14%
	Health professional	0%	4%	9%	9%	16%	13%	13%	13%	6%

Outcomes relating to adverse events




The result presented in percentage, both in total as well as by the different perspectives. The most common score for each group is highlighted.

Outcome	Perspective	1	2	3	4	5	6	7	8	9
Adverse event	All results	1%	0%	3%	12%	7%	13%	23%	9%	21%
	Researcher within the field	3%	0%	0%	13%	3%	10%	20%	7%	33%
	HTA-agencies policy makers and others	0%	0%	0%	0%	9%	9%	27%	9%	36%
	Patients	0%	0%	2%	19%	5%	9%	26%	9%	21%
	Health professional	1%	0%	4%	9%	10%	16%	21%	10%	12%
Spontaneous abortion or stillbirth	All results	0%	1%	5%	5%	5%	15%	13%	11%	30%
	Researcher within the field	0%	0%	7%	7%	3%	20%	10%	20%	20%

	HTA-agencies policy makers and others	0%	0%	0%	0%	9%	27%	0%	27%	18%
	Patients	0%	2%	5%	5%	0%	14%	16%	5%	40%
	Health professional	0%	1%	4%	4%	7%	12%	15%	7%	30%
Perinatal mortality	All results	0%	0%	4%	6%	3%	16%	9%	9%	37%
	Researcher within the field	0%	0%	7%	10%	3%	23%	3%	17%	23%
	HTA-agencies policy makers and others	0%	0%	0%	0%	0%	27%	9%	0%	45%
	Patients	0%	0%	5%	7%	0%	14%	9%	0%	49%
	Health professional	0%	0%	3%	4%	4%	12%	10%	13%	34%

Obstetric outcomes



The result presented in percentage, both in total as well as by the different perspectives. The most common score for each group is highlighted.

Outcome	Perspective	1	2	3	4	5	6	7	8	9
Infections during pregnancy	All results	3%	3%	18%	12%	11%	13%	5%	6%	9%
	Researcher within the field	0%	3%	30%	10%	13%	7%	10%	0%	7%


	HTA-agencies policy makers and others	0%	0%	27%	0%	9%	27%	0%	9%	9%
	Patients	5%	2%	12%	14%	9%	14%	0%	9%	16%
	Health professional	4%	3%	15%	13%	12%	12%	6%	6%	4%
Residual physical problems after delivery	All results	1%	1%	5%	8%	5%	19%	19%	5%	26%
	Researcher within the field	3%	0%	10%	13%	7%	13%	20%	0%	17%
	HTA-agencies policy makers and others	0%	0%	18%	18%	9%	0%	27%	9%	9%
	Patients	0%	0%	2%	2%	5%	7%	16%	7%	53%
	Health professional	0%	1%	1%	7%	4%	33%	18%	6%	15%
Birth injury	All results	1%	1%	6%	11%	9%	17%	13%	7%	23%
	Researcher within the field	3%	0%	13%	13%	17%	7%	17%	7%	7%
	HTA-agencies policy makers and others	0%	0%	9%	9%	9%	18%	27%	0%	9%
	Patients	0%	0%	2%	5%	2%	14%	14%	2%	51%
	Health professional	1%	1%	4%	13%	9%	22%	7%	12%	13%
Pain	All results	2%	1%	9%	15%	8%	18%	14%	9%	11%
	Researcher within the field	3%	3%	10%	17%	3%	27%	10%	10%	3%
	HTA-agencies policy makers and others	0%	0%	27%	0%	18%	18%	27%	0%	0%
	Patients	2%	0%	0%	19%	7%	9%	9%	16%	28%
	Health professional	1%	1%	12%	13%	9%	19%	16%	4%	6%
	All results	1%	2%	12%	15%	10%	17%	8%	7%	10%

Gestational hypertension or preeclampsia	Researcher within the field	0%	0%	17%	13%	10%	17%	10%	7%	10%
	HTA-agencies policy makers and others	0%	18%	9%	9%	9%	18%	9%	9%	0%
	Patients	5%	0%	9%	9%	14%	16%	9%	7%	19%
	Health professional	0%	1%	12%	19%	7%	18%	6%	7%	6%
Gestational age at delivery	All results	3%	1%	5%	9%	11%	20%	15%	9%	14%
	Researcher within the field	0%	0%	13%	3%	3%	27%	13%	7%	17%
	HTA-agencies policy makers and others	0%	0%	0%	9%	27%	27%	18%	0%	0%
	Patients	5%	0%	5%	9%	5%	21%	12%	9%	30%
	Health professional	3%	1%	3%	10%	16%	15%	18%	10%	4%
Gestational diabetes	All results	4%	3%	17%	12%	9%	13%	10%	5%	7%
	Researcher within the field	0%	0%	13%	7%	17%	23%	13%	3%	7%
	HTA-agencies policy makers and others	9%	0%	9%	9%	9%	18%	18%	9%	0%
	Patients	5%	5%	21%	14%	5%	9%	5%	5%	14%
	Health professional	4%	4%	18%	13%	7%	10%	10%	6%	3%
Estimated blood loss	All results	5%	3%	9%	11%	10%	18%	9%	5%	9%
	Researcher within the field	7%	7%	10%	10%	13%	20%	7%	3%	7%
	HTA-agencies policy makers and others	9%	0%	27%	18%	0%	18%	0%	9%	0%
	Patients	5%	0%	2%	19%	9%	12%	7%	9%	19%
	Health professional	4%	4%	10%	4%	10%	21%	13%	3%	6%

Induction of labour	All results	3%	3%	13%	12%	11%	16%	9%	9%	9%
	Researcher within the field	3%	3%	20%	13%	7%	23%	3%	10%	0%
	HTA-agencies policy makers and others	0%	18%	9%	18%	18%	18%	0%	0%	0%
	Patients	2%	0%	7%	2%	14%	7%	16%	14%	26%
	Health professional	3%	3%	13%	16%	9%	18%	7%	6%	4%
Mode of delivery	All results	2%	3%	3%	12%	8%	17%	11%	13%	19%
	Researcher within the field	0%	0%	3%	20%	7%	17%	13%	13%	13%
	HTA-agencies policy makers and others	0%	0%	18%	9%	18%	27%	0%	9%	9%
	Patients	2%	2%	0%	7%	14%	9%	14%	9%	35%
	Health professional	3%	4%	1%	12%	3%	21%	10%	15%	13%
Fetal growth assessment by Ultrasound	All results	2%	5%	11%	17%	8%	16%	9%	7%	8%
	Researcher within the field	3%	7%	17%	13%	3%	20%	3%	3%	10%
	HTA-agencies policy makers and others	0%	9%	9%	18%	0%	36%	9%	0%	0%
	Patients	2%	2%	9%	16%	5%	19%	9%	12%	16%
	Health professional	1%	4%	10%	19%	13%	9%	10%	6%	3%
Complications associated with the placenta	All results	5%	5%	15%	5%	10%	15%	5%	7%	11%
	Researcher within the field	7%	0%	17%	10%	20%	10%	3%	7%	3%
	HTA-agencies policy makers and others	9%	9%	18%	9%	0%	27%	9%	0%	0%
	Patients	5%	2%	9%	2%	12%	14%	7%	5%	26%

	Health professional	3%	7%	16%	4%	6%	16%	4%	9%	7%
Vaginal and/or faecal microbiota	All results	9%	8%	19%	8%	5%	9%	6%	2%	5%
	Researcher within the field	17%	7%	13%	17%	3%	3%	10%	3%	0%
	HTA-agencies policy makers and others	0%	18%	27%	0%	9%	9%	0%	0%	0%
	Patients	5%	2%	21%	5%	7%	12%	7%	5%	16%
	Health professional	10%	10%	18%	7%	4%	9%	4%	0%	1%
Transition of care from pregnancy to postpartum	All results	0%	1%	3%	7%	7%	11%	15%	17%	27%
	Researcher within the field	0%	0%	10%	13%	10%	3%	3%	27%	13%
	HTA-agencies policy makers and others	0%	9%	9%	9%	18%	9%	18%	9%	0%
	Patients	0%	0%	0%	0%	2%	14%	14%	9%	56%
	Health professional	0%	1%	0%	9%	7%	12%	21%	18%	19%

Outcomes relating to the child



The result presented in percentage, both in total as well as by the different perspectives. The most common score for each group is highlighted.

Outcome	Perspective	1	2	3	4	5	6	7	8	9
Breastfeeding	All results	3%	4%	9%	10%	10%	18%	11%	6%	17%
	Researcher within the field	0%	7%	20%	3%	13%	13%	7%	3%	23%
	HTA-agencies policy makers and others	0%	0%	9%	9%	9%	27%	18%	0%	9%
	Patients	7%	2%	5%	5%	7%	26%	7%	2%	28%
	Health professional	1%	4%	6%	16%	10%	13%	15%	10%	7%


Child attention	All results	3%	3%	11%	10%	17%	16%	13%	5%	5%
	Researcher within the field	0%	0%	10%	13%	27%	13%	7%	3%	10%
	HTA-agencies policy makers and others	9%	9%	36%	0%	27%	0%	9%	0%	0%
	Patients	5%	2%	9%	7%	14%	16%	16%	5%	12%
	Health professional	3%	4%	9%	12%	12%	19%	13%	7%	0%
Duration of crying episodes	All results	3%	2%	13%	8%	16%	19%	15%	5%	3%
	Researcher within the field	3%	3%	23%	13%	10%	17%	3%	10%	3%
	HTA-agencies policy makers and others	0%	0%	9%	9%	27%	27%	9%	9%	0%
	Patients	2%	0%	5%	5%	21%	19%	23%	2%	9%
	Health professional	4%	3%	13%	7%	13%	19%	15%	4%	0%
Child development	All results	1%	1%	3%	13%	11%	15%	21%	11%	13%
	Researcher within the field	0%	0%	0%	17%	13%	7%	17%	13%	20%
	HTA-agencies policy makers and others	0%	0%	9%	9%	9%	0%	36%	9%	18%
	Patients	2%	2%	2%	14%	9%	19%	21%	14%	12%
	Health professional	0%	1%	3%	10%	10%	19%	21%	7%	10%
Neurodevelopmental and neurobehavioral outcomes	All results	1%	1%	4%	11%	5%	18%	21%	12%	13%
	Researcher within the field	0%	0%	3%	7%	10%	17%	17%	10%	20%
	HTA-agencies policy makers and others	0%	9%	9%	0%	0%	27%	36%	0%	9%
	Patients	0%	0%	5%	9%	9%	21%	19%	12%	12%

	Health professional	1%	0%	3%	15%	1%	15%	22%	15%	10%
Sleep	All results	1%	3%	5%	9%	9%	20%	21%	11%	9%
	Researcher within the field	0%	3%	10%	10%	10%	27%	10%	10%	7%
	HTA-agencies policy makers and others	0%	0%	18%	9%	0%	45%	9%	9%	0%
	Patients	2%	0%	0%	2%	9%	7%	33%	19%	19%
	Health professional	1%	4%	3%	12%	10%	21%	21%	6%	4%
Vaccinations received/ Immunization status	All results	11%	11%	13%	10%	8%	15%	6%	3%	6%
	Researcher within the field	17%	7%	3%	13%	10%	17%	7%	3%	7%
	HTA-agencies policy makers and others	9%	27%	18%	0%	0%	27%	0%	0%	0%
	Patients	12%	5%	16%	7%	9%	19%	7%	2%	12%
	Health professional	7%	13%	15%	12%	7%	9%	6%	3%	3%
Infant or child's temperament	All results	3%	4%	11%	17%	12%	18%	11%	7%	5%
	Researcher within the field	0%	7%	17%	13%	17%	7%	10%	10%	7%
	HTA-agencies policy makers and others	18%	0%	9%	18%	9%	9%	18%	0%	9%
	Patients	2%	0%	5%	14%	16%	28%	9%	7%	9%
	Health professional	1%	6%	12%	19%	7%	18%	12%	7%	1%
Biological parameters in the infant	All results	5%	3%	11%	16%	9%	9%	9%	5%	6%
	Researcher within the field	3%	3%	7%	20%	13%	7%	13%	10%	3%
	HTA-agencies policy makers and others	18%	9%	0%	18%	9%	18%	9%	0%	0%

	Patients	2%	0%	9%	12%	9%	9%	9%	2%	16%
	Health professional	6%	4%	16%	16%	6%	7%	6%	6%	1%
Well-baby check-up status	All results	5%	2%	6%	11%	13%	19%	19%	5%	8%
	Researcher within the field	0%	3%	7%	10%	20%	23%	10%	3%	10%
	HTA-agencies policy makers and others	9%	9%	9%	9%	0%	18%	36%	0%	0%
	Patients	9%	2%	9%	7%	7%	14%	28%	5%	12%
	Health professional	3%	0%	3%	15%	15%	21%	15%	6%	6%
Congenital malformations	All results	2%	4%	7%	11%	9%	15%	12%	11%	12%
	Researcher within the field	0%	7%	3%	13%	13%	20%	7%	10%	10%
	HTA-agencies policy makers and others	9%	0%	18%	9%	0%	9%	27%	9%	0%
	Patients	2%	5%	9%	7%	7%	9%	12%	14%	16%
	Health professional	1%	3%	4%	12%	10%	18%	12%	10%	12%
Infant antibiotic use	All results	11%	7%	20%	10%	9%	11%	3%	2%	5%
	Researcher within the field	17%	3%	13%	20%	10%	10%	3%	3%	3%
	HTA-agencies policy makers and others	36%	0%	18%	0%	9%	27%	0%	0%	0%
	Patients	9%	7%	16%	5%	9%	9%	2%	2%	14%
	Health professional	6%	9%	25%	10%	7%	10%	4%	1%	0%
Birth weight, height and head circumference	All results	3%	4%	9%	15%	11%	23%	11%	3%	5%
	Researcher within the field	3%	0%	3%	10%	13%	30%	10%	10%	7%

	HTA-agencies policy makers and others	0%	27%	9%	27%	18%	9%	0%	0%	0%
	Patients	0%	2%	7%	16%	7%	26%	16%	2%	9%
	Health professional	4%	3%	12%	15%	12%	19%	10%	1%	1%
Small for gestational age (SGA)	All results	2%	6%	10%	12%	11%	20%	7%	9%	5%
	Researcher within the field	0%	7%	13%	10%	10%	20%	7%	17%	3%
	HTA-agencies policy makers and others	0%	9%	0%	0%	0%	73%	0%	0%	0%
	Patients	7%	0%	9%	9%	9%	21%	9%	7%	9%
	Health professional	0%	9%	10%	16%	13%	10%	7%	7%	4%
Apgar score	All results	3%	4%	11%	13%	11%	18%	12%	3%	7%
	Researcher within the field	0%	3%	13%	7%	13%	17%	17%	0%	10%
	HTA-agencies policy makers and others	0%	9%	18%	18%	0%	27%	9%	0%	0%
	Patients	5%	2%	7%	7%	16%	7%	19%	5%	14%
	Health professional	4%	4%	12%	18%	7%	24%	6%	3%	3%
Neonatal intensive care unit admission	All results	1%	0%	5%	15%	8%	20%	12%	9%	15%
	Researcher within the field	0%	0%	0%	13%	10%	30%	10%	10%	13%
	HTA-agencies policy makers and others	0%	0%	18%	18%	0%	36%	9%	0%	0%
	Patients	0%	0%	2%	14%	12%	14%	7%	9%	23%
	Health professional	1%	0%	6%	16%	6%	16%	16%	9%	12%

Outcomes relating to resource use




The result presented in percentage, both in total as well as by the different perspectives. The most common score for each group is highlighted.

Outcome	Perspective	1	2	3	4	5	6	7	8	9
Cost-effectiveness	All results	5%	3%	8%	10%	8%	21%	13%	11%	9%
	Researcher within the field	0%	7%	3%	7%	3%	17%	17%	27%	10%
	HTA-agencies policy makers and others	0%	0%	0%	9%	9%	36%	27%	0%	9%
	Patients	7%	5%	14%	2%	14%	21%	7%	7%	14%
	Health professional	6%	1%	7%	16%	6%	19%	12%	7%	4%
Use of health services	All results	3%	3%	9%	11%	11%	20%	13%	7%	9%
	Researcher within the field	3%	3%	10%	7%	17%	13%	17%	7%	13%
	HTA-agencies policy makers and others	0%	9%	0%	9%	18%	36%	9%	0%	9%
	Patients	2%	0%	7%	12%	12%	16%	21%	7%	12%

	Health professional	3%	3%	12%	13%	7%	22%	7%	9%	6%
Engagement and retention with health services	All results	3%	3%	11%	12%	14%	17%	14%	4%	10%
	Researcher within the field	0%	7%	10%	7%	23%	13%	17%	7%	7%
	HTA-agencies policy makers and others	9%	9%	0%	9%	9%	27%	18%	0%	9%
	Patients	2%	0%	7%	14%	12%	14%	19%	2%	23%
	Health professional	4%	3%	15%	13%	12%	19%	9%	4%	3%
Sick-leave	All results	3%	3%	10%	13%	11%	17%	15%	5%	12%
	Researcher within the field	3%	7%	10%	10%	17%	7%	17%	3%	17%
	HTA-agencies policy makers and others	0%	0%	9%	18%	0%	36%	9%	0%	9%
	Patients	2%	0%	12%	7%	12%	16%	14%	9%	21%
	Health professional	4%	3%	9%	18%	9%	19%	15%	3%	4%

Outcomes relating to experiences and level of satisfaction




The result presented in percentage, both in total as well as by the different perspectives. The most common score for each group is highlighted.

Outcome	Perspective	1	2	3	4	5	6	7	8	9
Satisfaction with intervention	All results	2%	1%	1%	5%	8%	17%	23%	16%	17%
	Researcher within the field	0%	3%	3%	17%	7%	20%	7%	13%	20%
	HTA-agencies policy makers and others	9%	0%	0%	0%	55%	9%	18%	0%	0%
	Patients	2%	0%	0%	0%	5%	19%	16%	14%	40%
	Health professional	1%	1%	1%	4%	3%	16%	34%	21%	3%
Therapeutic alliance	All results	1%	1%	4%	4%	9%	15%	17%	13%	26%
	Researcher within the field	3%	3%	7%	10%	10%	13%	20%	10%	13%

	HTA-agencies policy makers and others	0%	0%	18%	0%	27%	18%	27%	0%	0%
	Patients	0%	0%	0%	2%	2%	5%	14%	12%	58%
	Health professional	0%	1%	3%	3%	9%	22%	15%	16%	16%
Quality of care	All results	1%	4%	8%	7%	7%	19%	16%	10%	17%
	Researcher within the field	3%	13%	17%	10%	10%	10%	10%	3%	13%
	HTA-agencies policy makers and others	9%	0%	18%	0%	27%	18%	9%	9%	0%
	Patients	0%	0%	0%	7%	2%	16%	16%	16%	37%
	Health professional	0%	3%	7%	7%	6%	25%	19%	9%	9%
Parent experience	All results	1%	2%	5%	7%	5%	17%	17%	13%	22%
	Researcher within the field	0%	7%	7%	10%	3%	27%	10%	3%	13%
	HTA-agencies policy makers and others	0%	9%	0%	18%	36%	18%	0%	9%	0%
	Patients	2%	0%	2%	0%	0%	14%	16%	16%	44%
	Health professional	0%	0%	7%	7%	4%	15%	22%	15%	15%
Pregnancy / delivery experience	All results	0%	1%	2%	9%	11%	13%	17%	12%	23%
	Researcher within the field	0%	7%	3%	20%	13%	7%	20%	7%	13%
	HTA-agencies policy makers and others	0%	0%	0%	18%	36%	9%	9%	9%	0%
	Patients	0%	0%	0%	2%	9%	7%	9%	9%	56%
	Health professional	0%	0%	3%	7%	7%	21%	22%	16%	10%

Other outcomes



The result presented in percentage, both in total as well as by the different perspectives. The most common score for each group is highlighted.

Outcome	Perspective	1	2	3	4	5	6	7	8	9
Biological parameters	All results	5%	5%	8%	13%	11%	11%	11%	3%	9%
	Researcher within the field	3%	7%	13%	17%	10%	10%	10%	7%	7%
	HTA-agencies policy makers and others	18%	9%	0%	9%	9%	9%	18%	9%	0%
	Patients	2%	0%	2%	7%	12%	16%	12%	5%	21%
	Health professional	4%	6%	10%	16%	10%	7%	9%	0%	3%
Maternal vital signs/ physical examination	All results	3%	8%	14%	17%	7%	14%	7%	3%	6%
	Researcher within the field	3%	7%	23%	7%	7%	17%	10%	0%	7%
	HTA-agencies policy makers and others	0%	18%	9%	9%	18%	27%	0%	0%	0%
	Patients	2%	2%	12%	21%	7%	9%	9%	7%	16%

	Health professional	3%	10%	12%	19%	4%	13%	4%	1%	0%
Knowledge about anti-depressant treatment in pregnancy	All results	1%	2%	11%	17%	15%	9%	13%	8%	9%
	Researcher within the field	0%	3%	10%	17%	27%	7%	10%	10%	7%
	HTA-agencies policy makers and others	9%	18%	0%	9%	18%	9%	9%	9%	0%
	Patients	0%	0%	5%	16%	7%	12%	16%	7%	21%
	Health professional	1%	0%	18%	18%	13%	7%	13%	7%	4%
Partner's attendance	All results	1%	2%	5%	11%	10%	20%	23%	8%	9%
	Researcher within the field	0%	7%	10%	13%	3%	20%	27%	3%	7%
	HTA-agencies policy makers and others	9%	0%	9%	18%	18%	18%	9%	9%	0%
	Patients	0%	2%	2%	7%	9%	9%	33%	9%	21%
	Health professional	0%	0%	4%	12%	12%	27%	18%	9%	4%
Coping strategies among relatives	All results	1%	5%	12%	9%	15%	18%	12%	3%	9%
	Researcher within the field	0%	17%	17%	17%	17%	10%	0%	7%	3%
	HTA-agencies policy makers and others	9%	0%	36%	9%	9%	9%	9%	0%	9%
	Patients	0%	0%	5%	5%	7%	19%	21%	2%	26%
	Health professional	1%	3%	10%	7%	21%	22%	12%	3%	0%