



Bilaga 4 Sökstrategier för kvalitativa och kvantitativa studier

Appendix 4 Search strategies for qualitative and quantitative articles

Qualitative articles

CINAHL via EBSCO 19 August 2022	sidan 2
Cochrane Library via Wiley 19 August 2022 (CENTRAL)	sidan 4
Embase via Elsevier 19 August 2022	sidan 5
Medline via OvidSP 19 August 2022	sidan 6
PsycINFO via EBSCO 19 August 2022	sidan 8
Scopus via Elsevier 11 March 2021	sidan 8
Cited articles	sidan 9

Quantitative articles

CINAHL via EBSCO 15 August 2022	sid 12
Cochrane Library via Wiley 16 August 2022 (CENTRAL)	sidan 13
Embase via Elsevier 15 August 2022	sidan 14
Medline via OvidSP 15 August 2022	sidan 15
PsycINFO via EBSCO 16 August 2022	sidan 16
Scopus via Elsevier 2 February 2022	sidan 17
Cited articles	sidan 17

KVALITATIVA

Qualitative Strategies

CINAHL via EBSCO 19 August 2022

Title: Perinatal death – experiences in parents, siblings and health personnel

Search terms	Items found
Population: Perinatal death	
1. (MH "Perinatal Death")	9,008
2. TI (stillbirth* or "still birth*" or stillborn or "still born") OR AB (stillbirth* or "still birth*" or stillborn or "still born") OR SU (stillbirth* or "still birth*" or stillborn or "still born")	5,898
3. TI ((antenatal or "ante natal" or antepartal or "ante partal" or perinatal* or "perinatal" or peripartal or "peri partal" or fetal or foetal or intrauterine or "in utero") AND (bereave* or death or grief or griev* or demise or loss or mortality)) OR "pregnancy loss") OR AB ((antenatal or "ante natal" or antepartal or "ante partal" or perinatal* or "perinatal" or peripartal or "peri partal" or fetal or foetal or intrauterine or "in utero") N3 (bereave* or death or grief or griev* or demise or loss or mortality)) OR "pregnancy loss") OR SU ((antenatal or "ante natal" or antepartal or "ante partal" or perinatal* or "perinatal" or peripartal or "peri partal" or fetal or foetal or intrauterine or "in utero") N3 (bereave* or death or grief or griev* or demise or loss or mortality)) OR "pregnancy loss")	16,010
4. 1 OR 2 OR 3	18,315
Evaluation: Experiences, expectations and attitudes in parents and siblings of stillborn children	
5. (MH "Parents+/PF") OR (MH "Mothers+/PF") OR (MH "Fathers+/PF") OR (MH "Single Parent/PF") OR (MH "Parents of Disabled Children/PF") OR (MH "Siblings/PF") OR MH "Family Attitudes") OR (MH "Parental Attitudes") OR (MH "Maternal Attitudes") OR (MH "Paternal Attitudes") OR (MH "Attitude") OR (MH "Attitude to Health") OR (MH "Patient Satisfaction") OR (MH "Patient Preference") OR (MH "Personal Satisfaction") OR (MH "Attitude to Medical Treatment") OR (MH "Health Beliefs") OR (MH "Patient Attitudes") OR (MH "Consumer Satisfaction") OR (MH "Emotions") OR (MH "Patients/PF")	303,994
6. TI ((patient* or women or men or mother* or father* or brother* or sister* or sibling* or parent* or famil* or mum* or dad* or patern* or matern*) N15 (accept* or attitude* or belief* or believ* or disbelief* or disbeliev* or dissatif* or embodied or emotion* or encounter* or expectation* or experience* or improvement* or interpret* or involvement* or meaning or percept* or perspective* or psycholog* or satisf* or stigma* or relation or "self-report" or "treatment barrier*" or trust* or understanding* or unsatisf* or value or view*)) OR AB ((patient* or women or men or mother* or father* or brother* or sister* or sibling* or parent* or famil* or mum* or dad* or patern* or matern*) N15 (accept* or attitude* or belief* or believ* or disbelief* or disbeliev* or dissatif* or embodied or emotion* or encounter* or expectation* or experience* or improvement* or interpret* or involvement* or meaning or percept* or perspective* or psycholog* or satisf* or stigma* or relation or "self-report" or "treatment barrier*" or trust* or understanding* or unsatisf* or value or view*)) OR SU ((patient* or women or men or mother* or father* or brother* or sister* or sibling* or parent* or famil* or mum* or dad* or patern* or matern*) N15 (accept* or attitude* or belief* or believ* or disbelief* or disbeliev* or dissatif* or embodied or emotion* or encounter* or expectation* or experience* or improvement* or interpret* or involvement* or meaning or percept* or perspective* or psycholog* or satisf* or stigma* or relation or "self-report" or "treatment barrier*" or trust* or understanding* or unsatisf* or value or view*))	849,619
7. 5 OR 6	954,332
Evaluation: Experiences, expectations and attitudes in health personnel	
8. (MH "Attitude of Health Personnel+") OR (MH "Health Personnel+/PF")	155,696
9. TI (("childbirth professional*" or counselor* or counsellor* or "general practitioner*" or gynaecologist* or "health car*" or "maternity ward*" or midwi* or nurs* or obstetrician* or physician* or professional* or psychotherapist* or staff) N15 (accept* or attitude* or belief* or believ* or disbelief* or disbeliev* or dissatif* or embodied or emotion* or encounter* or expectation* or experience* or improvement* or interpret* or involvement*	426,689

	or meaning or percept* or perspective* or psycholog* or satisf* or stigma* or relation or "self-report" or "treatment barrier*" or trust* or understanding* or unsatisf* or value or view*) OR AB (("childbirth professional*" or counselor* or counsellor* or "general practitioner*" or gynaecologist* or "health car*" or "maternity ward*" or midwi* or nurs* or obstetrician* or physician* or professional* or psychotherapist* or staff) N15 (accept* or attitude* or belief* or believ* or disbelief* or disbeliev* or dissatif* or embodied or emotion* or encounter* or expectation* or experience* or improvement* or interpret* or involvement* or meaning or percept* or perspective* or psycholog* or satisf* or stigma* or relation or "self-report" or "treatment barrier*" or trust* or understanding* or unsatisf* or value or view*)) OR SU (("childbirth professional*" or counselor* or counsellor* or "general practitioner*" or gynaecologist* or "health car*" or "maternity ward*" or midwi* or nurs* or obstetrician* or physician* or professional* or psychotherapist* or staff) N15 (accept* or attitude* or belief* or believ* or disbelief* or disbeliev* or dissatif* or embodied or emotion* or encounter* or expectation* or experience* or improvement* or interpret* or involvement* or meaning or percept* or perspective* or psycholog* or satisf* or stigma* or relation or "self-report" or "treatment barrier*" or trust* or understanding* or unsatisf* or value or view*))	
10.	8 OR 9	489,159
Study types: studies with qualitative methods		
11.	(MH "Qualitative Studies+") OR (MH "Empirical Research") OR (MH "Questionnaires+") OR (MH "Videorecording") OR (MH "Research, Nursing") OR (MH "Research, Midwifery") OR (MH "Survey Research") OR (MH "Phenomenological Research") OR (MH "Grounded Theory") OR (MH "Multimethod Studies") OR (MH "Content Analysis") OR (MH "Constant Comparative Method") OR (MH "Discourse Analysis") OR (MH "Thematic Analysis") OR (MH "Audio recording") OR (MH "Focus Groups") OR (MH "Interviews+") OR (MH "Narratives+") OR (MH "Self Report+") OR (MH "Surveys+") OR (MH "Field Studies") OR (MH "Cluster Sample+") OR (MH "Comparative Studies") OR (MH "Life Experiences") OR (MH "Ethnology") OR (MH "Ethnonursing Research") OR (MH "Ethnological Research") OR (MH "Ethnographic Research") OR (MH "Phenomenology") OR (MH "Triangulation") OR (MH "Storytelling")	1,384,592
12.	TI (("action research" or "cluster sample*" or "constant comparative method*" or "key informant*" or "maximum variation sampling*" or qualitative* or interview* or "focus group*" or phenomeno* or ethnolog* or ethnograph* or ethnonurs* or ethnomethodolog* or "meta-ethnograph*" or hermeneutic* or "grounded theory" or observation or "lived experience*" or narrat* or "mixed method*" or "content analys*" or "discourse analys*" or poststructur* or "purposive sample*" or "social systems theor*" or "thematic analys*" or "theoretical sample*" or "triangulation design")) OR AB (("action research" or "cluster sample*" or "constant comparative method*" or "key informant*" or "maximum variation sampling*" or qualitative* or interview* or "focus group*" or phenomeno* or ethnolog* or ethnograph* or ethnonurs* or ethnomethodolog* or "meta-ethnograph*" or hermeneutic* or "grounded theory" or observation or "lived experience*" or narrat* or "mixed method*" or "content analys*" or "discourse analys*" or poststructur* or "purposive sample*" or "social systems theor*" or "thematic analys*" or "theoretical sample*" or "triangulation design")) OR SU (("action research" or "cluster sample*" or "constant comparative method*" or "key informant*" or "maximum variation sampling*" or qualitative* or interview* or "focus group*" or phenomeno* or ethnolog* or ethnograph* or ethnonurs* or ethnomethodolog* or "meta-ethnograph*" or hermeneutic* or "grounded theory" or observation or "lived experience*" or narrat* or "mixed method*" or "content analys*" or "discourse analys*" or poststructur* or "purposive sample*" or "social systems theor*" or "thematic analys*" or "theoretical sample*" or "triangulation design"))	690,854
13.	TI (((field N2 (research or study or studies or work)) or (grounded N2 (theor* or study or studies or research or analys*)) or ((lived or life) N2 (experience* or story or stories)) or ((video or tape) N2 record*) or (("semi-structured" or semistructured or unstructured or informal or "in-depth" or indepth or "face-to-face" or structured or guide) N3 (interview* or discussion* or open ended or questionnaire*)))) OR AB (((field N2 (research or study or studies or work)) or (grounded N2 (theor* or study or studies or research or analys*)) or ((lived or life) N2 (experience* or story or stories)) or ((video or tape) N2 record*) or (("semi-structured" or semistructured or unstructured or informal or "in-depth" or indepth or "face-to-face" or structured or guide) N3 (interview* or discussion* or open ended or questionnaire*)))) OR SU (((field N2 (research or study or studies or work)) or (grounded	238,044

	N2 (theor* or study or studies or research or analys*) or ((lived or life) N2 (experience* or story or stories)) or ((video or tape) N2 record*) or (("semi-structured" or semistructured or unstructured or informal or "in-depth" or indepth or "face-to-face" or structured or guide) N3 (interview* or discussion* or open ended or questionnaire*)))	
14.	TI ((mixed or multi) W1 (method* or stud*)) OR AB ((mixed or multi) W1 (method* or stud*)) OR SU ((mixed or multi) W1 (method* or stud*))	31,490
15.	11 OR 12 OR 13 OR 14	1,645,000
Combined sets:		
16.	4 AND (7 OR 10 OR 15)	7,441
Final result		
17.	16 Limiters - Language: Danish, English, Norwegian, Swedish	7,298

The final search result, usually found at the end of the documentation, forms the list of abstracts.

AB = Abstract; **AU** = Author; **DE** = Term from the thesaurus; **MH**= Exact Subject Heading from CINAHL Subject Headings; **MM** = Major Concept; **TI** = Title; **TX** = All Text. Performs a keyword search of all the database's searchable fields; **ZC** = Methodology Index; * = Truncation; " " = Citation Marks; searches for an exact phrase; **N** = Near Operator (N) finds the words if they are a maximum of x words apart from one another, regardless of the order in which they appear.; **W** = Within Operator (W) finds the words if they are within x words of one another, in the order in which you entered them.

Cochrane Library via Wiley 19 August 2022 (CENTRAL)

Title: Perinatal death – experiences in parents, siblings and health personel

Search terms	Items found
Population: Perinatal death	
1. MeSH descriptor: [Stillbirth] explode all trees	151
2. MeSH descriptor: [Fetal Death] explode all trees	394
3. (Stillbirth OR "still birth" OR stillborn OR "still born"):ti, ab, kw	1348
4. (((antenatal or "ante natal" or antepartal or "ante partal" or perinatal* or "peri natal" or peripartal or "peri partal" or fetal or foetal or intrauterine or "in utero") NEAR1 (bereave* or death or grief or griev* or demise or loss or mortality)):ti,ab,kw OR ("pregnancy loss"):ti,ab,kw	885
Final result	
5. 1 OR 2 OR 3 OR 4	CENTRAL/ 1794

The final search result, usually found at the end of the documentation, forms the list of abstracts.

:**au** = Author; **MeSH** = Term from the Medline controlled vocabulary, including terms found below this term in the MeSH hierarchy; **this term only** = Does not include terms found below this term in the MeSH hierarchy; **:ti** = Title; **:ab** = Abstract; **:kw** = Keyword; * = Truncation; " " = Citation Marks; searches for an exact phrase; **CDSR** = Cochrane Database of Systematic Review; **Cochrane Protocols** = Protocols of systematic reviews registered in Cochrane Library; **CENTRAL** = Cochrane Central Register of Controlled Trials, "trials"

Embase via Elsevier 19 August 2022

Title: Perinatal death – experiences in parents, siblings and health personnel

Search terms	Items found
Population: Perinatal death	
1. 'stillbirth'/mj OR 'fetus death'/mj	13,940
2. stillbirth:ti OR 'still birth':ti OR stillborn:ti OR 'still born':ti	3,528
3. (antenatal:ti OR 'ante natal':ti OR antepartal:ti OR 'ante partal':ti OR perinatal*:ti OR 'perinatal':ti OR peripartal:ti OR 'peri partal':ti OR fetal:ti OR foetal:ti OR intrauterine:ti OR 'in utero':ti) AND (bereave*:ti OR death:ti OR grief:ti OR griev*:ti OR demise:ti OR loss:ti OR mortality:ti) OR 'pregnancy loss':ti	14,384
4. 1 OR 2 OR 3	25,019
Evaluation: Experiences, expectations and attitudes in parents and siblings of stillborn children	
Evaluation: Experiences, expectations and attitudes in health personnel	
5. 'patient attitude' OR 'patient preference'/exp/mj OR 'patient satisfaction'/de OR 'family attitude'/exp OR 'health personnel attitude'/exp	473,012
6. (patient*:ti OR women:ti OR men:ti OR mother*:ti OR father*:ti OR brother*:ti OR sister*:ti OR sibling*:ti OR parent*:ti OR famil*:ti OR mum*:ti OR dad*:ti OR patern*:ti OR matern*:ti) AND (accept*:ti OR attitude*:ti OR belief*:ti OR believ*:ti OR disbelief*:ti OR disbeliev*:ti OR dissatif*:ti OR embodied:ti OR emotion*:ti OR encounter*:ti OR expectation*:ti OR experience*:ti OR improvement*:ti OR interpret*:ti OR involvement*:ti OR meaning:ti OR percept*:ti OR perspective*:ti OR psycholog*:ti OR satisf*:ti OR stigma*:ti OR relation:ti OR 'self-report':ti OR 'treatment barrier*:ti OR trust*:ti OR understanding*:ti OR unsatisf*:ti OR value:ti OR view*:ti)	319,370
7. ('childbirth professional*:ti OR counselor*:ti OR counsellor*:ti OR 'general practitioner*:ti OR gynaecologist*:ti OR 'health car*:ti OR 'maternity ward*:ti OR midwi*:ti OR nurs*:ti OR obstetrician*:ti OR physician*:ti OR professional*:ti OR psychotherapist*:ti OR staff:ti) AND (accept*:ti OR attitude*:ti OR belief*:ti OR believ*:ti OR disbelief*:ti OR disbeliev*:ti OR dissatif*:ti OR embodied:ti OR emotion*:ti OR encounter*:ti OR expectation*:ti OR experience*:ti OR improvement*:ti OR interpret*:ti OR involvement*:ti OR meaning:ti OR percept*:ti OR perspective*:ti OR psycholog*:ti OR satisf*:ti OR stigma*:ti OR relation:ti OR 'self-report':ti OR 'treatment barrier*:ti OR trust*:ti OR understanding*:ti OR unsatisf*:ti OR value:ti OR view*:ti)	96,072
8. 5 OR 6 OR 7	795,820
Study types: studies with qualitative method	
9. 'qualitative research'/exp OR 'empirical research'/exp OR 'questionnaire'/exp OR 'nursing research'/exp OR 'phenomenology'/exp OR 'survey methodology'/exp OR 'grounded theory'/exp OR 'content analysis'/exp OR 'multimethod study'/exp OR 'constant comparative method'/exp OR 'discourse analysis'/exp OR 'thematic analysis'/exp OR 'self report'/exp OR 'narrative'/exp OR 'interview'/exp OR 'comparative study'/de OR 'field study'/exp OR 'ethnology'/exp OR 'cluster analysis'/exp OR 'storytelling'/exp OR 'personal experience'/exp OR 'recording'/exp	2,591,482
10. 'action research' OR 'cluster sample*' OR 'constant comparative method*' OR 'key informant*' OR 'maximum variation sampling*' OR qualitative* OR interview* OR 'focus group*' OR phenomeno* OR ethnolog* OR ethnograph* OR ethnonurs* OR ethnomethodolog* OR 'meta-ethnograph*' OR hermeneutic* OR 'grounded theory' OR observation OR 'lived experience*' OR narrat* OR 'mixed method*' OR 'content analys*' OR 'discourse analys*' OR poststructur* OR 'purposive sample*' OR 'social systems theor*' OR 'thematic analys*' OR 'theoretical sample*' OR 'triangulation design'	1,660,948
11. (field NEAR/2 (research OR study OR studies OR work)):ti	46,946
12. (('semi-structured' OR semistructured OR unstructured OR informal OR 'in-depth' OR indepth OR 'face-to-face' OR structured OR guide) NEAR/3 (interview* OR discussion* OR 'open ended' OR questionnaire*)):ti	205,082
13. ((video OR tape) NEAR/2 record*):ti	22,120
14. ((lived OR life) NEAR/2 (experience* OR story OR stories)):ti	6,144
15. (grounded NEAR/2 (theor* OR study OR studies OR research OR analys*)):ti	18,829
16. 9 OR 10 OR 11 OR 12 OR 13 OR 14 OR 15	3,787,795

Limits: language, publication type	
17.	((danish)/lim OR [english]/lim OR [norwegian]/lim OR [swedish]/lim)
18.	[preprint]/lim
Combined sets	
19.	4 AND (8 OR 16) 2,662
Final result	
20.	(17 AND 19) NOT 18 2,383

The final search result, usually found at the end of the documentation, forms the list of abstracts.

/de = Term from the EMTREE controlled vocabulary; **/exp** = Includes terms found below this term in the EMTREE hierarchy
/mj = Major Topic; **:ab** = Abstract; **:au** = Author; **:ti** = Article Title; **:ti,ab** = Title or abstract; ***** = Truncation; **'** = Citation
Marks; searches for an exact phrase; **NEAR/n** = Requests terms that are within 'n' words of each other in either direction;
NEXT/n = Requests terms that are within 'n' words of each other in the order specified

Medline via OvidSP 19 August 2022

Title: Perinatal death – experiences in parents, siblings and health personnel

Search terms	Items found
Population: Perinatal death	
1. fetal death/ or stillbirth/	29847
2. (stillbirth* or "still birth*" or stillborn or "still born").ab,ti.	18142
3. (((antenatal or "ante natal" or antepartal or "ante partal" or perinatal* or "peri natal" or peripartal or "peri partal" or fetal or foetal or intrauterine or "in utero") adj3 (bereave* or death or grief or griev* or demise or loss or mortality)) or "pregnancy loss").ab,ti.	37357
4. 1 OR 2 OR 3	65840
Evaluation: Experiences, expectations and attitudes in parents and siblings of stillborn children	
5. exp Family/px [Psychology]	101249
6. Siblings/px [Psychology]	1653
7. Parents/px [Psychology]	31482
8. attitude to health/ or "patient acceptance of health care"/ or patient satisfaction/ or patient preference/	228079
9. Emotions/	79432
10. Expressed Emotion/	1835
11. exp Patients/px [Psychology]	18406
12. ((patient* or women or men or mother* or father* or brother* or sister* or sibling* or parent* or famil* or mum* or dad* or patern* or matern*) adj15 (accept* or attitude* or belief* or believ* or disbelief* or disbeliev* or dissatif* or embodied or emotion* or encounter* or expectation* or experience* or improvement* or interpret* or involvement* or meaning or percept* or perspective* or psycholog* or satisf* or stigma* or relation or "self-report" or "treatment barrier*" or trust* or understanding* or unsatisf* or value or view*))ti,kf.	1686763
13. 5 OR 6 OR 7 OR 8 OR 9 OR 10 OR 11 OR 12	1927682
Evaluation: Experiences, expectations and attitudes in health personnel	
14. exp Health Personnel/px [Psychology]	124551
15. attitude/ or "attitude of health personnel"/	181210
16. (("childbirth professional*" or counselor* or counsellor* or "general practitioner*" or gynaecologist* or "health car*" or "maternity ward*" or midwi* or nurs* or obstetrician* or physician* or professional* or psychotherapist* or staff) adj15 (accept* or attitude* or belief* or believ* or disbelief* or disbeliev* or dissatif* or embodied or emotion* or encounter* or expectation* or experience* or improvement* or interpret* or involvement* or meaning or percept* or perspective* or psycholog* or satisf* or stigma* or relation or "self-report" or "treatment barrier*" or trust* or understanding* or unsatisf* or value or view*))ti,kf.	378127
17. 14 OR 15 OR 16	572592

Study types: studies with qualitative method		
18.	exp empirical research/ or exp qualitative research/	81403
19.	data collection/ or focus groups/ or interviews as topic/ or narration/ or "surveys and questionnaires"/ or health care surveys/ or patient reported outcome measures/ or health surveys/ or patient health questionnaire/ or self report/	825230
20.	Ethnology/	1592
21.	exp Nursing Research/	53394
22.	exp comparative study/	1911467
23.	personal narratives as topic/	353
24.	personal narrative/	6198
25.	exp Video Recording/	44166
26.	("action research" or "cluster sample*" or "constant comparative method*" or "key informant*" or "maximum variation sampling*" or qualitative* or interview* or "focus group*" or phenomeno* or ethnolog* or ethnograph* or ethnouns* or ethnomethodolog* or "meta-ethnograph*" or hermeneutic* or "grounded theory" or observation or "lived experience*" or narrat* or "mixed method*" or "content analys*" or "discourse analys*" or poststructur* or "purposive sample*" or "social systems theor*" or "thematic analys*" or "theoretical sample*" or "triangulation design").ab,kf,ti.	1332965
27.	((field adj3 (research or study or studies or work)) or (grounded adj2 (theor* or study or studies or research or analys*)) or ((lived or life) adj2 (experience* or story or stories)) or ((video or tape) adj2 record*) or (("semi-structured" or semistructured or unstructured or informal or "in-depth" or indepth or "face-to-face" or structured or guide) adj3 (interview* or discussion* or open ended or questionnaire*))).ab,kf,ti.	273263
28.	18 OR 19 OR 20 OR 21 OR 22 OR 23 OR 24 OR 25 OR 26 OR 27	3907067
Limits: human studies		
29.	exp Animals/ NOT exp Humans/	5038073
Combined sets		
30.	4 AND (13 OR 17 OR 28)	9640
Final result		
31.	30 NOT 29 limit to (danish or english or norwegian or swedish)	7834

The final search result, usually found at the end of the documentation, forms the list of abstracts.

.ab. = Abstract; **.ab,ti.** = Abstract or title; **.af.** = All fields; **Exp** = Term from the Medline controlled vocabulary, including terms found below this term in the MeSH hierarchy; **.sh.** = Term from the Medline controlled vocabulary; **.ti.** = Title; **/** = Term from the Medline controlled vocabulary, but does not include terms found below this term in the MeSH hierarchy; ***** = Focus (if found in front of a MeSH-term); *** or \$** = Truncation (if found at the end of a free text term); **.mp** = Text, heading word, subject area node, title; **" "** = Citation Marks; searches for an exact phrase; **AD/n** = Positional operator that lets you retrieve records that contain your terms (in any order) within a specified number (n) of words of each other.

PsycINFO via EBSCO 19 August 2022

Title: Perinatal death – experiences in parents, siblings and health personnel

Search terms	Items found
Population: Perinatal death	
1. (DE "Death and Dying") AND (DE "Pregnancy Outcomes" OR DE "Birth" OR DE "Induced Abortion" OR DE "Obstetrical Complications" OR DE "Spontaneous Abortion")	539
2. TI (stillbirth* or "still birth*" or stillborn or "still born") OR AB (stillbirth* or "still birth*" or stillborn or "still born") OR SU (stillbirth* or "still birth*" or stillborn or "still born")	1,068
3. TI (((antenatal or "ante natal" or antepartal or "ante partal" or perinatal* or "peri natal" or peripartal or "peri partal" or fetal or foetal or intrauterine or "in utero") N1 (bereave* or death or grief or griev* or demise or loss or mortality)) OR "pregnancy loss") OR AB (((antenatal or "ante natal" or antepartal or "ante partal" or perinatal* or "perinatal" or peripartal or "peri partal" or fetal or foetal or intrauterine or "in utero") N1 (bereave* or death or grief or griev* or demise or loss or mortality)) OR "pregnancy loss") OR SU (((antenatal or "ante natal" or antepartal or "ante partal" or perinatal* or "perinatal" or peripartal or "peri partal" or fetal or foetal or intrauterine or "in utero") N1 (bereave* or death or grief or griev* or demise or loss or mortality)) OR "pregnancy loss")	1,834
4. 1 OR 2 OR 3	2,825
Final result	
5. 4 Limiters - Language: Danish, English, Norwegian, Swedish	2,711

The final search result, usually found at the end of the documentation, forms the list of abstracts.

AB = Abstract; **AU** = Author; **DE** = Term from the thesaurus; **MH** = Exact Subject Heading from CINAHL Subject Headings; **MM** = Major Concept; **TI** = Title; **TX** = All Text. Performs a keyword search of all the database's searchable fields; **ZC** = Methodology Index; * = Truncation; " " = Citation Marks; searches for an exact phrase; **N** = Near Operator (N) finds the words if they are a maximum of x words apart from one another, regardless of the order in which they appear.; **W** = Within Operator (W) finds the words if they are within x words of one another, in the order in which you entered them.

Scopus via Elsevier 11 March 2021

Title: Perinatal death – citation search

Search terms	Items found
Cited articles	
1. (DOI ("10.1007/s00404-017-4309-9" OR "10.1016/j.ejogrb.2018.01.007" OR "10.1016/j.ijnurstu.2012.10.014" OR "10.1016/j.jogn.2015.10.010" OR "10.1016/j.jogn.2020.09.156" OR "10.1016/j.midw.2013.10.021" OR "10.1016/j.midw.2018.06.011" OR "10.1016/j.midw.2018.08.010" OR "10.1016/j.midw.2019.06.013" OR "10.1016/j.wombi.2019.05.013" OR "10.1046/j.1523-536x.2001.00124.x")) OR (DOI ("10.1080/02646838.2012.661849" OR "10.1080/02682621.2020.1828724" OR "10.1080/07481187.2013.809034" OR "10.1080/07481187.2019.1626936" OR "10.1097/ANC.0000000000000703" OR "10.1097/NMC.0000000000000490" OR "10.1111/1471-0528.12695" OR "10.1111/1471-0528.16113" OR "10.1111/1552-6909.12040" OR "10.1111/ajo.12684" OR "10.1111/aogs.13354" OR "10.1111/birt.12335" OR "10.1111/j.0730-7659.2004.00291.x")) OR (DOI ("10.1111/j.0730-7659.2004.00304.x" OR "10.1111/j.1523-536x.2010.00457.x" OR "10.1111/j.1552-6909.1997.tb01506.x" OR "10.1111/scs.12816" OR "10.1136/bmjopen-2012-002237" OR "10.1186/1471-2393-12-137" OR "10.1186/s12884-020-02865-4" OR "10.1371/journal.pone.0191635" OR "10.2190/om.57.1.a" OR "10.12707/RV20037" OR "10.12968/bjom.2016.24.10.715" OR "10.20467/1091-5710.14.4.14" OR "10.1016/j.srhc.2011.02.002" OR "10.1177/00302228211050500" OR "10.1186/1471-2393-14-203")) OR (PMID (24968620 OR 17964235)) OR ((TITLE ("helping a woman") AND AUTHOR-NAME (radestad)))	40
2. View cited by	565

The final search result, usually found at the end of the documentation, forms the list of abstracts.

TITLE-ABS-KEY = Title or abstract or keywords; **ALL** = All fields; **PRE/n** = "precedes by". The first term in the search must precede the second by a specified number of terms (n).; **W/n** = "Within". The terms in the search must be within a specified number of terms (n) in any order.; * = Truncation; " " = Citation Marks; searches for an exact phrase; **LIMIT-TO (SRCTYPE, "j")** = Limit to source type journal; **LIMIT-TO (DOCTYPE, "ar")** = Limit to document type article; **LIMIT-TO (DOCTYPE, "re")** = Limit to document type review

Cited articles

1. Agwu Kalu F, Coughlan B, Larkin P. A mixed methods sequential explanatory study of the psychosocial factors that impact on midwives' confidence to provide bereavement support to parents who have experienced a perinatal loss. *Midwifery*. 2018;64:69-76.
2. Alghamdi R, Jarrett P. Experiences of student midwives in the care of women with perinatal loss: A qualitative descriptive study. *British Journal of Midwifery*. 2016;24(10):715-22.
3. Avelin P, Erlandsson K, Hildingsson I, Rådestad I. Swedish parents' experiences of parenthood and the need for support to siblings when a baby is stillborn. *Birth*. 2011;38(2):150-8.
4. Avelin P, Gyllenswärd G, Erlandsson K, Rådestad I. Adolescents' Experiences of Having a Stillborn Half-Sibling. *Death Studies*. 2014;38(9):557-62.
5. Bond D, Raynes-Greenow C, Gordon A. Bereaved parents' experience of care and follow-up after stillbirth in Sydney hospitals. *Australian and New Zealand Journal of Obstetrics and Gynaecology*. 2018;58(2):185-91.
6. Brierley-Jones L, Crawley R, Jones E, Gordon I, Knight J, Hinshaw K. Supporting parents through stillbirth: A qualitative study exploring the views of health professionals and health care staff in three hospitals in England. *European Journal of Obstetrics and Gynecology and Reproductive Biology*. 2018;222:45-51.
7. Cacciatore J, Erlandsson K, Rådestad I. Fatherhood and suffering: A qualitative exploration of Swedish men's experiences of care after the death of a baby. *International Journal of Nursing Studies*. 2013;50(5):664-70.
8. Camacho Ávila M, Fernández Medina IM, Jiménez-López FR, Granero-Molina J, Hernández-Padilla JM, Hernández Sánchez E, et al. Parents' Experiences about Support Following Stillbirth and Neonatal Death. *Adv Neonat Care*. 2020;20(2):151-60.
9. Cassidy PR. The Disenfranchisement of Perinatal Grief: How Silence, Silencing and Self-Censorship Complicate Bereavement (a Mixed Methods Study). *Omega (United States)*. 2021.
10. Downe S, Schmidt E, Kingdon C, Heazell AEP. Bereaved parents' experience of stillbirth in UK hospitals: A qualitative interview study. *BMJ Open*. 2013;3(2).
11. Fernández-Alcántara M, Schul-Martin L, García Caro MP, Montoya-Juárez R, Pérez-Marfil MN, Zech E. 'In the hospital there are no care guidelines': experiences and practices in perinatal loss in Spain. *Scandinavian Journal of Caring Sciences*. 2020;34(4):1063-73.
12. Gillis C, Wheatley V, Jones A, Roland B, Gill M, Marlett N, et al. Stillbirth, still life: A qualitative patient-led study on parents' unsilenced stories of stillbirth. *Bereavement Care*. 2020;39(3):124-32.
13. Hutti MH, Polivka B, White S, Hill J, Clark P, Cooke C, et al. Experiences of Nurses Who Care for Women After Fetal Loss. *JOGNN - Journal of Obstetric, Gynecologic, and Neonatal Nursing*. 2016;45(1):17-27.
14. Kavanaugh K. Parents' experience surrounding the death of a newborn whose birth is at the margin of viability. *Journal of obstetric, gynecologic, and neonatal nursing : JOGNN / NAACOG*. 1997;26(1):43-51.
15. Kelley MC, Trinidad SB. Silent loss and the clinical encounter: Parents' and physicians' experiences of stillbirth-a qualitative analysis. *BMC Pregnancy and Childbirth*. 2012;12.
16. King MQ, Oka M, Robinson WD. Pain without reward: A phenomenological exploration of stillbirth for couples and their hospital encounter. *Death Studies*. 2021;45(2):152-62.
17. Lee C. 'She was a person, she was here': The experience of late pregnancy loss in Australia. *Journal of Reproductive and Infant Psychology*. 2012;30(1):62-76.

18. Listermar KH, Sormunen T, Rådestad I. Perinatal palliative care after a stillbirth—Midwives' experiences of using Cubitus baby. *Women and Birth*. 2020;33(2):161-4.
19. Malm MC, Rådestad I, Erlandsson K, Lindgren H. Waiting in no-man's-land - Mothers' experiences before the induction of labour after their baby has died in utero. *Sexual and Reproductive Healthcare*. 2011;2(2):51-5.
20. Martínez-Serrano P, Palmar-Santos AM, Solís-Muñoz M, Álvarez-Plaza C, Pedraz-Marcos A. Midwives' experience of delivery care in late foetal death: A qualitative study. *Midwifery*. 2018;66:127-33.
21. Martínez-Serrano P, Pedraz-Marcos A, Solís-Muñoz M, Palmar-Santos AM. The experience of mothers and fathers in cases of stillbirth in Spain. A qualitative study. *Midwifery*. 2019;77:37-44.
22. McCreight BS. Perinatal loss: A qualitative study in Northern Ireland. *Omega: Journal of Death and Dying*. 2008;57(1):1-19.
23. McNamara K, Meaney S, O'Donoghue K. Intrapartum fetal death and doctors: a qualitative exploration. *Acta Obstetrica et Gynecologica Scandinavica*. 2018;97(7):890-8.
24. McNamara K, Meaney S, O'Connell O, McCarthy M, Greene RA, O'Donoghue K. Healthcare professionals' response to intrapartum death: a cross-sectional study. *Archives of Gynecology and Obstetrics*. 2017;295(4):845-52.
25. Miranda AMC, Zangão MOB. Mothers' experiences of fetal death. *Revista de Enfermagem Referencia*. 2020;2020(3):1-8.
26. Nurse-Clarke N. Managing Ambiguity When Caring for Women Who Experience Stillbirth. *JOGNN - Journal of Obstetric, Gynecologic, and Neonatal Nursing*. 2021;50(2):143-53.
27. Nuzum D, Meaney S, O'Donoghue K. The impact of stillbirth on consultant obstetrician gynaecologists: A qualitative study. *BJOG: An International Journal of Obstetrics and Gynaecology*. 2014;121(8):1020-8.
28. Nuzum D, Meaney S, O'Donoghue K. The impact of stillbirth on bereaved parents: A qualitative study. *PLoS ONE*. 2018;13(1).
29. Puia DM, Lewis L, Beck CT. Experiences of obstetric nurses who are present for a perinatal loss. *JOGNN - Journal of Obstetric, Gynecologic, and Neonatal Nursing*. 2013;42(3):321-31.
30. Redshaw M, Henderson J. Care associated with stillbirth for the most disadvantaged women: A multi-method study of care in England. *Birth*. 2018;45(3):275-85.
31. Ryninks K, Roberts-Collins C, McKenzie-McHarg K, Horsch A. Mothers' experience of their contact with their stillborn infant: An interpretative phenomenological analysis. *BMC Pregnancy and Childbirth*. 2014;14(1).
32. Rådestad I, Christoffersen L. Helping a woman meet her stillborn baby while it is soft and warm. *British Journal of Midwifery*. 2008;16(9):588-91.
33. Rådestad I, Malm MC, Lindgren H, Pettersson K, Larsson LLF. Being alone in silence - Mothers' experiences upon confirmation of their baby's death in utero. *Midwifery*. 2014;30(3):e91-e5.
34. Samuelsson M, Rådestad I, Segesten K. A waste of life: Fathers' experience of losing a child before birth. *Birth*. 2001;28(2):124-30.
35. Skupski D. We need bereavement training and support for stillbirth providers. *BJOG: An International Journal of Obstetrics and Gynaecology*. 2014;121(8):1028.
36. Smith LK, Dickens J, Bender Atik R, Bevan C, Fisher J, Hinton L. Parents' experiences of care following the loss of a baby at the margins between miscarriage, stillbirth and neonatal death: a UK qualitative study. *BJOG: An International Journal of Obstetrics and Gynaecology*. 2020;127(7):868-74.
37. Smith P, Vasileiou K, Jordan A. Healthcare professionals' perceptions and experiences of using a cold cot following the loss of a baby: A qualitative study in maternity and neonatal units in the UK. *BMC Pregnancy and Childbirth*. 2020;20(1).

38. Säflund K, Sjögren B, Wredling R. The role of caregivers after a stillbirth: Views and experiences of parents. *Birth*. 2004;31(2):132-7.

39. Trulsson O, Rådestad I. The silent child - Mothers' experiences before, during, and after stillbirth. *Birth*. 2004;31(3):189-95.

40. Willis P. Nurses' Perspective on Caring for Women Experiencing Perinatal Loss. *MCN The American Journal of Maternal/Child Nursing*. 2019;44(1):46-51.

KVANTITATIVA

Quantitative Strategies

CINAHL via EBSCO 15 August 2022

Title: Perinatal death - support

Search terms	Items found
Population: Perinatal death	
1. (MH "Perinatal Death")	8,999
2. TI ((stillbirth* or "still birth*" or stillborn or "still born") OR AB ((stillbirth* or "still birth*" or stillborn or "still born"))	5,889
3. TI (((antenatal or "ante natal" or antepartal or "ante partal" or perinatal* or "perinatal" or peripartal or "peri partal" or fetal or foetal or intrauterine or "in utero") AND (bereave* or death or grief or griev* or demise or mortality) or "pregnancy loss") OR AB (((antenatal or "ante natal" or antepartal or "ante partal" or perinatal* or "perinatal" or peripartal or "peri partal" or fetal or foetal or intrauterine or "in utero") N3 (bereave* or death or grief or griev* or demise or mortality) or "pregnancy loss")	10,709
4. <i>S1 OR S2 OR S3</i>	19,136
Population: Siblings of stillborn children	
5. (MH "Siblings")	6,530
6. TI (brother* or sister* or sibling*) OR AB (brother* or sister* or sibling*) OR SU (brother* or sister* or sibling*)	20,131
7. <i>S5 OR S6</i>	20,131
Intervention: Support	
8. MH "Bereavement Support (Saba CCC)" OR MH "Psychosocial Care (Saba CCC)" OR MH "Psychotherapy+" OR MH "Midwifery+" OR MH "Grief Work Facilitation (Iowa NIC)" OR (MH "Support, Psychosocial") OR (MH "Counseling") OR (MH "Mental Health Services+")	312,586
9. TI (abreaction or "acceptance and commitment" or ACT or "applied behav*" or care or caring or CBT or consultation* or counsel* or "crisis intervention*" or DBT or "emotion focused" or "free association" or hypnosis or hypnotherapy or intervent* or IPT or "life review" or "listening visit*" or logotherap* or mindfulness or "mom power" or "motivational interview*" or "peer group" or "peer support" or "problem solv*" or program* or psychoanaly* or "psycho-analy*" or psychodynamic* or "psycho dynamic*" or psychodrama or "psycho-drama*" or psychoeducat* or "psychologic desenzitization" or "psychological feedback" or psychosocial or psychotherap* or "psycho therap*" or "rational emotive" or "reminiscence" or "self control" or "self management" or "self system" or service* or "socioenvironmental therapy" or support* or therap* or "transactional analysis" or train* or treat* or holding or "cold cot")	1,460,767
10. TI ((behav* N3 (activat* or component* or contracting or defusion or guidance or management or modif* or rehab* or restructur* or technique* or train*))) OR AB ((behav* N3 (activat* or component* or contracting or defusion or guidance or management or modif* or rehab* or restructur* or technique* or train*))) OR SU ((behav* N3 (activat* or component* or contracting or defusion or guidance or management or modif* or rehab* or restructur* or technique* or train*)))	22,043
11. TI (((cognitiv* or dialect*) N3 (behav* or component* or contracting or defusion or guidance or management or modif* or rehab* or restructur* or technique* or train*))) OR AU (((cognitiv* or dialect*) N3 (behav* or component* or contracting or defusion or guidance or management or modif* or rehab* or restructur* or technique* or train*))) OR SU (((cognitiv* or dialect*) N3 (behav* or component* or contracting or defusion or guidance or management or modif* or rehab* or restructur* or technique* or train*)))	33,666
12. TI (((home or house) N3 (based or call* or care or service* or visit*))) OR AB (((home or house) N3 (based or call* or care or service* or visit*))) OR SU (((home or house) N3 (based or call* or care or service* or visit*)))	76,180
13. <i>S8 OR S9 OR S10 OR S11 OR S12</i>	1,669,572

Outcome: Bereavement		
14.	(MH "Bereavement+") OR (MH "Grief+") OR (MH "Psychological Distress") OR (MH "Depression, Reactive") OR (MM "Adolescent Parents+/PF") OR (MM "Mothers+/PF") OR (MM "Fathers+/PF") OR (MM "Parents+/PF") OR (MM "Single Parent/PF") OR (MH "Expectant Parents+/PF") OR (MH "Grieving (Saba CCC)+") OR (MH "Grieving (NANDA)+") OR (MH "Grief Resolution (Iowa NOC)") OR (MH "Grief (Omaha)")	44,621
15.	TI (bereav* or grief* or griev* or mourn* or sorrow* or sadness)	9,375
16.	(MH "Perinatal Death/PF")	888
17.	S14 OR S15	46,630
Combined sets:		
18.	S4 AND S7	150
19.	S4 AND S13	3,668
20.	(S4 AND S17)	1,667
21.	S16 OR S18 OR S19 OR S20	4,691
Final result		
22.	S21 Limiters - Language: Danish, English, Norwegian, Swedish	4,585

The final search result, usually found at the end of the documentation, forms the list of abstracts.

AB = Abstract; **AU** = Author; **DE** = Term from the thesaurus; **MH** = Exact Subject Heading from CINAHL Subject Headings; **MM** = Major Concept; **TI** = Title; **TX** = All Text. Performs a keyword search of all the database's searchable fields; **ZC** = Methodology Index; ***** = Truncation; **" "** = Citation Marks; searches for an exact phrase; **N** = Near Operator (N) finds the words if they are a maximum of x words apart from one another, regardless of the order in which they appear.; **W** = Within Operator (W) finds the words if they are within x words of one another, in the order in which you entered them.

Cochrane Library via Wiley 16 August 2022 (CENTRAL)

Title: Perinatal death - support

Search terms	Items found
Population: Perinatal death	
1. MeSH descriptor: [Stillbirth] explode all trees	151
2. MeSH descriptor: [Fetal Death] explode all trees	394
3. (Stillbirth OR "still birth" OR stillborn OR "still born"):ti, ab, kw	1348
4. (((antenatal or "ante natal" or antepartal or "ante partal" or perinatal* or "peri natal" or peripartal or "peri partal" or fetal or foetal or intrauterine or "in utero") NEAR1 (bereave* or death or grief or griev* or demise or loss or mortality)):ti,ab,kw OR ("pregnancy loss"):ti,ab,kw	885
Final result	
5. 1 OR 2 OR 3 OR 4	CENTRAL/ 1794

The final search result, usually found at the end of the documentation, forms the list of abstracts.

:au = Author; **MeSH** = Term from the Medline controlled vocabulary, including terms found below this term in the MeSH hierarchy; **this term only** = Does not include terms found below this term in the MeSH hierarchy; **:ti** = Title; **:ab** = Abstract; **:kw** = Keyword; ***** = Truncation; **" "** = Citation Marks; searches for an exact phrase; **CDSR** = Cochrane Database of Systematic Review; **Cochrane Protocols** = Protocols of systematic reviews registered in Cochrane Library; **CENTRAL** = Cochrane Central Register of Controlled Trials, "trials"

Embase via Elsevier 15 August 2022

Title: Perinatal death - support

Search terms	Items found
Population: Perinatal death	
1. 'stillbirth'/mj OR 'fetus death'/mj	13,937
2. stillbirth:ti OR 'still birth':ti OR stillborn:ti OR 'still born':ti	3,538
3. (antenatal:ti OR 'ante natal':ti OR antepartal:ti OR 'ante partal':ti OR perinatal*:ti OR 'perinatal':ti OR peripartal:ti OR 'peri partal':ti OR fetal:ti OR foetal:ti OR intrauterine:ti OR 'in utero':ti) AND (bereave*:ti OR death:ti OR grief:ti OR griev*:ti OR demise:ti OR loss:ti OR mortality:ti) OR 'pregnancy loss':ti	14,382
4. 1 OR 2 OR 3	24,749
Population: Siblings of stillborn children	
5. 'sibling'/exp	51,319
6. brother*:ti OR sister*:ti OR sibling*:ti	27,906
7. 5 OR 6	67,476
Intervention: Support	
8. 'psychotherapy'/exp OR 'counseling'/exp OR 'home care'/exp OR 'peer group'/exp OR 'social support'/exp OR 'self care'/exp OR 'self help'/exp OR 'nursing'/exp	1,124,569
9. abreaction:ti OR 'acceptance and commitment':ti OR act:ti OR 'applied behav*':ti OR care:ti OR cbt:ti OR consultation*:ti OR counsel*:ti OR 'crisis intervention':ti OR dbt:ti OR 'emotion focused':ti OR 'free association':ti OR hypnosis:ti OR hypnotherapy:ti OR intervent*:ti OR ipt:ti OR 'life review':ti OR 'listening visit*':ti OR logotherap*:ti OR mindfulness:ti OR 'mom power':ti OR 'motivational interview*':ti OR 'peer group':ti OR 'peer support':ti OR 'problem solv*':ti OR program*:ti OR psychoanaly*:ti OR 'psychoanaly*':ti OR psychodynamic*:ti OR 'psycho dynamic*':ti OR psychodrama:ti OR 'psycho-drama*':ti OR psychoeducat*:ti OR 'psychologic desenzitization':ti OR 'psychological feedback':ti OR psychosocial:ti OR psychotherap*:ti OR 'psycho therap*':ti OR 'rational emotive':ti OR 'reminiscence':ti OR 'self control':ti OR 'self management':ti OR 'self system':ti OR service*:ti OR 'socioenvironmental therapy':ti OR support*:ti OR therap*:ti OR 'transactional analysis':ti OR train*:ti OR treat*:ti OR holding:ti OR 'cold cot':ti	4,918,168
10. (behav* NEAR/3 (activat* OR component* OR contracting OR defusion OR guidance OR management OR modif* OR rehab* OR restructur* OR technique* OR train*)):ti,ab	56,732
11. ((cognitiv* or dialect*) adj3 (behav* or component? or contracting or defusion or guidance or management or modif* or rehab* or restructur* or technique* or train*)):ab,ti,kf.	91,917
12. ((home OR house) NEAR/3 (based OR call* OR care OR service* OR visit*)):ti,ab	82,097
13. 8 OR 9 OR 10 OR 11 OR 12	5,680,816
Outcome: Bereavement	
14. 'bereavement'/exp OR 'grief'/exp OR 'reactive depression'/exp	22,610
15. griev*:ti OR grief*:ti OR sadness:ti OR sorrow:ti OR bereav*:ti OR mourn*:ti	10,826
16. 14 OR 15	24,817
Combined sets	
17. 4 AND (7 OR 13 OR 16)	3,203
Final result	
18. 17 AND ((danish)/lim OR [english]/lim OR [norwegian]/lim OR [swedish]/lim)	2,855

The final search result, usually found at the end of the documentation, forms the list of abstracts.

/de = Term from the EMTREE controlled vocabulary; **/exp** = Includes terms found below this term in the EMTREE hierarchy
/mj = Major Topic; **:ab** = Abstract; **:au** = Author; **:ti** = Article Title; **:ti,ab** = Title or abstract; ***** = Truncation; **' '** = Citation Marks; searches for an exact phrase; **NEAR/n** = Requests terms that are within 'n' words of each other in either direction; **NEXT/n** = Requests terms that are within 'n' words of each other in the order specified

Medline via OvidSP 15 August 2022

Title: Perinatal death - support

Search terms	Items found
Population: Perinatal death	
1. fetal death/ or stillbirth/	29843
2. (stillbirth* or "still birth*" or stillborn or "still born").ti.	3616
3. (((antenatal or "ante natal" or antepartal or "ante partal" or perinatal* or "peri natal" or peripartal or "peri partal" or fetal or foetal or intrauterine or "in utero") adj1 (bereave* or death or grief or griev* or demise or loss or mortality)) or "pregnancy loss").ti.	9456
4. 1 OR 2 OR 3 OR 4	35462
Population: Siblings of stillborn children	
5. Siblings/	12879
6. (brother* or sister* or sibling*).ti.	23734
7. 5 OR 6	29601
Intervention: Support	
8. exp Psychotherapy/	213331
9. exp Counseling/	47836
10. Home Care Services/	35763
11. peer group/	23350
12. exp social support/	78108
13. Self Care/	35503
14. Self-Management/	4741
15. Self-Help Groups/	9493
16. nursing/ or midwifery/	71636
17. (abreaction or "acceptance and commitment" or ACT or "applied behav*" or care or CBT or consultation* or counsel* or "crisis intervention" or DBT or "emotion focused" or "free association" or hypnosis or hypnotherapy or intervent* or IPT or "life review" or "listening visit*" or logotherap* or mindfulness or "mom power" or "motivational interview*" or "peer group" or "peer support" or "problem solv*" or program* or psychoanaly* or "psycho-analy*" or psychodynamic* or "psycho dynamic*" or psychodrama or "psycho-drama*" or psychoeducat* or "psychologic desenzitization" or "psychological feedback" or psychosocial or psychotherap* or "psycho therap*" or "rational emotive" or "reminiscence" or "self control" or "self management" or "self system" or service* or "socioenvironmental therapy" or support* or therap* or "transactional analysis" or train* or treat* or holding or "cold cot").ti.	3751874
18. (behav* adj3 (activat* or component? or contracting or defusion or guidance or management or modif* or rehab* or restructur* or technique* or train*)).ab,ti,kf.	45904
19. ((cognitiv* or dialect*) adj3 (behav* or component? or contracting or defusion or guidance or management or modif* or rehab* or restructur* or technique* or train*)).ab,ti,kf.	67733
20. ((home or house) adj3 (based or call* or care or service* or visit*)).ab,ti,kf.	63568
21. 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19 or 20	4081763
Outcome: Bereavement	
22. exp bereavement/ or exp grief/ or exp psychological distress/ or sadness/	21087
23. Adjustment Disorders/	4295
24. parents/ or fathers/ or mothers/ or single parent/ or siblings/ or single-parent family/	144858
25. Parents/px [Psychology]	31466
26. Mothers/px [Psychology]	23210
27. Fathers/px [Psychology]	4019
28. Siblings/px [Psychology]	1653
29. Family/px [Psychology]	23725
30. Stillbirth/px [Psychology]	383
31. (griev* or grief* or sadness or sorrow or bereav* or mourn*).ti.	9258
32. 22 or 23 or 24 or 25 or 26 or 27 or 28 or 29 or 31	189853

Limits: human studies		
33.	exp Animals/ NOT exp Humans/	5036395
Combined sets		
34.	4 and 7	66
35.	4 and 21	2987
36.	4 and 32	1487
Final result		
37.	(30 OR 34 OR 35 OR 36) NOT 33 limit 37 to (danish or english or norwegian or swedish)	3286

The final search result, usually found at the end of the documentation, forms the list of abstracts.

.ab. = Abstract; **.ab,ti.** = Abstract or title; **.af.** = All fields; **Exp** = Term from the Medline controlled vocabulary, including terms found below this term in the MeSH hierarchy; **.sh.** = Term from the Medline controlled vocabulary; **.ti.** = Title; **/** = Term from the Medline controlled vocabulary, but does not include terms found below this term in the MeSH hierarchy; ***** = Focus (if found in front of a MeSH-term); *** or \$** = Truncation (if found at the end of a free text term); **.mp** = Text, heading word, subject area node, title; **" "** = Citation Marks; searches for an exact phrase; **ADJn** = Positional operator that lets you

PsycINFO via EBSCO 16 August 2022

Title: Perinatal death - support

Search terms	Items found
Population: Perinatal death	
1. (DE "Death and Dying") AND (DE "Pregnancy Outcomes" OR DE "Birth" OR DE "Obstetrical Complications" OR DE "Spontaneous Abortion")	537
2. TI (stillbirth* or "still birth*" or stillborn or "still born") OR AB (stillbirth* or "still birth*" or stillborn or "still born") OR SU (stillbirth* or "still birth*" or stillborn or "still born")	1,067
3. TI (((antenatal or "ante natal" or antepartal or "ante partal" or perinatal* or "peri natal" or peripartal or "peri partal" or fetal or foetal or intrauterine or "in utero") N1 (bereave* or death or grief or griev* or demise or loss or mortality)) OR "pregnancy loss") OR AB (((antenatal or "ante natal" or antepartal or "ante partal" or perinatal* or "perinatal" or peripartal or "peri partal" or fetal or foetal or intrauterine or "in utero") N1 (bereave* or death or grief or griev* or demise or loss or mortality)) OR "pregnancy loss") OR SU (((antenatal or "ante natal" or antepartal or "ante partal" or perinatal* or "perinatal" or peripartal or "peri partal" or fetal or foetal or intrauterine or "in utero") N1 (bereave* or death or grief or griev* or demise or loss or mortality)) OR "pregnancy loss")	1,834
4. S1 OR S2 OR S3	2,823
Limits: human studies	
5. DE "Animals" OR DE "Animal Limb" OR DE "Animal Offspring" OR DE "Female Animals" OR DE "Infants (Animal)" OR DE "Invertebrates" OR DE "Male Animals" OR DE "Pets" OR DE "Service Animals" OR DE "Species Differences" OR DE "Vertebrates"	282,605
Final result	
6. S4 NOT S5 Limiters - Language: Danish, English, Norwegian, Swedish	2,552

The final search result, usually found at the end of the documentation, forms the list of abstracts.

AB = Abstract; **AU** = Author; **DE** = Term from the thesaurus; **MH** = Exact Subject Heading from CINAHL Subject Headings; **MM** = Major Concept; **TI** = Title; **TX** = All Text. Performs a keyword search of all the database's searchable fields; **ZC** = Methodology Index; ***** = Truncation; **" "** = Citation Marks; searches for an exact phrase; **N** = Near Operator (N) finds the words if they are a maximum of x words apart from one another, regardless of the order in which they appear.; **W** = Within Operator (W) finds the words if they are within x words of one another, in the order in which you entered them.

Scopus via Elsevier 2 February 2022

Title: Perinatal death – citation search

Search terms	Items found
Cited articles	
1. (DOI (10.1080/07481187.2011.553318) OR DOI (10.1136/bmj.285.6353.1475) OR DOI (10.1016/s0002-9378(87)80295-8) OR DOI (10.1080/074811800200469))	4
2. <i>View cited by</i>	254
3. <i>View references</i>	132
Final result	
4. 2 Refined to: LIMIT-TO (LANGUAGE , "English") OR LIMIT-TO (LANGUAGE , "Norwegian")	239
5. 3	132

The final search result, usually found at the end of the documentation, forms the list of abstracts.

TITLE-ABS-KEY = Title or abstract or keywords; **ALL** = All fields; **PRE/n** = "precedes by". The first term in the search must precede the second by a specified number of terms (n).; **W/n** = "Within". The terms in the search must be within a specified number of terms (n) in any order.; * = Truncation; " " = Citation Marks; searches for an exact phrase; **LIMIT-TO (SRCTYPE, "j"** = Limit to source type journal; **LIMIT-TO (DOCTYPE, "ar"** = Limit to document type article; **LIMIT-TO (DOCTYPE, "re"** = Limit to document type review

Cited articles:

1. Aho AL, Tarkka MT, Åstedt-Kurki P, Sorvari L, Kaunonen M. Evaluating a bereavement follow-up intervention for grieving fathers and their experiences of support after the death of a child-a pilot study. *Death Studies*. 2011;35(10):879-904.
2. Forrest GC, Standish E, Baum JD. Support after perinatal death: A study of support and counselling after perinatal bereavement. *Br Med J*. 1982;285(6353):1475-9.
3. Lake MF, Johnson TM, Murphy J, Knuppel RA. Evaluation of a perinatal grief support team. *American Journal of Obstetrics and Gynecology*. 1987;157(5):1203-6.
4. Murray JA, Terry DJ, Vance JC, Battistutta D, Connolly Y. Effects of a program of intervention on parental distress following infant death. *Death Studies*. 2000;24(4):275-305.