

Bilaga 3B - Exkluderade studier

Studier som efter fulltextgranskning ej uppfyllt inklusionskriterierna och således inte ligger till grund för evidensbaserade resultat.

Kapitel 3.1

Acharya S, Maskarinec G, Williams AE, Oshiro C, Hebshi S, Murphy SP. Nutritional changes among premenopausal women undertaking a soya based dietary intervention study in Hawaii. *Journal of Human Nutrition and Dietetics* 2004;17:413-419.

Acheson KJ. Carbohydrate and weight control: where do we stand? *Curr Opin Clin Nutr Metab Care* 2004;7:485-92.

Agurs-Collins T, Rosenberg L, Makambi K, Palmer JR, Adams-Campbell L. Dietary patterns and breast cancer risk in women participating in the Black Women's Health Study. *Am J Clin Nutr*. 2009;90:621-8.

Akbaraly TN, Ferrie JE, Berr C, Brunner EJ, Head J, Marmot MG, et al. Alternative Healthy Eating Index and mortality over 18 y of follow-up: results from the Whitehall II cohort. *Am J Clin Nutr* 2011;94:247-53.

Akbaraly TN, Singh-Manoux A, Tabak AG, Jokela M, Virtanen M, Ferrie JE, et al. Overall diet history and reversibility of the metabolic syndrome over 5 years: The Whitehall II prospective cohort study. *Diabetes Care* 2010;33:2339-2341.

Alhassan S, Kim S, Bersamin A, King AC, Gardner CD. Dietary adherence and weight loss success among overweight women: results from the A TO Z weight loss study. *Int J Obes (Lond)* 2008;32:985-91.

Alonso A, de la Fuente C, Martin-Arnau AM, de Irala J, Martinez JA, Martinez-Gonzalez MA. Fruit and vegetable consumption is inversely associated with blood pressure in a Mediterranean population with a high vegetable-fat intake: the Seguimiento Universidad de Navarra (SUN) Study. *Br J Nutr* 2004;92:311-9.

Altorf-van der Kuil W, Engberink MF, van Rooij FJ, Hofman A, van't Veer P, Witteman JC, et al. Dietary protein and risk of hypertension in a Dutch older population: the Rotterdam study. *J Hypertens*. 2010;28:2394-400.

Anderson R, Anderson D, Hurst C. Modeling factors that influence exercise and dietary change among midlife Australian women: results from the Healthy Aging of Women Study. *Maturitas*. 2010;67:151-8.

Andrews RC, Cooper AR, Montgomery AA, Norcross AJ, Peters TJ, Sharp DJ, et al. Diet or diet plus physical activity versus usual care in patients with newly diagnosed type 2 diabetes: the Early ACTID randomised controlled trial. *Lancet* 2011;378:129-39.

Anton S, Han H, Bray G, Sacks F, Williamson D. Demographic and macronutrient diet characteristics influence adherence to calorie restricted diets: Findings from the pounds lost trial. *SOURCE Obesity*

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

(2011) 19 SUPPL. 1 (S101). Date of Publication: November 2011 CONFERENCE NAME 29th Annual Scientific Meeting of the Obesity Society, Obesity 2011 CONFERENCE LOCATION Orlando, FL, United States CONFERENCE DATE 2011-10-01 to 2011-10-05.

Armendariz-Anguiano AL, Jimenez-Cruz A, Bacardi-Gascon M, Hurtado-Ayala L. Effect of a low glycemic load on body composition and Homeostasis Model Assessment (HOMA) in overweight and obese subjects. *Nutr Hosp* 2011;26:170-5.

Ascherio A, Rimm EB, Giovannucci EL, Colditz GA, Rosner B, Willett WC, et al. A prospective study of nutritional factors and hypertension among US men. *Circulation*. 1992;86:1475-84.

Ash S, Koutsoukos M, Capra S, Yeo S, Kerr G. Intensive diet therapy improves weight and glycaemic control in overweight men with type II diabetes. *American Journal of Clinical Nutrition* 2002;382s.

Astrup A, Grunwald GK, Melanson EL, Saris WH, Hill JO. The role of low-fat diets in body weight control: a meta-analysis of ad libitum dietary intervention studies. *Int J Obes Relat Metab Disord* 2000;24:1545-52.

Augustsson K, Michaud DS, Rimm EB, Leitzmann MF, Stampfer MJ, Willett WC, et al. A prospective study of intake of fish and marine fatty acids and prostate cancer. *Cancer Epidemiol Biomarkers Prev* 2003;12:64-7.

Azadbakht L, Mirmiran P, Esmailzadeh A, Azizi F. Better dietary adherence and weight maintenance achieved by a long-term moderate-fat diet. *Br J Nutr* 2007;97:399-404.

Bahadori B, Yazdani-Biuki B, Krippel P, Brath H, Uitz E, Wascher TC. Low-fat, high-carbohydrate (low-glycaemic index) diet induces weight loss and preserves lean body mass in obese healthy subjects: results of a 24-week study. *Diabetes Obes Metab* 2005;7:290-3.

Balkau B, Vierron E, Vernay M, Born C, Arondel D, Petrella A, et al. The impact of 3-year changes in lifestyle habits on metabolic syndrome parameters: the D.E.S.I.R study. *Eur J Cardiovasc Prev Rehabil*. 2006;13:334-40.

Baran D, Sorensen A, Grimes J, Lew R, Karellas A, Johnson B, et al. Dietary modification with dairy products for preventing vertebral bone loss in premenopausal women: a three-year prospective study. *J Clin Endocrinol Metab*. 1990;70:264-70.

Barclay AW, Petocz P, McMillan-Price J, Flood VM, Prvan T, Mitchell P, et al. Glycemic index, glycemic load, and chronic disease risk--a meta-analysis of observational studies. *Am J Clin Nutr*. 2008;87:627-37.

Barnard ND, Cohen J, Jenkins DJ, Turner-McGrievy G, Gloede L, Green A, et al. A low-fat vegan diet and a conventional diabetes diet in the treatment of type 2 diabetes: a randomized, controlled, 74-wk clinical trial. *Am J Clin Nutr* 2009;89:1588S-1596S.

Bassett DR, Abel M, Moellering RC, Jr., Rosenblatt G, Stokes J, 3rd. Coronary heart disease in Hawaii: dietary intake, depot fat, "stress", smoking, and energy balance in Hawaiian and Japanese men. *Am J Clin Nutr* 1969;22:1483-503.

Beasley JM, Lacroix AZ, Neuhaus ML, Huang Y, Tinker L, Woods N, et al. Protein intake and incident frailty in the women's health initiative observational study. *Journal of the American Geriatrics Society*

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

2010;58:1063-1071.

Bemelmans WJ, Broer J, Hulshof KF, Siero FW, May JF, Meyboom-de Jong B. Long-term effects of nutritional group education for persons at high cardiovascular risk. *Eur J Public Health* 2004;14:240-5.

Ben-Avraham S, Harman-Boehm I, Schwarzfuchs D, Shai I. Dietary strategies for patients with type 2 diabetes in the era of multi-approaches; review and results from the Dietary Intervention Randomized Controlled Trial (DIRECT). *Diabetes Res Clin Pract* 2009;86 Suppl 1:S41-8.

Beresford SA, Johnson KC, Ritenbaugh C, Lasser NL, Snetselaar LG, Black HR, et al. Low-fat dietary pattern and risk of colorectal cancer: the Women's Health Initiative Randomized Controlled Dietary Modification Trial. *JAMA* 2006;295:643-54.

Bergqvist AG, Schall JJ, Stallings VA, Zemel BS. Progressive bone mineral content loss in children with intractable epilepsy treated with the ketogenic diet. *Am J Clin Nutr*. 2008;88:1678-84.

Berkey CS, Rockett HR, Field AE, Gillman MW, Frazier AL, Camargo CA, Jr., et al. Activity, dietary intake, and weight changes in a longitudinal study of preadolescent and adolescent boys and girls. *Pediatrics* 2000;105:E56.

Berkey CS, Rockett HR, Willett WC, Colditz GA. Milk, dairy fat, dietary calcium, and weight gain: a longitudinal study of adolescents. *Arch Pediatr Adolesc Med* 2005;159:543-50.

Bertrais S, Galan P, Renault N, Zarebska M, Preziosi P, Hercberg S. Consumption of soup and nutritional intake in French adults: consequences for nutritional status. *J Hum Nutr Diet*. 2001;14:121-8.

Bertz F, Brekke H, Elleganullrd L, Wennergren M, Rasmussen KM, Winkvist A. Dietary restriction or dietary restriction and exercise, but not exercise intervention alone, reduces weight and fat mass in overweight and obese women postpartum. SOURCE FASEB Journal (2010) 24 Meeting Abstracts. Date of Publication: April, 2010 CONFERENCE NAME Experimental Biology 2010, EB CONFERENCE LOCATION Anaheim, CA, United States CONFERENCE DATE 2010-04-24 to 2010-04-28.

Bes-Rastrollo M, van Dam RM, Martinez-Gonzalez MA, Li TY, Sampson LL, Hu FB. Prospective study of dietary energy density and weight gain in women. *Am J Clin Nutr* 2008;88:769-77.

Beulens JW, de Bruijne LM, Stolk RP, Peeters PH, Bots ML, Grobbee DE, et al. High dietary glycemic load and glycemic index increase risk of cardiovascular disease among middle-aged women: a population-based follow-up study. *J Am Coll Cardiol*. 2007;50:14-21.

Beunza JJ, Toledo E, Hu FB, Bes-Rastrollo M, Serrano-Martinez M, Sanchez-Villegas A, et al. Adherence to the Mediterranean diet, long-term weight change, and incident overweight or obesity: the Seguimiento Universidad de Navarra (SUN) cohort. *Am J Clin Nutr* 2010;92:1484-93.

Biel RK, Friedenreich CM, Cszimadi I, Robson PJ, McLaren L, Faris P, et al. Case-control study of dietary patterns and endometrial cancer risk. *Nutr Cancer* 2011;63:673-86.

Bierenbaum ML, Fleischman AI, Green DP, Raichelson RI, Hayton T, Watson PB, et al. The 5-year experience of modified fat diets on younger men with coronary heart disease. *Circulation* 1970;42:943-52.

SBU. Mat vid fetma. En systematisk litteraturöversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Bingham S, Luben R, Welch A, Tasevska N, Wareham N, Khaw KT. Epidemiologic assessment of sugars consumption using biomarkers: comparisons of obese and nonobese individuals in the European prospective investigation of cancer Norfolk. *Cancer Epidemiol Biomarkers Prev* 2007;16:1651-4.

Bingham SA, Day NE, Luben R, Ferrari P, Slimani N, Norat T, et al. Dietary fibre in food and protection against colorectal cancer in the European Prospective Investigation into Cancer and Nutrition (EPIC): an observational study. *Lancet* 2003;361:1496-501.

Blackburn GL, Copeland T, Khaodhiar L, Buckley RB. Diet and breast cancer. *J Womens Health (Larchmt)* 2003;12:183-92.

Bladbjerg EM, Larsen TM, Due A, Jespersen J, Stender S, Astrup A. Long-term effects on haemostatic variables of three ad libitum diets differing in type and amount of fat and carbohydrate: a 6-month randomised study in obese individuals. *Br J Nutr* 2010;104:1824-30.

Bladbjerg EM, Larsen TM, Due A, Stender S, Astrup A, Jespersen J. Effects on markers of inflammation and endothelial cell function of three ad libitum diets differing in type and amount of fat and carbohydrate: a 6-month randomised study in obese individuals. *Br J Nutr* 2011:1-8.

Blissmer B, Riebe D, Dye G, Ruggiero L, Greene G, Caldwell M. Health-related quality of life following a clinical weight loss intervention among overweight and obese adults: intervention and 24 month follow-up effects. *Health Qual Life Outcomes* 2006;4:43.

Blumenthal JA, Babyak MA, Hinderliter A, Watkins LL, Craighead L, Lin PH, et al. Effects of the DASH diet alone and in combination with exercise and weight loss on blood pressure and cardiovascular biomarkers in men and women with high blood pressure: the ENCORE study. *Arch Intern Med* 2010;170:126-35.

Blumenthal JA, Babyak MA, Sherwood A, Craighead L, Lin PH, Johnson J, et al. Effects of the dietary approaches to stop hypertension diet alone and in combination with exercise and caloric restriction on insulin sensitivity and lipids. *Hypertension* 2010;55:1199-205.

Boden-Albala B, Elkind MS, White H, Szumski A, Paik MC, Sacco RL. Dietary total fat intake and ischemic stroke risk: the Northern Manhattan Study. *Neuroepidemiology*. 2009;32:296-301.

Boniface DR, Tefft ME. Dietary fats and 16-year coronary heart disease mortality in a cohort of men and women in Great Britain. *Eur J Clin Nutr* 2002;56:786-92.

Bopp MJ, Houston DK, Lenchik L, Easter L, Kritchevsky SB, Nicklas BJ. Lean mass loss is associated with low protein intake during dietary-induced weight loss in postmenopausal women. *J Am Diet Assoc* 2008;108:1216-20.

Borgquist S, Wirfalt E, Jirstrom K, Anagnostaki L, Gullberg B, Berglund G, et al. Diet and body constitution in relation to subgroups of breast cancer defined by tumour grade, proliferation and key cell cycle regulators. *Breast Cancer Res*. 2007;9:R11.

Botterweck AA, van den Brandt PA, Goldbohm RA. Vitamins, carotenoids, dietary fiber, and the risk of gastric carcinoma: results from a prospective study after 6.3 years of follow-up. *Cancer* 2000;88:737-48.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Bourn DM, Mann JI, McSkimming BJ, Waldron MA, Wishart JD. Impaired glucose tolerance and NIDDM: Does a lifestyle intervention program have an effect? *Diabetes Care* 1994;17:1311-1319.

Bray GA, Lovejoy JC, Most-Windhauser M, Smith SR, Volaufova J, Denkins Y, et al. A 9-mo randomized clinical trial comparing fat-substituted and fat-reduced diets in healthy obese men: the Ole Study. *Am J Clin Nutr* 2002;76:928-34.

Bray GA, Smith SR, de Jonge L, Xie H, Rood J, Martin CK, et al. Effect of dietary protein content on weight gain, energy expenditure, and body composition during overeating: a randomized controlled trial. *JAMA* 2012;307:47-55.

Brinkworth GD, Buckley JD, Noakes M, Clifton PM, Wilson CJ. Long-term effects of a very low-carbohydrate diet and a low-fat diet on mood and cognitive function. *Arch Intern Med* 2009;169:1873-80.

Brinkworth GD, Buckley JD, Noakes M, Clifton PM. Renal function following long-term weight loss in individuals with abdominal obesity on a very-low-carbohydrate diet vs high-carbohydrate diet. *J Am Diet Assoc* 2010;110:633-8.

Brinkworth GD, Noakes M, Buckley JD, Keogh JB, Clifton PM. Long-term effects of a very-low-carbohydrate weight loss diet compared with an isocaloric low-fat diet after 12 mo. *Am J Clin Nutr* 2009;90:23-32.

Brinkworth GD, Noakes M, Parker B, Foster P, Clifton PM. Long-term effects of advice to consume a high-protein, low-fat diet, rather than a conventional weight-loss diet, in obese adults with type 2 diabetes: one-year follow-up of a randomised trial. *Diabetologia* 2004;47:1677-86.

Buckland G, Bach A, Serra-Majem L. Obesity and the Mediterranean diet: a systematic review of observational and intervention studies. *Obes Rev* 2008;9:582-93.

Burke LE, Choo J, Music E, Warziski M, Styn MA, Kim Y, et al. PREFER study: a randomized clinical trial testing treatment preference and two dietary options in behavioral weight management--rationale, design and baseline characteristics. *Contemp Clin Trials* 2006;27:34-48.

Burke LE, Hudson AG, Warziski MT, Styn MA, Music E, Elci OU, et al. Effects of a vegetarian diet and treatment preference on biochemical and dietary variables in overweight and obese adults: a randomized clinical trial. *Am J Clin Nutr* 2007;86:588-96.

Burke LE, Warziski M, Styn MA, Music E, Hudson AG, Sereika SM. A randomized clinical trial of a standard versus vegetarian diet for weight loss: the impact of treatment preference. *Int J Obes (Lond)* 2008;32:166-76.

Burke V, Lee AH, Hunter E, Spargo R, Smith R, Beilin LJ, et al. Alcohol intake and incidence of coronary disease in Australian aborigines. *Alcohol Alcohol* 2007;42:119-24.

Burrows T, Janet WM, Collins CE. Long-term changes in food consumption trends in overweight children in the HIKCUPS intervention. *J Pediatr Gastroenterol Nutr* 2011;53:543-7.

Buyken AE, Toeller M, Heitkamp G, Vitelli F, Stehle P, Scherbaum WA, et al. Relation of fibre intake to HbA1c and the prevalence of severe ketoacidosis and severe hypoglycaemia. *EURODIAB IDDM*

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Complications Study Group. *Diabetologia* 1998;41:882-90.

Byers T. Nutritional risk factors for breast cancer. *Cancer* 1994;74:288-95.

Cade JE, Burley VJ, Greenwood DC. Dietary fibre and risk of breast cancer in the UK Women's Cohort Study. *Int J Epidemiol* 2007;36:431-8.

Cade JE, Taylor EF, Burley VJ, Greenwood DC. Does the Mediterranean dietary pattern or the Healthy Diet Index influence the risk of breast cancer in a large British cohort of women? *Eur J Clin Nutr* 2011;65:920-8.

Cairella G, Cairella M, Marchini G. Effect of dietary fibre on weight correction after modified fasting. *Eur J Clin Nutr* 1995;49 Suppl 3:S325-7.

Camhi SM, Stefanick ML, Katzmarzyk PT, Young DR. Metabolic syndrome and changes in body fat from a low-fat diet and/or exercise randomized control trial. *Obesity*;18:548-554.

Cardillo S, Seshadri P, Iqbal N. The effects of a low-carbohydrate versus low-fat diet on adipocytokines in severely obese adults: three-year follow-up of a randomized trial. *Eur Rev Med Pharmacol Sci* 2006;10:99-106.

Carels RA, Darby LA, Douglass OM, Cacciapaglia HM, Rydin S. Education on the glycemic index of foods fails to improve treatment outcomes in a behavioral weight loss program. *Eat Behav* 2005;6:145-50.

Carty CL, Kooperberg C, Neuhaus ML, Tinker L, Howard B, Wactawski-Wende J, et al. Low-fat dietary pattern and change in body-composition traits in the Women's Health Initiative Dietary Modification Trial. *American Journal of Clinical Nutrition* 2011;93:516-524.

Champagne CM, Broyles ST, Moran LD, Cash KC, Levy EJ, Lin PH, et al. Dietary intakes associated with successful weight loss and maintenance during the Weight Loss Maintenance trial. *J Am Diet Assoc* 2011;111:1826-35.

Chang-Claude J, Hermann S, Eilber U, Steindorf K. Lifestyle determinants and mortality in German vegetarians and health-conscious persons: results of a 21-year follow-up. *Cancer Epidemiol Biomarkers Prev* 2005;14:963-8.

Chao D, Espeland MA, Farmer D, Register TC, Lenchik L, Applegate WB, et al. Effect of voluntary weight loss on bone mineral density in older overweight women. *J Am Geriatr Soc* 2000;48:753-9.

Cheatham RA, Roberts SB, Das SK, Gilhooly CH, Golden JK, Hyatt R, et al. Long-term effects of provided low and high glycemic load low energy diets on mood and cognition. *Physiol Behav* 2009;98:374-9.

Cheatham PJ, Katz AE. Diet and prostate cancer - a holistic approach to management. *Arch Esp Urol* 2011;64:720-34.

Chen G, Heilbrun LK, Venkatramanamoorthy R, Maranci V, Redd JN, Klurfeld DM, et al. Effects of low-fat and/or high-fruit-and-vegetable diets on plasma levels of 8-isoprostane-F2alpha in the Nutrition and Breast Health study. *Nutr Cancer* 2004;50:155-60.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Cheng G, Karaolis-Danckert N, Libuda L, Bolzenius K, Remer T, Buyken AE. Relation of dietary glycemic index, glycemic load, and fiber and whole-grain intakes during puberty to the concurrent development of percent body fat and body mass index. *Am J Epidemiol.* 2009;169:667-77.

Cho E, Spiegelman D, Hunter DJ, Chen WY, Colditz GA, Willett WC. Premenopausal dietary carbohydrate, glycemic index, glycemic load, and fiber in relation to risk of breast cancer. *Cancer Epidemiol Biomarkers Prev* 2003;12:1153-8.

Choi HK, Atkinson K, Karlson EW, Willett W, Curhan G. Purine-rich foods, dairy and protein intake, and the risk of gout in men. *N Engl J Med.* 2004;350:1093-103.

Christensen P, Bartels EM, Riecke BF, Bliddal H, Leeds AR, Astrup A, et al. Improved nutritional status and bone health after diet-induced weight loss in sedentary osteoarthritis patients: a prospective cohort study. *SOURCE European Journal of Clinical Nutrition* (2011). Date of Publication: 21 Dec 2011.

Christensen R, Astrup A, Bliddal H. Weight loss: the treatment of choice for knee osteoarthritis? A randomized trial. *Osteoarthritis Cartilage* 2005;13:20-7.

Christiansen T, Bruun JM, Madsen EL, Richelsen B. Weight loss maintenance in severely obese adults after an intensive lifestyle intervention: 2- to 4-year follow-up. *Obesity* 2007;15:413-420.

Chrysohoou C, Panagiotakos DB, Aggelopoulos P, Kastorini CM, Kehagia I, Pitsavos C, et al. The Mediterranean diet contributes to the preservation of left ventricular systolic function and to the long-term favorable prognosis of patients who have had an acute coronary event. *Am J Clin Nutr* 2010;92:47-54.

Clifton PM, Keogh JB, Noakes M. Long-term effects of a high-protein weight-loss diet. *Am J Clin Nutr* 2008;87:23-9.

Colditz GA, Manson JE, Stampfer MJ, Rosner B, Willett WC, Speizer FE. Diet and risk of clinical diabetes in women. *Am J Clin Nutr* 1992;55:1018-23.

Coppell KJ, Kataoka M, Williams SM, Chisholm AW, Vorgers SM, Mann JI. Nutritional intervention in patients with type 2 diabetes who are hyperglycaemic despite optimised drug treatment - Lifestyle over and above drugs in diabetes (LOADD) study: Randomised controlled trial. *BMJ* 2010;341:237.

Cornier MA, Donahoo WT, Pereira R, Gurevich I, Westergren R, Enerback S, et al. Insulin sensitivity determines the effectiveness of dietary macronutrient composition on weight loss in obese women. *Obes Res* 2005;13:703-9.

Crawley HF, While D. The diet and body weight of British teenage smokers at 16-17 years. *Eur J Clin Nutr.* 1995;49:904-14.

Croft PR, Brigg D, Smith S, Harrison CB, Branthwaite A, Collins MF. How useful is weight reduction in the management of hypertension? *J R Coll Gen Pract* 1986;36:445-8.

Crowe FL, Roddam AW, Key TJ, Appleby PN, Overvad K, Jakobsen MU, et al. Fruit and vegetable intake and mortality from ischaemic heart disease: results from the European Prospective Investigation into Cancer and Nutrition (EPIC)-Heart study. *Eur Heart J* 2011;32:1235-43.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Dale KS, McAuley KA, Taylor RW, Williams SM, Farmer VL, Hansen P, et al. Determining optimal approaches for weight maintenance: a randomized controlled trial. *CMAJ* 2009;180:E39-46.

Daly ME, Paisey R, Millward BA, Eccles C, Williams K, Hammersley S, et al. Short-term effects of severe dietary carbohydrate-restriction advice in Type 2 diabetes--a randomized controlled trial. *Diabet Med* 2006;23:15-20.

Dargent-Molina P, Sabia S, Touvier M, Kesse E, Breart G, Clavel-Chapelon F, et al. Proteins, dietary acid load, and calcium and risk of postmenopausal fractures in the E3N French women prospective study. *J Bone Miner Res* 2008;23:1915-22.

Dashti HM, Bo-Abbas YY, Asfar SK, Mathew TC, Hussein T, Behbahani A, et al. Ketogenic diet modifies the risk factors of heart disease in obese patients. *Nutrition* 2003;19:901-2.

Davis NJ, Crandall JP, Gajavelli S, Berman JW, Tomuta N, Wylie-Rosett J, et al. Differential effects of low-carbohydrate and low-fat diets on inflammation and endothelial function in diabetes. *J Diabetes Complications* 2011;25:371-6.

Davis NJ, Tomuta N, Isasi CR, Leung V, Wylie-Rosett J. Diabetes-specific quality of life after a low-carbohydrate and low-fat dietary intervention. *Diabetes Educ* 2012;38:250-5.

Davis NJ, Tomuta N, Schechter C, Isasi CR, Segal-Isaacson CJ, Stein D, et al. Comparative study of the effects of a 1-year dietary intervention of a low-carbohydrate diet versus a low-fat diet on weight and glycemic control in type 2 diabetes. *Diabetes Care* 2009;32:1147-52.

Day RS, Fulton JE, Dai S, Mihalopoulos NL, Barradas DT. Nutrient intake, physical activity, and CVD risk factors in children: Project HeartBeat! *Am J Prev Med*. 2009;37:S25-33.

de Goede J, Geleijnse JM, Boer JM, Kromhout D, Verschuren WM. Marine (n-3) fatty acids, fish consumption, and the 10-year risk of fatal and nonfatal coronary heart disease in a large population of Dutch adults with low fish intake. *J Nutr* 2010;140:1023-8.

de las Fuentes L, Waggoner AD, Mohammed BS, Stein RI, Miller BV, 3rd, Foster GD, et al. Effect of moderate diet-induced weight loss and weight regain on cardiovascular structure and function. *J Am Coll Cardiol* 2009;54:2376-81.

de Lorgeril M, Salen P, Martin JL, Monjaud I, Boucher P, Mamelle N. Mediterranean dietary pattern in a randomized trial: prolonged survival and possible reduced cancer rate. *Arch Intern Med* 1998;158:1181-7.

de Lorgeril M, Salen P, Martin JL, Monjaud I, Delaye J, Mamelle N. Mediterranean diet, traditional risk factors, and the rate of cardiovascular complications after myocardial infarction: final report of the Lyon Diet Heart Study. *Circulation* 1999;99:779-85.

de Lorgeril M, Salen P. Mediterranean type of diet for the prevention of coronary heart disease. A global perspective from the seven countries study to the most recent dietary trials. *Int J Vitam Nutr Res* 2001;71:166-72.

de Souza RJ, Bray GA, Carey VJ, Hall KD, LeBoff MS, Loria CM, et al. Effects of 4 weight-loss diets differing in fat, protein, and carbohydrate on fat mass, lean mass, visceral adipose tissue, and hepatic

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

fat: results from the POUNDS LOST trial. *Am J Clin Nutr* 2012;95:614-25.

Deheeger M, Rolland-Cachera MF, Fontvieille AM. Physical activity and body composition in 10 year old French children: linkages with nutritional intake? *Int J Obes Relat Metab Disord*. 1997;21:372-9.

deKoning L, Anand SS. Adherence to a Mediterranean diet and survival in a Greek population. Trichopoulos A, Costacou T, Bamia C, Trichopoulos D. *N Engl J Med* 2003; 348: 2599-608. *Vasc Med* 2004;9:145-6.

Delahanty LM, Hayden D, Ammerman A, Nathan DM. Medical nutrition therapy for hypercholesterolemia positively affects patient satisfaction and quality of life outcomes. *Ann Behav Med*. 2002;24:269-78.

Delbridge EA, Prendergast LA, Pritchard JE, Proietto J. One-year weight maintenance after significant weight loss in healthy overweight and obese subjects: does diet composition matter? *Am J Clin Nutr* 2009;90:1203-14.

D'Elia L, Barba G, Cappuccio FP, Strazzullo P. Potassium intake, stroke, and cardiovascular disease: A meta-analysis of prospective studies. *Journal of the American College of Cardiology* 2011;57:1210-1219.

Demol S, Yackobovitch-Gavan M, Shalitin S, Nagelberg N, Gillon-Keren M, Phillip M. Low-carbohydrate (low & high-fat) versus high-carbohydrate low-fat diets in the treatment of obesity in adolescents. *Acta Paediatr* 2009;98:346-51.

Dietary adherence in the Women's Health Initiative Dietary Modification Trial. *J Am Diet Assoc* 2004;104:654-8.

Djousse L, Levy D, Murabito JM, Cupples LA, Ellison RC. Alcohol consumption and risk of intermittent claudication in the Framingham Heart Study. *Circulation* 2000;102:3092-7.

Djousse L, Rudich T, Gaziano JM. Nut consumption and risk of hypertension in US male physicians. *Clin Nutr* 2009;28:10-4.

Due A, Larsen TM, Hermansen K, Stender S, Holst JJ, Toubro S, et al. Comparison of the effects on insulin resistance and glucose tolerance of 6-mo high-monounsaturated-fat, low-fat, and control diets. *Am J Clin Nutr* 2008;87:855-62.

Due A, Larsen TM, Mu H, Hermansen K, Stender S, Astrup A. Comparison of 3 ad libitum diets for weight-loss maintenance, risk of cardiovascular disease, and diabetes: a 6-mo randomized, controlled trial. *Am J Clin Nutr* 2008;88:1232-41.

Due A, Toubro S, Skov AR, Astrup A. Effect of normal-fat diets, either medium or high in protein, on body weight in overweight subjects: a randomised 1-year trial. *Int J Obes Relat Metab Disord* 2004;28:1283-90.

Due A, Toubro S, Stender S, Skov AR, Astrup A. The effect of diets high in protein or carbohydrate on inflammatory markers in overweight subjects. *Diabetes Obes Metab* 2005;7:223-9.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Dwyer T, Calvert GD, Baghurst KI, Leitch DR. Diet, other lifestyle factors and HDL cholesterol in a population of Australian male service recruits. *Am J Epidemiol* 1981;114:683-96.

Dyson PA, Beatty S, Matthews DR. An assessment of low-carbohydrate or low-fat diets for weight loss at 2 year's follow-up. *Diabet Med* 2010;27:363-4.

Ebbeling CB, Leidig MM, Feldman HA, Lovesky MM, Ludwig DS. Effects of a low-glycemic load vs low-fat diet in obese young adults: a randomized trial. *JAMA* 2007;297:2092-102.

Ebbeling CB, Leidig MM, Sinclair KB, Hangen JP, Ludwig DS. A reduced-glycemic load diet in the treatment of adolescent obesity. *Arch Pediatr Adolesc Med* 2003;157:773-9.

Ebbeling CB, Leidig MM, Sinclair KB, Seger-Shippie LG, Feldman HA, Ludwig DS. Effects of an ad libitum low-glycemic load diet on cardiovascular disease risk factors in obese young adults. *Am J Clin Nutr* 2005;81:976-82.

Elias MC, Parise ER, Carvalho LD, Szejnfeld D, Netto JP. Effect of 6-month nutritional intervention on non-alcoholic fatty liver disease. *Nutrition* 2010;26:1094-1099.

Elkan AC, Sjöberg B, Kolsrud B, Ringertz B, Hafstrom I, Frostegard J. Gluten-free vegan diet induces decreased LDL and oxidized LDL levels and raised atheroprotective natural antibodies against phosphorylcholine in patients with rheumatoid arthritis: A randomized study. *SOURCE Arthritis Research and Therapy* (2008) 10:2 Article Number: R34. Date of Publication: 18 Mar 2008 2008.

Ellingsen I, Hjerkin EM, Arnesen H, Seljeflot I, Hjermann I, Tonstad S. Follow-up of diet and cardiovascular risk factors 20 years after cessation of intervention in the Oslo Diet and Antismoking Study. *Eur J Clin Nutr* 2006;60:378-85.

Elliott SA, Truby H, Lee A, Harper C, Abbott RA, Davies PS. Associations of body mass index and waist circumference with: energy intake and percentage energy from macronutrients, in a cohort of Australian children. *Nutr J* 2011;10:58.

Ello-Martin JA, Roe LS, Ledikwe JH, Beach AM, Rolls BJ. Dietary energy density in the treatment of obesity: a year-long trial comparing 2 weight-loss diets. *Am J Clin Nutr* 2007;85:1465-77.

Ely AC, Banitt A, Befort C, Hou Q, Rhode PC, Grund C, et al. Kansas primary care weighs in: a pilot randomized trial of a chronic care model program for obesity in 3 rural Kansas primary care practices. *J Rural Health* 2008;24:125-32.

Engeset D, Alsaker E, Ciampi A, Lund E. Dietary patterns and lifestyle factors in the Norwegian EPIC cohort: the Norwegian Women and Cancer (NOWAC) study. *Eur J Clin Nutr*. 2005;59:675-84.

Eriksson KF, Lindgärde F. No excess 12-year mortality in men with impaired glucose tolerance who participated in the Malmö Preventive Trial with diet and exercise. *Diabetologia* 1998;1010-6.

Erkkila AT, Lehto S, Pyorala K, Uusitupa MI. n-3 Fatty acids and 5-y risks of death and cardiovascular disease events in patients with coronary artery disease. *Am J Clin Nutr* 2003;78:65-71.

Esposito K, Ima Maiorino M, Ciotola M, Di Palo C, Scognamiglio P, Gicchino M, et al. Effects of a mediterranean-style diet on the need for antihyperglycemic drug therapy in patients with newly diagnosed type 2 diabetes: A randomized trial. *Obstetrical and Gynecological Survey* 2010;65:379-

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

380.

Esposito K, Kastorini CM, Panagiotakos DB, Giugliano D. Mediterranean diet and weight loss: meta-analysis of randomized controlled trials. *Metab Syndr Relat Disord* 2011;9:1-12.

Esposito K, Maiorino MI, Ciotola M, Di Palo C, Scognamiglio P, Gicchino M, et al. Effects of a Mediterranean-style diet on the need for antihyperglycemic drug therapy in patients with newly diagnosed type 2 diabetes: a randomized trial. *Ann Intern Med* 2009;151:306-14.

Esposito K, Marfella R, Ciotola M, Di Palo C, Giugliano F, Giugliano G, et al. Effect of a Mediterranean-style diet on endothelial dysfunction and markers of vascular inflammation in the metabolic syndrome: A randomized trial. *Journal of the American Medical Association* 2004;292:1440-1446.

Esrey KL, Joseph L, Grover SA. Relationship between dietary intake and coronary heart disease mortality: lipid research clinics prevalence follow-up study. *J Clin Epidemiol* 1996;49:211-6.

Estruch R, Martinez-Gonzalez MA, Corella D, Salas-Salvado J, Ruiz-Gutierrez V, Covas MI, et al. Effects of a Mediterranean-style diet on cardiovascular risk factors: a randomized trial. *Ann Intern Med* 2006;145:1-11.

Faria SL, de Oliveira Kelly E, Lins RD, Faria OP. Nutritional management of weight regain after bariatric surgery. *Obes Surg* 2010;20:135-9.

Faria SL, Faria OP, Lopes TC, Galvao MV, de Oliveira Kelly E, Ito MK. Relation between carbohydrate intake and weight loss after bariatric surgery. *Obes Surg* 2009;19:708-16.

Farnsworth E, Luscombe ND, Noakes M, Wittert G, Argyiou E, Clifton PM. Effect of a high-protein, energy-restricted diet on body composition, glycemic control, and lipid concentrations in overweight and obese hyperinsulinemic men and women. *Am J Clin Nutr* 2003;78:31-9.

Fehily AM, Vaughan-Williams E, Shiels K, Williams AH, Horner M, Bingham G, et al. Factors influencing compliance with dietary advice: The diet and reinfarction trial (DART). *Journal of Human Nutrition and Dietetics* 1991;4:33-42.

Feskens EJ, van Dam RM. Dietary fat and the etiology of type 2 diabetes: an epidemiological perspective. *Nutr Metab Cardiovasc Dis* 1999;9:87-95.

Fidanza F, Alberti A, Lanti M, Menotti A. Mediterranean diet score: correlation with 25-year mortality from coronary heart disease in the Seven Countries Study. *Nutr Metab Cardiovasc Dis* 2004;14:397.

Field AE, Aneja P, Austin SB, Shrier LA, de Moor C, Gordon-Larsen P. Race and gender differences in the association of dieting and gains in BMI among young adults. *Obesity (Silver Spring)* 2007;15:456-64.

Fischman ME. CONTROL OF OBESITY IN CHILDREN. *J Med Soc N J* 1965;62:18-22.

Fitzgibbon ML, Stolley MR, Schiffer L, Sharp LK, Singh V, Dyer A. Obesity Reduction Black Intervention Trial (ORBIT): 18-month results. *Obesity* 2010;18:2317-2325.

Fitzwater SL, Weinsier RL, Wooldridge NH, Birch R, Liu C, Bartolucci AA. Evaluation of long-term weight changes after a multidisciplinary weight control program. *J Am Diet Assoc* 1991;91:421-6,

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

429.

Flechtner-Mors M, Boehm BO, Wittmann R, Thoma U, Ditschuneit HH. Enhanced weight loss with protein-enriched meal replacements in subjects with the metabolic syndrome. *Diabetes Metab Res Rev* 2010;26:393-405.

Fleming RM. The effect of high-, moderate-, and low-fat diets on weight loss and cardiovascular disease risk factors. *Prev Cardiol* 2002;5:110-8.

Flood A, Mitchell N, Jaeb M, Finch EA, Laqua PS, Welsh EM, et al. Energy density and weight change in a long-term weight-loss trial. *The International Journal of Behavioral Nutrition and Physical Activity* 2009;6.

Focht BC, Rejeski WJ, Ambrosius WT, Katula JA, Messier SP. Exercise, self-efficacy, and mobility performance in overweight and obese older adults with knee osteoarthritis. *Arthritis Rheum* 2005;53:659-65.

Folsom AR, Demissie Z. Fish intake, marine omega-3 fatty acids, and mortality in a cohort of postmenopausal women. *Am J Epidemiol* 2004;160:1005-10.

Forster M, Veerman JL, Barendregt JJ, Vos T. Cost-effectiveness of diet and exercise interventions to reduce overweight and obesity. *Int J Obes (Lond)* 2011;35:1071-8.

Franceschi S, Dal Maso L, Augustin L, Negri E, Parpinel M, Boyle P, et al. Dietary glycemic load and colorectal cancer risk. *Ann Oncol* 2001;12:173-8.

Freeland KR, Wilson C, Wolever TMS. Adaptation of colonic fermentation and glucagon-like peptide-1 secretion with increased wheat fibre intake for 1 year in hyperinsulinaemic human subjects. *British Journal of Nutrition* 2010;103:82-90.

Friedman AN, Ogden LG, Foster GD, Klein S, Stein R, Miller B, et al. Comparative effects of low-carbohydrate high-protein versus low-fat diets on the kidney. *Clinical Journal of the American Society of Nephrology* 2012;7:1103-1111.

Frisch S, Zittermann A, Berthold HK, Gotting C, Kuhn J, Kleesiek K, et al. A randomized controlled trial on the efficacy of carbohydrate-reduced or fat-reduced diets in patients attending a telemedically guided weight loss program. *Cardiovasc Diabetol* 2009;8:36.

Fung TT, Schulze M, Manson JE, Willett WC, Hu FB. Dietary patterns, meat intake, and the risk of type 2 diabetes in women. *Arch Intern Med* 2004;164:2235-40.

Fung TT, van Dam RM, Hankinson SE, Stampfer M, Willett WC, Hu FB. Low-carbohydrate diets and all-cause and cause-specific mortality: two cohort studies. *Ann Intern Med* 2010;153:289-98.

Gann D. A low-carbohydrate diet in overweight patients undergoing stable statin therapy raises high-density lipoprotein and lowers triglycerides substantially. *Clin Cardiol* 2004;27:563-4.

Gann PH, Chatterton RT, Gapstur SM, Liu K, Garside D, Giovanazzi S, et al. The effects of a low-fat/high-fiber diet on sex hormone levels and menstrual cycling in premenopausal women: a 12-month randomized trial (the diet and hormone study). *Cancer* 2003;98:1870-9.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Gao SK, Beresford SA, Frank LL, Schreiner PJ, Burke GL, Fitzpatrick AL. Modifications to the Healthy Eating Index and its ability to predict obesity: the Multi-Ethnic Study of Atherosclerosis. *Am J Clin Nutr* 2008;88:64-9.

Gardner CD, Kiazand A, Alhassan S, Kim S, Stafford RS, Balise RR, et al. Comparison of the Atkins, Zone, Ornish, and LEARN diets for change in weight and related risk factors among overweight premenopausal women: the A TO Z Weight Loss Study: a randomized trial. *JAMA* 2007;297:969-77.

Gately PJ, King NA, Greatwood HC, Humphrey LC, Radley D, Cooke CB, et al. Does a high-protein diet improve weight loss in overweight and obese children? *Obesity (Silver Spring)* 2007;15:1527-34.

Genkinger JM, Hunter DJ, Spiegelman D, Anderson KE, Buring JE, Freudenheim JL, et al. Alcohol intake and ovarian cancer risk: a pooled analysis of 10 cohort studies. *Br J Cancer*. 2006;94:757-62.

Gilliam ML. Long-term effects of weight-reducing diets in hypertensive patients. *Obstetrics and gynecology* 2012;119:180-181.

Giugliano D, Ceriello A, Esposito K. The Effects of Diet on Inflammation. Emphasis on the Metabolic Syndrome. *Journal of the American College of Cardiology* 2006;48:677-685.

Gogebakan O, Kohl A, Osterhoff MA, van Baak MA, Jebb SA, Papadaki A, et al. Effects of weight loss and long-term weight maintenance with diets varying in protein and glycemic index on cardiovascular risk factors: the diet, obesity, and genes (DIOGenes) study: a randomized, controlled trial. *Circulation* 2011;124:2829-38.

Golan R, Schwarzfuchs D, Stampfer MJ, Shai I, group D. Halo effect of a weight-loss trial on spouses: the DIRECT-Spouse study. *Public Health Nutr* 2009;13:544-9.

Gonzalez CA, Riboli E. Diet and cancer prevention: Contributions from the European Prospective Investigation into Cancer and Nutrition (EPIC) study. *Eur J Cancer* 2010;46:2555-62.

Goodman MT, Nomura AM, Wilkens LR, Hankin J. The association of diet, obesity, and breast cancer in Hawaii. *Cancer Epidemiol Biomarkers Prev* 1992;1:269-75.

Gorin AA, Phelan S, Wing RR, Hill JO. Promoting long-term weight control: does dieting consistency matter? *Int J Obes Relat Metab Disord* 2004;28:278-81.

Goyenechea E, Holst C, van Baak MA, Saris WHM, Jebb S, Kafatos A, et al. Effects of different protein content and glycaemic index of ad libitum diets on diabetes risk factors in overweight adults: The DIOGenes multicentre, randomized, dietary intervention trial. *Diabetes/metabolism research and reviews* 2011;27:705-716.

IGreenberg I, Stampfer MJ, Schwarzfuchs D, Shai I, Group D. Adherence and success in long-term weight loss diets: the dietary intervention randomized controlled trial (DIRECT). *J Am Coll Nutr* 2009;28:159-68.

IGriffin HJ, O'Connor HT, Rooney KB, Steinbeck KS, Cheng HL, Petocz P. Comparison of higher-protein and higher-carbohydrate diets in overweight and obese young women. *Obesity Reviews* 2010;11:823.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

IGuldbrand H, Dizdar B, Bunjaku B, Lindstrom T, Bachrach-Lindstrom M, Fredrikson M, et al. In type 2 diabetes, randomisation to advice to follow a low-carbohydrate diet transiently improves glycaemic control compared with advice to follow a low-fat diet producing a similar weight loss. *Diabetologia* 2012;55:2118-2127.

IGunther AL, Buyken AE, Kroke A. The influence of habitual protein intake in early childhood on BMI and age at adiposity rebound: results from the DONALD Study. *Int J Obes (Lond)*. 2006;30:1072-9.

IHailu A, Knutsen SF, Fraser GE. Associations between meat consumption and the prevalence of degenerative arthritis and soft tissue disorders in the adventist health study, California U.S.A. *J Nutr Health Aging*. 2006;10:7-14.

IHakanen M, Lagstrom H, Pahkala K, Sillanmaki L, Saarinen M, Niinikoski H, et al. Dietary and lifestyle counselling reduces the clustering of overweight-related cardiometabolic risk factors in adolescents. *Acta Paediatr* 2009;99:888-95.

IHalkjaer J, Olsen A, Overvad K, Jakobsen MU, Boeing H, Buijsse B, et al. Intake of total, animal and plant protein and subsequent changes in weight or waist circumference in European men and women: the Diogenes project. *Int J Obes (Lond)* 2011;35:1104-13.

IHall WD, Feng Z, George VA, Lewis CE, Oberman A, Huber M, et al. Low-fat diet: effect on anthropometrics, blood pressure, glucose, and insulin in older women. *Ethn Dis* 2003;13:337-43.

Hall Y, Stamler J, Cohen DB, Mojonier L, Epstein MB, Berkson DM, et al. Effectiveness of a low saturated fat, low cholesterol, weight-reducing diet for the control of hypertriglyceridemia. *Atherosclerosis* 1972;16:389-403.

Halton TL, Liu S, Manson JE, Hu FB. Low-carbohydrate-diet score and risk of type 2 diabetes in women. *Am J Clin Nutr* 2008;87:339-46.

Halton TL, Willett WC, Liu S, Manson JE, Albert CM, Rexrode K, et al. Low-carbohydrate-diet score and the risk of coronary heart disease in women. *N Engl J Med* 2006;355:1991-2002.

Hannan MT, Tucker KL, Dawson-Hughes B, Cupples LA, Felson DT, Kiel DP. Effect of dietary protein on bone loss in elderly men and women: the Framingham Osteoporosis Study. *J Bone Miner Res*. 2000;15:2504-12.

Hardy DS, Hoelscher DM, Aragaki C, Stevens J, Steffen LM, Pankow JS, et al. Association of glycemic index and glycemic load with risk of incident coronary heart disease among Whites and African Americans with and without type 2 diabetes: the Atherosclerosis Risk in Communities study. *Ann Epidemiol* 2010;20:610-6.

Hare-Bruun H, Flint A, Heitmann BL. Glycemic index and glycemic load in relation to changes in body weight, body fat distribution, and body composition in adult Danes. *Am J Clin Nutr* 2006;84:871-9; quiz 952-3.

Harvey-Berino J. Calorie restriction is more effective for obesity treatment than dietary fat restriction. *Ann Behav Med* 1999;21:35-9.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Harvey-Berino J. The efficacy of dietary fat vs. total energy restriction for weight loss. *Obes Res* 1998;6:202-7.

Haufe S, Engeli S, Kast P, Bohnke J, Utz W, Haas V, et al. Randomized comparison of reduced fat and reduced carbohydrate hypocaloric diets on intrahepatic fat in overweight and obese human subjects. *Hepatology* 2011;53:1504-14.

Haufe S, Utz W, Engeli S, Kast P, Bohnke J, Pofahl M, et al. Left ventricular mass and function with reduced-fat or reduced-carbohydrate hypocaloric diets in overweight and obese subjects. *Hypertension* 2012;59:70-5.

He M, van Dam RM, Rimm E, Hu FB, Qi L. Whole-grain, cereal fiber, bran, and germ intake and the risks of all-cause and cardiovascular disease-specific mortality among women with type 2 diabetes mellitus. *Circulation* 2010;121:2162-8.

Heitmann BL, Lissner L, Sorensen TI, Bengtsson C. Dietary fat intake and weight gain in women genetically predisposed for obesity. *Am J Clin Nutr* 1995;61:1213-7.

Heshka S, Greenway F, Anderson JW, Atkinson RL, Hill JO, Phinney SD, et al. Self-help weight loss versus a structured commercial program after 26 weeks: a randomized controlled study. *Am J Med* 2000;109:282-7.

Hinton PS, Rector RS, Donnelly JE, Smith BK, Bailey B. Total body bone mineral content and density during weight loss and maintenance on a low- or recommended-dairy weight-maintenance diet in obese men and women. *Eur J Clin Nutr* 2010;64:392-9.

Hjermann I, Leren P, Norman N, Helgeland A, Holme I. Serum insulin response to oral glucose load during a dietary intervention trial in healthy coronary high risk men: the Oslo study. *Scand J Clin Lab Invest* 1980;40:89-94.

Hockaday TD, Hockaday JM, Mann JJ, Turner RC. Prospective comparison of modified fat-high-carbohydrate with standard low-carbohydrate dietary advice in the treatment of diabetes: one year follow-up study. *Br J Nutr* 1978;39:357-62.

Hodge AM, English DR, O'Dea K, Giles GG. Glycemic index and dietary fiber and the risk of type 2 diabetes. *Diabetes Care* 2004;27:2701-6.

Hoffman JA, Thompson DR, Franko DL, Power TJ, Leff SS, Stallings VA. Decaying behavioral effects in a randomized, multi-year fruit and vegetable intake intervention. *Prev Med* 2011;52:370-5.

Hollis JF, Gullion CM, Stevens VJ, Brantley PJ, Appel LJ, Ard JD, et al. Weight loss during the intensive intervention phase of the weight-loss maintenance trial. *Am J Prev Med* 2008;35:118-26.

Hooper L, Abdelhamid A, Moore HJ, Douthwaite W, Skeaff CM, Summerbell CD. Effect of reducing total fat intake on body weight: systematic review and meta-analysis of randomised controlled trials and cohort studies. *BMJ* 2012;345:e7666.

Hooper L, Thompson RL, Harrison RA, Summerbell CD, Ness AR, Moore HJ, et al. Risks and benefits of omega 3 fats for mortality, cardiovascular disease, and cancer: systematic review. *BMJ* 2006;332:752-60.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Hopping BN, Erber E, Grandinetti A, Verheus M, Kolonel LN, Maskarinec G. Dietary fiber, magnesium, and glycemic load alter risk of type 2 diabetes in a multiethnic cohort in Hawaii. *J Nutr* 2010;140:68-74.

Howard BV, Curb JD, Eaton CB, Kooperberg C, Ockene J, Kostis JB, et al. Low-fat dietary pattern and lipoprotein risk factors: the Women's Health Initiative Dietary Modification Trial. *Am J Clin Nutr* 2010;91:860-74.

Howard BV, Manson JE, Stefanick ML, Beresford SA, Frank G, Jones B, et al. Low-fat dietary pattern and weight change over 7 years: the Women's Health Initiative Dietary Modification Trial. *JAMA* 2006;295:39-49.

Howard BV, Van Horn L, Hsia J, Manson JE, Stefanick ML, Wassertheil-Smoller S, et al. Low-fat dietary pattern and risk of cardiovascular disease: the Women's Health Initiative Randomized Controlled Dietary Modification Trial. *JAMA* 2006;295:655-66.

Howarth NC, Murphy SP, Wilkens LR, Hankin JH, Kolonel LN. Dietary energy density is associated with overweight status among 5 ethnic groups in the multiethnic cohort study. *J Nutr* 2006;136:2243-8.

Hoy MK, Winters BL, Chlebowski RT, Papoutsakis C, Shapiro A, Lubin MP, et al. Implementing a low-fat eating plan in the Women's Intervention Nutrition Study. *J Am Diet Assoc* 2009;109:688-96.

Hsieh LJ, Carter HB, Landis PK, Tucker KL, Metter EJ, Newschaffer CJ, et al. Association of energy intake with prostate cancer in a long-term aging study: Baltimore Longitudinal Study of Aging (United States). *Urology*. 2003;61:297-301.

Hu FB, Rimm EB, Stampfer MJ, Ascherio A, Spiegelman D, Willett WC. Prospective study of major dietary patterns and risk of coronary heart disease in men. *Am J Clin Nutr* 2000;72:912-21.

Hu FB, Stampfer MJ, Manson JE, Grodstein F, Colditz GA, Speizer FE, et al. Trends in the incidence of coronary heart disease and changes in diet and lifestyle in women. *N Engl J Med* 2000;343:530-7.

Hu T, Mills KT, Yao L, Demanelis K, Eloustaz M, Yancy WS, et al. Effects of low-carbohydrate diets versus low-fat diets on metabolic risk factors: A meta-analysis of randomized controlled clinical trials. *American Journal of Epidemiology* 2012;176:S44-S54.

Huijbregts P, Feskens E, Rasanen L, Fidanza F, Nissinen A, Menotti A, et al. Dietary pattern and 20 year mortality in elderly men in Finland, Italy, and The Netherlands: longitudinal cohort study. *BMJ* 1997;315:13-7.

Hulley SB, Wilson WS, Burrows MI, Nichaman MZ. Lipid and lipoprotein responses of hypertriglyceridaemic outpatients to a low-carbohydrate modification of the A.H.A. fat-controlled diet. *Lancet* 1972;2:551-5.

Hyden S, Tyroler HA, Hames CG, Bartel A, Thompson JW, Krishan I, et al. Diet treatment of obese hypertensives. *Clin Sci Mol Med Suppl* 1973;45 Suppl 1:209s-12.

Imayama I, Alfano CM, Kong A, Foster-Schubert KE, Bain CE, Xiao L, et al. Dietary weight loss and exercise interventions effects on quality of life in overweight/obese postmenopausal women: a randomized controlled trial. *Int J Behav Nutr Phys Act* 2011;8:118.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Insull W, Jr., Henderson MM, Prentice RL, Thompson DJ, Clifford C, Goldman S, et al. Results of a randomized feasibility study of a low-fat diet. *Arch Intern Med* 1990;150:421-7.

Iqbal SI, Helge JW, Heitmann BL. Do energy density and dietary fiber influence subsequent 5-year weight changes in adult men and women? *Obesity (Silver Spring)* 2006;14:106-14.

Iribarren C, Markovitz JH, Jacobs DR, Jr., Schreiner PJ, Daviglius M, Hibbeln JR. Dietary intake of n-3, n-6 fatty acids and fish: relationship with hostility in young adults--the CARDIA study. *Eur J Clin Nutr* 2004;58:24-31.

Iago R, Nicklas T, Yang SJ, Baranowski T, Zakeri I, Berenson GS. Physical activity and health enhancing dietary behaviors in young adults: Bogalusa Heart Study. *Prev Med* 2005;41:194-202.

Jain MG, Rohan TE, Howe GR, Miller AB. A cohort study of nutritional factors and endometrial cancer. *Eur J Epidemiol* 2000;16:899-905.

Jarvinen R, Knekt P, Rissanen H, Reunanen A. Intake of fish and long-chain n-3 fatty acids and the risk of coronary heart mortality in men and women. *Br J Nutr* 2006;95:824-9.

Jazet IM, de Craen AJ, van Schie EM, Meinders AE. Sustained beneficial metabolic effects 18 months after a 30-day very low calorie diet in severely obese, insulin-treated patients with type 2 diabetes. *Diabetes Res Clin Pract* 2007;77:70-6.

Jebb SA, Lovegrove JA, Griffin BA, Frost GS, Moore CS, Chatfield MD, et al. Effect of changing the amount and type of fat and carbohydrate on insulin sensitivity and cardiovascular risk: the RISCK (Reading, Imperial, Surrey, Cambridge, and Kings) trial. *Am J Clin Nutr* 2010;92:748-58.

Jeffery RW, Hellerstedt WL, French SA, Baxter JE. A randomized trial of counseling for fat restriction versus calorie restriction in the treatment of obesity. *Int J Obes Relat Metab Disord* 1995;19:132-7.

Jenkins DJ, Kendall CW, Augustin LS, Franceschi S, Hamidi M, Marchie A, et al. Glycemic index: overview of implications in health and disease. *Am J Clin Nutr* 2002;76:266S-73S.

Jiao L, Flood A, Subar AF, Hollenbeck AR, Schatzkin A, Stolzenberg-Solomon R. Glycemic index, carbohydrates, glycemic load, and the risk of pancreatic cancer in a prospective cohort study. *Cancer Epidemiol Biomarkers Prev*. 2009;18:1144-51.

Johnson L, Mander AP, Jones LR, Emmett PM, Jebb SA. Energy-dense, low-fiber, high-fat dietary pattern is associated with increased fatness in childhood. *Am J Clin Nutr* 2008;87:846-54.

Johnson ST, Newton AS, Chopra M, Buckingham J, Huang TTK, Franks PW, et al. In search of quality evidence for lifestyle management and glycemic control in children and adolescents with type 2 diabetes: A systematic review. *SOURCE BMC Pediatrics* (2010) 10 Article Number: 97. Date of Publication: 23 Dec 2010 2010.

Jones DW, Miller ME, Wofford MR, Anderson DC, Jr., Cameron ME, Willoughby DL, et al. The effect of weight loss intervention on antihypertensive medication requirements in the hypertension Optimal Treatment (HOT) study. *Am J Hypertens* 1999;12:1175-80.

Josse AR, Atkinson SA, Tarnopolsky MA, Phillips SM. Diets higher in dairy foods and dietary protein support bone health during diet- and exercise-induced weight loss in overweight and obese

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

premenopausal women. *J Clin Endocrinol Metab* 2012;97:251-60.

Kabagambe EK, Baylin A, Campos H. Nonfatal acute myocardial infarction in Costa Rica: modifiable risk factors, population-attributable risks, and adherence to dietary guidelines. *Circulation* 2007;115:1075-81.

Kafatos A, Diacatou A, Voukiklaris G, Nikolakakis N, Vlachonikolis J, Kounali D, et al. Heart disease risk-factor status and dietary changes in the Cretan population over the past 30 y: the Seven Countries Study. *Am J Clin Nutr* 1997;65:1882-6.

Kant AK, Graubard BI, Schatzkin A, Ballard-Barbash R. Proportion of energy intake from fat and subsequent weight change in the NHANES I Epidemiologic Follow-up Study. *Am J Clin Nutr* 1995;61:11-7.

Kant AK, Leitzmann MF, Park Y, Hollenbeck A, Schatzkin A. Patterns of recommended dietary behaviors predict subsequent risk of mortality in a large cohort of men and women in the United States. *J Nutr*. 2009;139:1374-80.

Kant AK. Weight-loss attempts and reporting of foods and nutrients, and biomarkers in a national cohort. *Int J Obes Relat Metab Disord* 2002;26:1194-204.

Karlsson J, Hallgren P, Kral J, Lindroos AK, Sjostrom L, Sullivan M. Predictors and effects of long-term dieting on mental well-being and weight loss in obese women. *Appetite* 1994;23:15-26.

Kato I, Toniolo P, Zeleniuch-Jacquotte A, Shore RE, Koenig KL, Akhmedkhanov A, et al. Diet, smoking and anthropometric indices and postmenopausal bone fractures: a prospective study. *Int J Epidemiol* 2000;29:85-92.

Kelemen LE, Kushi LH, Jacobs DR, Jr., Cerhan JR. Associations of dietary protein with disease and mortality in a prospective study of postmenopausal women. *Am J Epidemiol* 2005;161:239-49.

Kelishadi R, Zemel MB, Hashemipour M, Hosseini M, Mohammadifard N, Poursafa P. Can a dairy-rich diet be effective in long-term weight control of young children? *J Am Coll Nutr* 2009;28:601-10.

Kerksick CM, Wismann-Bunn J, Fogt D, Thomas AR, Taylor L, Campbell BI, et al. Changes in weight loss, body composition and cardiovascular disease risk after altering macronutrient distributions during a regular exercise program in obese women. *Nutr J* 2010;9:59.

Khaw KT, Barrett-Connor E. Dietary fiber and reduced ischemic heart disease mortality rates in men and women: a 12-year prospective study. *Am J Epidemiol* 1987;126:1093-102.

Khoo J, Piantadosi C, Duncan R, Worthley SG, Jenkins A, Noakes M, et al. Comparing effects of a low-energy diet and a high-protein low-fat diet on sexual and endothelial function, urinary tract symptoms, and inflammation in obese diabetic men. *J Sex Med* 2011;8:2868-75.

Kinlen LJ. Diet and breast cancer. *Br Med Bull* 1991;47:462-9.

Kirk S, Brehm B, Saelens BE, Woo JG, Kissel E, D'Alessio D, et al. Role of Carbohydrate Modification in Weight Management among Obese Children: A Randomized Clinical Trial. *SOURCE Journal of Pediatrics* (2012).

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Kleber M, Schaefer A, Winkel K, Hoffmann D, Wunsch R, Kersting M, et al. Lifestyle intervention "Obeldicks Mini" for obese children aged 4 to 7 years. *Klin Padiatr* 2009;221:290-4.

Klein S. Clinical Trial Experience with Fat-Restricted vs. Carbohydrate-Restricted Weight-Loss Diets. *Obesity Research* 2004;12:141S-144S.

Klemsdal TO, Holme I, Nerland H, Pedersen TR, Tonstad S. Effects of a low glycemic load diet versus a low-fat diet in subjects with and without the metabolic syndrome. *Nutr Metab Cardiovasc Dis* 2010;20:195-201.

Klesges RC, Klesges LM, Haddock CK, Eck LH. A longitudinal analysis of the impact of dietary intake and physical activity on weight change in adults. *Am J Clin Nutr* 1992;55:818-22.

Knight JA, Martin LJ, Greenberg CV, Lockwood GA, Byng JW, Yaffe MJ, et al. Macronutrient intake and change in mammographic density at menopause: results from a randomized trial. *Cancer Epidemiol Biomarkers Prev* 1999;8:123-8.

Koh-Banerjee P, Franz M, Sampson L, Liu S, Jacobs DR, Jr., Spiegelman D, et al. Changes in whole-grain, bran, and cereal fiber consumption in relation to 8-y weight gain among men. *Am J Clin Nutr*. 2004;80:1237-45.

Koletzko B, Toschke AM. Meal patterns and frequencies: do they affect body weight in children and adolescents? *Crit Rev Food Sci Nutr* 2010;50:100-5.

Kong A, Beresford SA, Alfano CM, Foster-Schubert KE, Neuhauser ML, Johnson DB, et al. Associations between snacking and weight loss and nutrient intake among postmenopausal overweight to obese women in a dietary weight-loss intervention. *J Am Diet Assoc* 2011;111:1898-903.

Koppes LL, Boon N, Nooyens AC, van Mechelen W, Saris WH. Macronutrient distribution over a period of 23 years in relation to energy intake and body fatness. *Br J Nutr*. 2009;101:108-15.

Korhonen M, Kastarinen M, Uusitupa M, Puska P, Nissinen A. The effect of intensified diet counseling on the diet of hypertensive subjects in primary health care: A 2-year open randomized controlled trial of lifestyle intervention against hypertension in eastern Finland. *Preventive Medicine* 2003;36:8-16.

Kouris-Blazos A, Gnardellis C, Wahlqvist ML, Trichopoulos D, Lukito W, Trichopoulou A. Are the advantages of the Mediterranean diet transferable to other populations? A cohort study in Melbourne, Australia. *Br J Nutr* 1999;82:57-61.

Kretsch MJ, Green MW, Fong AK, Elliman NA, Johnson HL. Cognitive effects of a long-term weight reducing diet. *Int J Obes Relat Metab Disord* 1997;21:14-21.

Kring SI, Heitmann BL. Fiber intake, not dietary energy density, is associated with subsequent change in BMI z-score among sub-groups of children. *Obes Facts* 2008;1:331-8.

Krishnan S, Rosenberg L, Singer M, Hu FB, Djousse L, Cupples LA, et al. Glycemic index, glycemic load, and cereal fiber intake and risk of type 2 diabetes in US black women. *Arch Intern Med* 2007;167:2304-9.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Kromhout D, Bosschieter EB, de Lezenne Coulander C. Dietary fibre and 10-year mortality from coronary heart disease, cancer, and all causes. The Zutphen study. *Lancet* 1982;2:518-22.

Kromhout D, Saris WH, Horst CH. Energy intake, energy expenditure, and smoking in relation to body fatness: the Zutphen Study. *Am J Clin Nutr.* 1988;47:668-74.

Kuczmarski MF, Cremer Sees A, Hotchkiss L, Cotugna N, Evans MK, Zonderman AB. Higher Healthy Eating Index-2005 scores associated with reduced symptoms of depression in an urban population: findings from the Healthy Aging in Neighborhoods of Diversity Across the Life Span (HANDLS) study. *J Am Diet Assoc* 2010;110:383-9.

Kuller LH, Simkin-Silverman LR, Wing RR, Meilahn EN, Ives DG. Women's Healthy Lifestyle Project: A randomized clinical trial: results at 54 months. *Circulation* 2001;103:32-7.

Kumanyika SK, Espeland MA, Bahnson JL, Bottom JB, Charleston JB, Folmar S, et al. Ethnic comparison of weight loss in the Trial of Nonpharmacologic Interventions in the Elderly. *Obes Res* 2002;10:96-106.

Laaksonen DE, Nyyssonen K, Niskanen L, Rissanen TH, Salonen JT. Prediction of cardiovascular mortality in middle-aged men by dietary and serum linoleic and polyunsaturated fatty acids. *Arch Intern Med* 2005;165:193-9.

Lagiou P, Sandin S, Lof M, Trichopoulos D, Adami HO, Weiderpass E. Low carbohydrate-high protein diet and incidence of cardiovascular diseases in Swedish women: prospective cohort study. *BMJ* 2012;344:e4026.

Lagiou P, Sandin S, Weiderpass E, Lagiou A, Mucci L, Trichopoulos D, et al. Low carbohydrate-high protein diet and mortality in a cohort of Swedish women. *J Intern Med* 2007;261:366-74.

Lagiou P, Trichopoulos D, Sandin S, Lagiou A, Mucci L, Wolk A, et al. Mediterranean dietary pattern and mortality among young women: a cohort study in Sweden. *Br J Nutr* 2006;96:384-92.

Lapointe A, Provencher V, Weisnagel SJ, Begin C, Blanchet R, Dufour-Bouchard AA, et al. Dietary intervention promoting high intakes of fruits and vegetables: short-term effects on eating behaviors in overweight-obese postmenopausal women. *Eat Behav* 2010;11:305-8.

Lapointe A, Weisnagel SJ, Provencher V, Begin C, Dufour-Bouchard AA, Trudeau C, et al. Comparison of a dietary intervention promoting high intakes of fruits and vegetables with a low-fat approach: long-term effects on dietary intakes, eating behaviours and body weight in postmenopausal women. *Br J Nutr* 2010;104:1080-90.

Lapointe A, Weisnagel SJ, Provencher V, Begin C, Dufour-Bouchard AA, Trudeau C, et al. Using restrictive messages to limit high-fat foods or nonrestrictive messages to increase fruit and vegetable intake: what works better for postmenopausal women? *Eur J Clin Nutr* 2010;64:194-202.

Larsen TM, Dalskov SM, van Baak M, Jebb SA, Papadaki A, Pfeiffer AF, et al. Diets with high or low protein content and glycemic index for weight-loss maintenance. *N Engl J Med* 2010;363:2102-13.

Larsson SC, Bergkvist L, Wolk A. Fruit and vegetable consumption and incidence of gastric cancer: a prospective study. *Cancer Epidemiol Biomarkers Prev* 2006;15:1998-2001.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Larsson SC, Bergkvist L, Wolk A. Glycemic load, glycemic index and breast cancer risk in a prospective cohort of Swedish women. *Int J Cancer*. 2009;125:153-7.

Larsson SC, Friberg E, Wolk A. Carbohydrate intake, glycemic index and glycemic load in relation to risk of endometrial cancer: A prospective study of Swedish women. *Int J Cancer* 2007;120:1103-7.

Lasker DAW, Evans EM, Layman DK. Moderate carbohydrate, moderate protein weight loss diet reduces cardiovascular disease risk compared to high carbohydrate, low protein diet in obese adults: A randomized clinical trial. *SOURCE Nutrition and Metabolism* (2008) 5:1 Article Number: 30. Date of Publication: 2008 2008.

Lavery MA, Loewy JW, Kapadia AS, Nichaman MZ, Foreyt JP, Gee M. Long-term follow-up of weight status of subjects in a behavioral weight control program. *J Am Diet Assoc* 1989;89:1259-64.

Layman DK, Evans EM, Erickson D, Seyler J, Weber J, Bagshaw D, et al. A moderate-protein diet produces sustained weight loss and long-term changes in body composition and blood lipids in obese adults. *J Nutr* 2009;139:514-21.

Lean MEJ, Han TS, Prvan T, Richmond PR, Avenell A. Weight loss with high and low carbohydrate 1200 kcal diets in free living women. *European Journal of Clinical Nutrition* 1997;51:243-248.

Leichtle AB, Helmschrodt C, Ceglarek U, Shai I, Henkin Y, Schwarzfuchs D, et al. Effects of a 2-y dietary weight-loss intervention on cholesterol metabolism in moderately obese men. *Am J Clin Nutr* 2011;94:1189-95.

Leidy HJ, Carnell NS, Mattes RD, Campbell WW. Higher protein intake preserves lean mass and satiety with weight loss in pre-obese and obese women. *Obesity (Silver Spring)* 2007;15:421-9.

Lejeune MP, Kovacs EM, Westerterp-Plantenga MS. Additional protein intake limits weight regain after weight loss in humans. *Br J Nutr* 2005;93:281-9.

Lemaitre RN, King IB, Mozaffarian D, Kuller LH, Tracy RP, Siscovick DS. n-3 Polyunsaturated fatty acids, fatal ischemic heart disease, and nonfatal myocardial infarction in older adults: the Cardiovascular Health Study. *Am J Clin Nutr* 2003;77:319-25.

Leosdottir M, Nilsson P, Nilsson JA, Mansson H, Berglund G. The association between total energy intake and early mortality: data from the Malmo Diet and Cancer Study. *J Intern Med* 2004;256:499-509.

Lepe M, Bacardi Gascon M, Jimenez Cruz A. Long-term efficacy of high-protein diets: a systematic review. *Nutr Hosp* 2011;26:1256-9.

Leren P. The effect of plasma cholesterol lowering diet in male survivors of myocardial infarction. A controlled clinical trial. *Acta Med Scand Suppl* 1966;466:1-92.

Lewis KJ, Doyle MD. Nutrient intake and weight response of women on weight-control diets. *J Am Diet Assoc* 1970;56:119-25.

Levitan EB, Mittleman MA, Wolk A. Dietary glycaemic index, dietary glycaemic load and incidence of myocardial infarction in women. *Br J Nutr* 2010;103:1049-55.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Levitan EB, Wolk A, Mittleman MA. Fish consumption, marine omega-3 fatty acids, and incidence of heart failure: a population-based prospective study of middle-aged and elderly men. *Eur Heart J* 2009;30:1495-500.

Li Z, Treyzon L, Chen S, Yan E, Thames G, Carpenter CL. Protein-enriched meal replacements do not adversely affect liver, kidney or bone density: an outpatient randomized controlled trial. *Nutr J* 2011;9:72.

Li Z, Treyzon L, Chen S, Yan E, Thames G, Carpenter CL. Protein-enriched meal replacements do not adversely affect liver, kidney or bone density: an outpatient randomized controlled trial. *Nutr J* 2010;9:72.

Lien LF, Brown AJ, Ard JD, Loria C, Erlinger TP, Feldstein AC, et al. Effects of PREMIER lifestyle modifications on participants with and without the metabolic syndrome. *Hypertension* 2007;50:609-16.

Liese AD, Weis KE, Schulz M, Tooze JA. Food intake patterns associated with incident type 2 diabetes: the Insulin Resistance Atherosclerosis Study. *Diabetes Care* 2009;32:263-8.

Lim SS, Noakes M, Keogh JB, Clifton PM. Long-term effects of a low carbohydrate, low fat or high unsaturated fat diet compared to a no-intervention control. *Nutr Metab Cardiovasc Dis* 2010;20:599-607.

Lin J, Zhang SM, Cook NR, Rexrode KM, Liu S, Manson JE, et al. Dietary intakes of fruit, vegetables, and fiber, and risk of colorectal cancer in a prospective cohort of women (United States). *Cancer Causes Control* 2005;16:225-33.

Lin PH, Wang Y, Grambow SC, Goggins W, Almirall D. Dietary saturated fat intake is negatively associated with weight maintenance among the PREMIER participants. *Obesity (Silver Spring)* 2012;20:571-5.

Linos E, Holmes MD, Willett WC. Diet and breast cancer. *Current Oncology Reports* 2007;9:31-41.

Lissner L, Heitmann BL. Dietary fat and obesity: evidence from epidemiology. *Eur J Clin Nutr* 1995;49:79-90.

Liu S, Willett WC, Stampfer MJ, Hu FB, Franz M, Sampson L, et al. A prospective study of dietary glycemic load, carbohydrate intake, and risk of coronary heart disease in US women. *Am J Clin Nutr*. 2000;71:1455-61.

Logan KJ, Woodside JV, Young IS, McKinley MC, Perkins-Porras L, McKeown PP. Adoption and maintenance of a Mediterranean diet in patients with coronary heart disease from a Northern European population: a pilot randomised trial of different methods of delivering Mediterranean diet advice. *Journal of Human Nutrition & Dietetics* 2010;23:30-37.

Ludwig DS, Pereira MA, Kroenke CH, Hilner JE, Van Horn L, Slattery ML, et al. Dietary fiber, weight gain, and cardiovascular disease risk factors in young adults. *JAMA* 1999;282:1539-46.

Lundgren H, Bengtsson C, Blohme G, Isaksson B, Lapidus L, Lenner RA, et al. Dietary habits and incidence of noninsulin-dependent diabetes mellitus in a population study of women in Gothenburg,

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Sweden. *Am J Clin Nutr*. 1989;49:708-12.

Luscombe-Marsh ND, Noakes M, Wittert GA, Keogh JB, Foster P, Clifton PM. Carbohydrate-restricted diets high in either monounsaturated fat or protein are equally effective at promoting fat loss and improving blood lipids. *Am J Clin Nutr* 2005;81:762-72.

Ma Y, Olendzki B, Chiriboga D, Hebert JR, Li Y, Li W, et al. Association between dietary carbohydrates and body weight. *Am J Epidemiol* 2005;161:359-67.

Ma Y, Olendzki BC, Chiriboga D, Rosal M, Sinagra E, Crawford S, et al. PDA-assisted low glycemic index dietary intervention for type II diabetes: A pilot study. *European Journal of Clinical Nutrition* 2006;60:1235-1243.

Ma Y, Olendzki BC, Merriam PA, Chiriboga DE, Culver AL, Li W, et al. A randomized clinical trial comparing low-glycemic index versus ADA dietary education among individuals with type 2 diabetes. *Nutrition* 2008;24:45-56.

Macdiarmid JI, Cade JE, Blundell JE. High and low fat consumers, their macronutrient intake and body mass index: further analysis of the National Diet and Nutrition Survey of British Adults. *Eur J Clin Nutr* 1996;50:505-12.

Maclure KM, Hayes KC, Colditz GA, Stampfer MJ, Speizer FE, Willett WC. Weight, diet, and the risk of symptomatic gallstones in middle-aged women. *N Engl J Med* 1989;321:563-9.

Maclure M, Travis LB, Willett W, MacMahon B. A prospective cohort study of nutrient intake and age at menarche. *Am J Clin Nutr*. 1991;54:649-56.

Maki KC, Rains TM, Kaden VN, Raneri KR, Davidson MH. Effects of a reduced-glycemic-load diet on body weight, body composition, and cardiovascular disease risk markers in overweight and obese adults. *Am J Clin Nutr* 2007;85:724-34.

Mann JI, Appleby PN, Key TJ, Thorogood M. Dietary determinants of ischaemic heart disease in health conscious individuals. *Heart* 1997;78:450-5.

Marniemi J, Seppanen A, Hakala P. Long-term effects on lipid metabolism of weight reduction on lactovegetarian and mixed diet. *Int J Obes* 1990;14:113-25.

Martin CK, Rosenbaum D, Han H, Geiselman PJ, Wyatt HR, Hill JO, et al. Change in food cravings, food preferences, and appetite during a low-carbohydrate and low-fat diet. *Obesity (Silver Spring)* 2011;19:1963-70.

Martinez-Gonzalez MA, Garcia-Lopez M, Bes-Rastrollo M, Toledo E, Martinez-Lapiscina EH, Delgado-Rodriguez M, et al. Mediterranean diet and the incidence of cardiovascular disease: a Spanish cohort. *Nutr Metab Cardiovasc Dis* 2011;21:237-44.

Maruthur NM, Wang NY, Appel LJ. Lifestyle interventions reduce coronary heart disease risk: results from the PREMIER Trial. *Circulation* 2009;119:2026-31.

Masala G, Ceroti M, Pala V, Krogh V, Vineis P, Sacerdote C, et al. A dietary pattern rich in olive oil and raw vegetables is associated with lower mortality in Italian elderly subjects. *Br J Nutr* 2007;98:406-15.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Masheb RM, Grilo CM, Rolls BJ. A randomized controlled trial for obesity and binge eating disorder: Low-energy-density dietary counseling and cognitive-behavioral therapy. *Behaviour Research and Therapy*;49:821-829.

Matsuo T, Kato Y, Murotake Y, Kim MK, Unno H, Tanaka K. An increase in high-density lipoprotein cholesterol after weight loss intervention is associated with longterm maintenance of reduced visceral abdominal fat. *International Journal of Obesity*;34:1742-1751.

Mayer-Davis EJ, Sparks KC, Hirst K, Costacou T, Lovejoy JC, Regensteiner JG, et al. Dietary intake in the diabetes prevention program cohort: baseline and 1-year post randomization. *Ann Epidemiol*. 2004;14:763-72.

Mazaraki A, Tsioufis C, Dimitriadis K, Tsiachris D, Stefanadi E, Zampelas A, et al. Adherence to the Mediterranean diet and albuminuria levels in Greek adolescents: data from the Leontio Lyceum ALbuminuria (3L study). *Eur J Clin Nutr*. 2011;65:219-25.

McAuley KA, Smith KJ, Taylor RW, McLay RT, Williams SM, Mann JI. Long-term effects of popular dietary approaches on weight loss and features of insulin resistance. *Int J Obes (Lond)* 2006;30:342-9.

McCarl M, Harnack L, Limburg PJ, Anderson KE, Folsom AR. Incidence of colorectal cancer in relation to glycemic index and load in a cohort of women. *Cancer Epidemiol Biomarkers Prev* 2006;15:892-6.

McClernon FJ, Yancy WS, Jr., Eberstein JA, Atkins RC, Westman EC. The effects of a low-carbohydrate ketogenic diet and a low-fat diet on mood, hunger, and other self-reported symptoms. *Obesity (Silver Spring)* 2007;15:182-7.

McConahy KL, Smiciklas-Wright H, Birch LL, Mitchell DC, Picciano MF. Food portions are positively related to energy intake and body weight in early childhood. *J Pediatr* 2002;140:340-7.

McCullough ML, Feskanich D, Stampfer MJ, Rosner BA, Hu FB, Hunter DJ, et al. Adherence to the Dietary Guidelines for Americans and risk of major chronic disease in women. *Am J Clin Nutr*. 2000;72:1214-22.

McDoniel SO, Hammond RS. A 24-week randomised controlled trial comparing usual care and metabolic-based diet plans in obese adults. *Int J Clin Pract* 2010;64:1503-11.

McGuire HL, Svetkey LP, Harsha DW, Elmer PJ, Appel LJ, Ard JD. Comprehensive lifestyle modification and blood pressure control: a review of the PREMIER trial. *J Clin Hypertens (Greenwich)* 2004;6:383-90.

McKeown NM, Meigs JB, Liu S, Rogers G, Yoshida M, Saltzman E, et al. Dietary carbohydrates and cardiovascular disease risk factors in the Framingham offspring cohort. *J Am Coll Nutr*. 2009;28:150-8.

McLaughlin T, Carter S, Lamendola C, Abbasi F, Schaaf P, Basina M, et al. Clinical efficacy of two hypocaloric diets that vary in overweight patients with type 2 diabetes: comparison of moderate fat versus carbohydrate reductions. *Diabetes Care* 2007;30:1877-9.

McManus K, Antinoro L, Sacks F. A randomized controlled trial of a moderate-fat, low-energy diet compared with a low fat, low-energy diet for weight loss in overweight adults. *Int J Obes Relat Metab*

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Disord 2001;25:1503-11.

McMillan-Price J, Petocz P, Atkinson F, O'Neill K, Samman S, Steinbeck K, et al. Comparison of 4 diets of varying glycemic load on weight loss and cardiovascular risk reduction in overweight and obese young adults: a randomized controlled trial. *Arch Intern Med* 2006;166:1466-75.

McTiernan A, Wactawski-Wende J, Wu L, Rodabough RJ, Watts NB, Tylavsky F, et al. Low-fat, increased fruit, vegetable, and grain dietary pattern, fractures, and bone mineral density: the Women's Health Initiative Dietary Modification Trial. *Am J Clin Nutr* 2009;89:1864-76.

Meinhold CL, Dodd KW, Jiao L, Flood A, Shikany JM, Genkinger JM, et al. Available carbohydrates, glycemic load, and pancreatic cancer: is there a link? *Am J Epidemiol*. 2010;171:1174-82.

Melander A. Review of previous impaired glucose tolerance intervention studies. *Diabet Med* 1996;13:S20-2.

Melanson EL, Astrup A, Donahoo WT. The relationship between dietary fat and fatty acid intake and body weight, diabetes, and the metabolic syndrome. *Annals of Nutrition and Metabolism* 2009;55:229-243.

Mendez MA, Popkin BM, Jakszyn P, Berenguer A, Tormo MJ, Sanchez MJ, et al. Adherence to a Mediterranean diet is associated with reduced 3-year incidence of obesity. *J Nutr* 2006;136:2934-8.

Messier SP, Gutekunst DJ, Davis C, DeVita P. Weight loss reduces knee-joint loads in overweight and obese older adults with knee osteoarthritis. *Arthritis Rheum* 2005;52:2026-32.

Metz JA, Stern JS, Kris-Etherton P, Reusser ME, Morris CD, Hatton DC, et al. A randomized trial of improved weight loss with a prepared meal plan in overweight and obese patients: impact on cardiovascular risk reduction. *Arch Intern Med* 2000;160:2150-8.

Meyer HE, Pedersen JI, Loken EB, Tverdal A. Dietary factors and the incidence of hip fracture in middle-aged Norwegians. A prospective study. *Am J Epidemiol* 1997;145:117-23.

Meyer J, Doring A, Herder C, Roden M, Koenig W, Thorand B. Dietary patterns, subclinical inflammation, incident coronary heart disease and mortality in middle-aged men from the MONICA/KORA Augsburg cohort study. *Eur J Clin Nutr* 2011;65:800-7.

Micha R, Mozaffarian D. Saturated fat and cardiometabolic risk factors, coronary heart disease, stroke, and diabetes: A fresh look at the evidence. *Lipids* 2010;45:893-905.

Michaud DS, Fuchs CS, Liu S, Willett WC, Colditz GA, Giovannucci E. Dietary glycemic load, carbohydrate, sugar, and colorectal cancer risk in men and women. *Cancer Epidemiol Biomarkers Prev*. 2005;14:138-47.

Michels KB, Fuchs CS, Giovannucci E, Colditz GA, Hunter DJ, Stampfer MJ, et al. Fiber intake and incidence of colorectal cancer among 76,947 women and 47,279 men. *Cancer Epidemiol Biomarkers Prev* 2005;14:842-9.

Mikkila V, Rasanen L, Raitakari OT, Pietinen P, Viikari J. Longitudinal changes in diet from childhood into adulthood with respect to risk of cardiovascular diseases: The Cardiovascular Risk in Young Finns

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Study. *Eur J Clin Nutr*. 2004;58:1038-45.

Miller GD, Rejeski WJ, Williamson JD, Morgan T, Sevick MA, Loeser RF, et al. The Arthritis, Diet and Activity Promotion Trial (ADAPT): design, rationale, and baseline results. *Control Clin Trials* 2003;24:462-80.

Mitrou PN, Kipnis V, Thiebaut AC, Reedy J, Subar AF, Wirfalt E, et al. Mediterranean dietary pattern and prediction of all-cause mortality in a US population: results from the NIH-AARP Diet and Health Study. *Arch Intern Med* 2007;167:2461-8.

Montonen J, Knekt P, Jarvinen R, Aromaa A, Reunanen A. Whole-grain and fiber intake and the incidence of type 2 diabetes. *Am J Clin Nutr* 2003;77:622-9.

Moore C, Gitau R, Goff L, Lewis FJ, Griffin MD, Chatfield MD, et al. Successful manipulation of the quality and quantity of fat and carbohydrate consumed by free-living individuals using a food exchange model. *Journal of Nutrition* 2009;139:1534-1540.

Moore CS, Lindroos AK, Kreutzer M, Larsen TM, Astrup A, van Baak MA, et al. Dietary strategy to manipulate ad libitum macronutrient intake, and glycaemic index, across eight European countries in the Diogenes Study. *Obes Rev* 2010;11:67-75.

Moretti L, Canada T. A randomized study comparing the effects of a low-carbohydrate diet and a conventional diet on lipoprotein subfractions and C-reactive protein levels in patients with severe obesity: Commentary. *Nutrition in Clinical Practice* 2006;21:187-188.

Morgan LM, Griffin BA, Millward DJ, DeLooy A, Fox KR, Baic S, et al. Comparison of the effects of four commercially available weight-loss programmes on lipid-based cardiovascular risk factors. *Public Health Nutr* 2009;12:799-807.

Morgan PJ, Lubans DR, Collins CE, Warren JM, Callister R. 12-month outcomes and process evaluation of the SHED-IT RCT: an internet-based weight loss program targeting men. *Obesity (Silver Spring)* 2011;19:142-51.

Mozaffarian D, Aro A, Willett WC. Health effects of trans-fatty acids: Experimental and observational evidence. *European Journal of Clinical Nutrition* 2009;63:S5-S21.

Mozaffarian D, Cao H, King IB, Lemaitre RN, Song X, Siscovick DS, et al. Trans-palmitoleic acid, metabolic risk factors, and new-onset diabetes in U.S. adults: a cohort study. *Ann Intern Med* 2010;153:790-9.

Mueller-Cunningham WM, Quintana R, Kasim-Karakas SE. An ad libitum, very low-fat diet results in weight loss and changes in nutrient intakes in postmenopausal women. *J Am Diet Assoc* 2003;103:1600-6.

Munger RG, Cerhan JR, Chiu BC. Prospective study of dietary protein intake and risk of hip fracture in postmenopausal women. *Am J Clin Nutr* 1999;69:147-52.

Musunuru K. Atherogenic dyslipidemia: Cardiovascular risk and dietary intervention. *Lipids* 2010;45:907-914.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Nagle CM, Kolahdooz F, Ibiebele TI, Olsen CM, Lahmann PH, Green AC, et al. Carbohydrate intake, glycemic load, glycemic index, and risk of ovarian cancer. *Ann Oncol* 2011;22:1332-8.

Ness AR. The Avon Longitudinal Study of Parents and Children (ALSPAC)--a resource for the study of the environmental determinants of childhood obesity. *Eur J Endocrinol* 2004;151 Suppl 3:U141-9.

Neuhouser ML, Barnett MJ, Kristal AR, Ambrosone CB, King I, Thornquist M, et al. (n-6) PUFA increase and dairy foods decrease prostate cancer risk in heavy smokers. *J Nutr* 2007;137:1821-7.

Neuhouser ML, Howard B, Lu J, Tinker LF, Van Horn L, Caan B, et al. A low-fat dietary pattern and risk of metabolic syndrome in postmenopausal women: The Women's Health Initiative. *Metabolism: Clinical and Experimental* 2012;61:1572-1581.

Newby PK, Peterson KE, Berkey CS, Leppert J, Willett WC, Colditz GA. Dietary composition and weight change among low-income preschool children. *Arch Pediatr Adolesc Med.* 2003;157:759-64.

Newby PK, Tucker KL, Wolk A. Risk of overweight and obesity among semivegetarian, lactovegetarian, and vegan women. *Am J Clin Nutr* 2005;81:1267-74.

Nicklas BJ, Katzel LI, Bunyard LB, Dennis KE, Goldberg AP. Effects of an American Heart Association diet and weight loss on lipoprotein lipids in obese, postmenopausal women. *Am J Clin Nutr* 1997;66:853-9.

Nield L, Moore H, Hooper L, Cruickshank K, Vyas A, Whittaker V, et al. Dietary advice for treatment of type 2 diabetes mellitus in adults. *Cochrane Database of Systematic Reviews* 2007.

Nielsen JV, Jonsson E, Nilsson AK. Lasting improvement of hyperglycaemia and bodyweight: low-carbohydrate diet in type 2 diabetes--a brief report. *Ups J Med Sci* 2005;110:69-73.

Nilsson LM, Winkvist A, Eliasson M, Jansson JH, Hallmans G, Johansson I, et al. Low-carbohydrate, high-protein score and mortality in a northern Swedish population-based cohort. *Eur J Clin Nutr* 2012;66:694-700.

Nkondjock A, Robidoux A, Paredes Y, Narod SA, Ghadirian P. Diet, lifestyle and BRCA-related breast cancer risk among French-Canadians. *Breast Cancer Res Treat.* 2006;98:285-94.

Nobels F, van Gaal L, de Leeuw I. Weight reduction with a high protein, low carbohydrate, calorie-restricted diet: effects on blood pressure, glucose and insulin levels. *Neth J Med* 1989;35:295-302.

Noble RE. A controlled study of a weight reduction regimen. *Curr Ther Res Clin Exp* 1971;13:685-91.

Nomura AMY, Hankin JH, Henderson BE, Wilkens LR, Murphy SP, Pike MC, et al. Dietary fiber and colorectal cancer risk: The multiethnic cohort study. *Cancer Causes and Control* 2007;18:753-764.

Nordmann AJ, Nordmann A, Briel M, Keller U, Yancy WS, Jr., Brehm BJ, et al. Effects of low-carbohydrate vs low-fat diets on weight loss and cardiovascular risk factors: a meta-analysis of randomized controlled trials. *Arch Intern Med* 2006;166:285-93.

Nystrom FH, Ostgren CJ, Lindstrom T, Bachrach-Lindstrom M, Schold AK, Dizdar B, et al. A high fat diet improves glycaemic control compared with low fat diet: A 24-month randomised prospective study of patients with type 2 diabetes in primary health care. *SOURCE Diabetologia* (2011) 54 SUPPL.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

1 (S358). Date of Publication: September 2011 CONFERENCE NAME 47th Annual Meeting of the European Association for the Study of Diabetes, EASD 2011 CONFERENCE LOCATION Lisbon, Portugal CONFERENCE DATE 2011-09-12 to 2011-09-16.

Obarzanek E, Proschan MA, Vollmer WM, Moore TJ, Sacks FM, Appel LJ, et al. Individual blood pressure responses to changes in salt intake: Results from the DASH-sodium trial. *Hypertension* 2003;42:459-467.

Ochner CN, Lowe MR. Self-reported changes in dietary calcium and energy intake predict weight regain following a weight loss diet in obese women. *J Nutr* 2007;137:2324-8.

Oh K, Hu FB, Cho E, Rexrode KM, Stampfer MJ, Manson JE, et al. Carbohydrate intake, glycemic index, glycemic load, and dietary fiber in relation to risk of stroke in women. *Am J Epidemiol*. 2005;161:161-9.

Oh K, Willett WC, Fuchs CS, Giovannucci EL. Glycemic index, glycemic load, and carbohydrate intake in relation to risk of distal colorectal adenoma in women. *Cancer Epidemiol Biomarkers Prev*. 2004;13:1192-8.

Osler M, Heitmann BL, Gerdes LU, Jorgensen LM, Schroll M. Dietary patterns and mortality in Danish men and women: a prospective observational study. *Br J Nutr* 2001;85:219-25.

Paeratakul S, Popkin BM, Keyou G, Adair LS, Stevens J. Changes in diet and physical activity affect the body mass index of Chinese adults. *Int J Obes Relat Metab Disord*. 1998;22:424-31.

Paineau DL, Beaufile F, Boulier A, Cassuto DA, Chwalow J, Combris P, et al. Family dietary coaching to improve nutritional intakes and body weight control: a randomized controlled trial. *Arch Pediatr Adolesc Med* 2008;162:34-43.

Panagiotakos DB, Pitsavos C, Chrysohoou C, Skoumas I, Stefanadis C. Five-year incidence of cardiovascular disease and its predictors in Greece: the ATTICA study. *Vasc Med* 2008;13:113-21.

Papadaki A, Linardakis M, Larsen TM, van Baak MA, Lindroos AK, Pfeiffer AF, et al. The effect of protein and glycemic index on children's body composition: the DiOGenes randomized study. *Pediatrics* 2010;126:e1143-52.

Papadaki A, Linardakis M, Larsen TM, van Baak MA, Lindroos AK, Pfeiffer AF, et al. The effect of protein and glycemic index on children's body composition: the DiOGenes randomized study. *Pediatrics* 2010;126:e1143-52.

Partsalaki I, Karvela A, Spiliotis BE. Metabolic impact of a ketogenic diet compared to a hypocaloric diet in obese children and adolescents. *Journal of Pediatric Endocrinology and Metabolism* 2012;25:697-704.

Partsalaki I, Karvela A, Spiliotis BE. Metabolic impact of a ketogenic diet compared to a hypocaloric diet in obese children and adolescents. *Journal of Pediatric Endocrinology and Metabolism* 2012;25:697-704.

Pascale RW, Wing RR, Butler BA, Mullen M, Bononi P. Effects of a behavioral weight loss program stressing calorie restriction versus calorie plus fat restriction in obese individuals with NIDDM or a

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

family history of diabetes. *Diabetes Care* 1995;18:1241-8.

Pearce ML, Dayton S. Incidence of cancer in men on a diet high in polyunsaturated fat. *Lancet* 1971;1:464-7.

Pereira MA, Swain J, Goldfine AB, Rifai N, Ludwig DS. Effects of a low-glycemic load diet on resting energy expenditure and heart disease risk factors during weight loss. *JAMA* 2004;292:2482-90.

Perez-Martinez P, Moreno-Conde M, Cruz-Teno C, Ruano J, Fuentes F, Delgado-Lista J, et al. Dietary fat differentially influences regulatory endothelial function during the postprandial state in patients with metabolic syndrome: From the LIPGENE study. *Atherosclerosis* 2010;209:533-538.

Philippou E, Bovill-Taylor C, Rajkumar C, Vampa ML, Ntatsaki E, Brynes AE, et al. Preliminary report: the effect of a 6-month dietary glycemic index manipulation in addition to healthy eating advice and weight loss on arterial compliance and 24-hour ambulatory blood pressure in men: a pilot study. *Metabolism* 2009;58:1703-8.

Pieper C, Redman L, Racette S, Roberts S, Bhapkar M, Rochon J, et al. Development of adherence metrics for caloric restriction interventions. *Clinical Trials* 2011;8:155-164.

Pittas AG, Das SK, Hajduk CL, Golden J, Saltzman E, Stark PC, et al. A low-glycemic load diet facilitates greater weight loss in overweight adults with high insulin secretion but not in overweight adults with low insulin secretion in the CALERIE Trial. *Diabetes Care* 2005;28:2939-41.

Pittas AG, Roberts SB, Das SK, Gilhooly CH, Saltzman E, Golden J, et al. The effects of the dietary glycemic load on type 2 diabetes risk factors during weight loss. *Obesity (Silver Spring)* 2006;14:2200-9.

Poppitt SD, Keogh GF, Prentice AM, Williams DE, Sonnemans HM, Valk EE, et al. Long-term effects of ad libitum low-fat, high-carbohydrate diets on body weight and serum lipids in overweight subjects with metabolic syndrome. *Am J Clin Nutr* 2002;75:11-20.

Poppitt SD, Prentice AM. Energy density and its role in the control of food intake: evidence from metabolic and community studies. *Appetite* 1996;26:153-74.

Prentice RL, Caan B, Chlebowski RT, Patterson R, Kuller LH, Ockene JK, et al. Low-fat dietary pattern and risk of invasive breast cancer: the Women's Health Initiative Randomized Controlled Dietary Modification Trial. *JAMA* 2006;295:629-42.

Prentice RL, Thomson CA, Caan B, Hubbell FA, Anderson GL, Beresford SA, et al. Low-fat dietary pattern and cancer incidence in the Women's Health Initiative Dietary Modification Randomized Controlled Trial. *J Natl Cancer Inst* 2007;99:1534-43.

Pritchard JE, Nowson CA, Wark JD. Bone loss accompanying diet-induced or exercise-induced weight loss: a randomised controlled study. *Int J Obes Relat Metab Disord* 1996;20:513-20.

Psaltopoulou T, Kyrozis A, Stathopoulos P, Trichopoulos D, Vassilopoulos D, Trichopoulou A. Diet, physical activity and cognitive impairment among elders: the EPIC-Greece cohort (European Prospective Investigation into Cancer and Nutrition). *Public Health Nutr*. 2008;11:1054-62.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Qi L, Meigs JB, Liu S, Manson JE, Mantzoros C, Hu FB. Dietary fibers and glycemic load, obesity, and plasma adiponectin levels in women with type 2 diabetes. *Diabetes Care* 2006;29:1501-5.

Qi L, van Dam RM, Rexrode K, Hu FB. Heme iron from diet as a risk factor for coronary heart disease in women with type 2 diabetes. *Diabetes Care* 2007;30:101-6.

Qi Q, Bray GA, Smith SR, Hu FB, Sacks FM, Qi L. Insulin Receptor Substrate 1 Gene Variation Modifies Insulin Resistance Response to Weight-Loss Diets in a 2-Year Randomized Trial: The Preventing Overweight Using Novel Dietary Strategies (POUNDS LOST) Trial. *Circulation*;124:563-571.

Qi Q, Hu FB, Sacks FM, Qi L. Common variant near GIPR gene modifies fasting glucose and related traits response to weight-loss diets in a two-year randomized trial. *SOURCE Diabetes* (2011) 60 SUPPL. 1 (A32). Date of Publication: July 2011 CONFERENCE NAME 71st Scientific Sessions of the American Diabetes Association CONFERENCE LOCATION San Diego, CA, United States CONFERENCE DATE 2011-06-24 to 2011-06-28.

Raatz SK, Torkelson CJ, Redmon JB, Reck KP, Kwong CA, Swanson JE, et al. Reduced glycemic index and glycemic load diets do not increase the effects of energy restriction on weight loss and insulin sensitivity in obese men and women. *J Nutr* 2005;135:2387-91.

Randi G, Ferraroni M, Talamini R, Garavello W, Deandrea S, Decarli A, et al. Glycemic index, glycemic load and thyroid cancer risk. *Ann Oncol.* 2008;19:380-3.

Rapuri PB, Gallagher JC, Haynatzka V. Protein intake: effects on bone mineral density and the rate of bone loss in elderly women. *Am J Clin Nutr.* 2003;77:1517-25.

Raynor H, Anderson E, Hecht J, Wing RR. Does limiting variety in snack foods assist with reducing energy intake and weight loss during a lifestyle intervention? *SOURCE Obesity* (2011) 19 SUPPL. 1 (S62). Date of Publication: November 2011 CONFERENCE NAME 29th Annual Scientific Meeting of the Obesity Society, Obesity 2011 CONFERENCE LOCATION Orlando, FL, United States CONFERENCE DATE 2011-10-01 to 2011-10-05.

Raynor HA, Jeffery RW, Ruggiero AM, Clark JM, Delahanty LM. Weight loss strategies associated with BMI in overweight adults with type 2 diabetes at entry into the Look AHEAD (Action for Health in Diabetes) trial. *Diabetes Care* 2008;31:1299-304.

Raynor HA, Looney SM, Steeves EA, Spence M, Gorin AA. The effects of an energy density prescription on diet quality and weight loss: a pilot randomized controlled trial. *J Acad Nutr Diet* 2012;112:1397-402.

Raynor HA, Van Walleghen EL, Bachman JL, Looney SM, Phelan S, Wing RR. Dietary energy density and successful weight loss maintenance. *Eating Behaviors* 2011;12:119-125.

Raynor HA, Van Walleghen EL, Bachman JL, Looney SM, Phelan S, Wing RR. Dietary energy density and successful weight loss maintenance. *Eat Behav* 2011;12:119-25.

Razquin C, Alfredo Martinez J, Martinez-Gonzalez MA, Corella D, Santos JM, Marti A. The Mediterranean diet protects against waist circumference enlargement in 12Ala carriers for the PPARgamma gene: 2 years' follow-up of 774 subjects at high cardiovascular risk. *Br J Nutr.* 2009;102:672-9.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Razquin C, Martinez JA, Martinez-Gonzalez MA, Bes-Rastrollo M, Fernandez-Crehuet J, Marti A. A 3-year intervention with a Mediterranean diet modified the association between the rs9939609 gene variant in FTO and body weight changes. *Int J Obes (Lond)* 2010;34:266-72.

Razquin C, Martinez JA, Martinez-Gonzalez MA, Mitjavila MT, Estruch R, Marti A. A 3 years follow-up of a Mediterranean diet rich in virgin olive oil is associated with high plasma antioxidant capacity and reduced body weight gain. *Eur J Clin Nutr* 2009;63:1387-93.

Rebholz CM, Friedman EE, Powers LJ, Arroyave WD, He J, Kelly TN. Dietary protein intake and blood pressure: A meta-analysis of randomized controlled trials. *American Journal of Epidemiology* 2012;176:S27-S43.

Redman LM, Heilbronn LK, Martin CK, Alfonso A, Smith SR, Ravussin E. Effect of calorie restriction with or without exercise on body composition and fat distribution. *J Clin Endocrinol Metab* 2007;92:865-72.

Reinehr T, Kleber M, Toschke AM. Lifestyle intervention in obese children is associated with a decrease of the metabolic syndrome prevalence. *Atherosclerosis* 2009:174-80.

Reinehr T, Schmidt C, Toschke AM, Andler W. Lifestyle intervention in obese children with nonalcoholic fatty liver disease: 2-Year follow-up study. *Archives of Disease in Childhood* 2009;94:437-442.

Rejeski WJ, Focht BC, Messier SP, Morgan T, Pahor M, Penninx B. Obese, older adults with knee osteoarthritis: weight loss, exercise, and quality of life. *Health Psychol* 2002;21:419-26.

Riebe D, Blissmer B, Greene G, Caldwell M, Ruggiero L, Stillwell KM, et al. Long-term maintenance of exercise and healthy eating behaviors in overweight adults. *Prev Med* 2005;40:769-78.

Roderick P. A randomized trial to evaluate the effectiveness of dietary advice by practice nurses in lowering diet-related coronary heart disease risk. *British Journal of General Practice* 1997;47:7-12.

Rodriguez-Hernandez H, Cervantes-Huerta M, Rodriguez-Moran M, Guerrero-Romero F. Decrease of aminotransferase levels in obese women is related to body weight reduction, irrespective of type of diet. *Ann Hepatol* 2011;10:486-92.

Rodriguez-Hernandez H, Morales-Amaya UA, Rosales-Valdez R, Rivera-Hinojosa F, Rodriguez-Moran M, Guerrero-Romero F. Adding cognitive behavioural treatment to either low-carbohydrate or low-fat diets: differential short-term effects. *Br J Nutr* 2009;102:1847-53.

Rohan TE, Howe GR, Friedenreich CM, Jain M, Miller AB. Dietary fiber, vitamins A, C, and E, and risk of breast cancer: a cohort study. *Cancer Causes Control*. 1993;4:29-37.

Rolland C, Hession M, Murray S, Wise A, Broom I. Randomized clinical trial of standard dietary treatment versus a low-carbohydrate/high-protein diet or the LighterLife Programme in the management of obesity*. *J Diabetes* 2009;1:207-17.

Rolland-Cachera MF, Thibault H, Souberbielle JC, Soulie D, Carbonel P, Deheeger M, et al. Massive obesity in adolescents: dietary interventions and behaviours associated with weight regain at 2 y follow-up. *Int J Obes Relat Metab Disord* 2004;28:514-9.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Rolls BJ, Roe LS, Beach AM, Kris-Etherton PM. Provision of foods differing in energy density affects long-term weight loss. *Obes Res* 2005;13:1052-60.

Romaguera D, Norat T, Mouw T, May AM, Bamia C, Slimani N, et al. Adherence to the Mediterranean diet is associated with lower abdominal adiposity in European men and women. *J Nutr* 2009;139:1728-37.

Romaguera D, Norat T, Vergnaud A, Mouw T, May AM, Agudo A, et al. Mediterranean dietary patterns and prospective weight change in participants of the EPIC-PANACEA project. *American Journal of Clinical Nutrition*;92:912-921.

Ros E. Dietary cis-monounsaturated fatty acids and metabolic control in type 2 diabetes. *Am J Clin Nutr* 2003;78:617S-625S.

Ross LJ, Tapsell LC, Probst Y. Optimizing dietary fat in a weight-loss trial requires advice based on a structured "whole-of-diet" model. *Nutr Res* 2011;31:683-90.

Ryttig KR, Rossner S. Weight maintenance after a very low calorie diet (VLCD) weight reduction period and the effects of VLCD supplementation. A prospective, randomized, comparative, controlled long-term trial. *J Intern Med* 1995;238:299-306.

Sacks FM, Bray GA, Carey VJ, Smith SR, Ryan DH, Anton SD, et al. Comparison of weight-loss diets with different compositions of fat, protein, and carbohydrates. *N Engl J Med* 2009;360:859-73.

Sacks FM, Bray GA, Carey VJ, Smith SR, Ryan DH, Anton SD, et al. The new England journal of medicine: Comparison of weight-loss diets with different compositions of fat, protein, and carbohydrates. *New England Journal of Medicine* 2009;360:859-873.

Sahni S, Cupples LA, McLean RR, Tucker KL, Broe KE, Kiel DP, et al. Protective effect of high protein and calcium intake on the risk of hip fracture in the Framingham offspring cohort. *J Bone Miner Res* 2010;25:2494-500.

Salamone LM, Cauley JA, Black DM, Simkin-Silverman L, Lang W, Gregg E, et al. Effect of a lifestyle intervention on bone mineral density in premenopausal women: a randomized trial. *Am J Clin Nutr* 1999;70:97-103.

Salas-Salvado J, Bullo M, Babio N, Martinez-Gonzalez MA, Ibarrola-Jurado N, Basora J, et al. Reduction in the incidence of type 2 diabetes with the mediterranean diet: Results of the PREDIMED-Reus nutrition intervention randomized trial. *Diabetes Care* 2011;34:14-19.

Salas-Salvado J, Fernandez-Ballart J, Ros E, Martinez-Gonzalez MA, Fito M, Estruch R, et al. Effect of a Mediterranean diet supplemented with nuts on metabolic syndrome status: one-year results of the PREDIMED randomized trial. *Arch Intern Med* 2008;168:2449-58.

Salmeron J, Manson JE, Stampfer MJ, Colditz GA, Wing AL, Willett WC. Dietary fiber, glycemic load, and risk of non-insulin-dependent diabetes mellitus in women. *JAMA* 1997;277:472-7.

Sanchez-Tainta A, Estruch R, Bullo M, Corella D, Gomez-Gracia E, Fiol M, et al. Adherence to a Mediterranean-type diet and reduced prevalence of clustered cardiovascular risk factors in a cohort of 3,204 high-risk patients. *Eur J Cardiovasc Prev Rehabil* 2008;15:589-93.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Sanchez-Villegas A, Delgado-Rodriguez M, Alonso A, Schlatter J, Lahortiga F, Serra Majem L, et al. Association of the Mediterranean dietary pattern with the incidence of depression: the Seguimiento Universidad de Navarra/University of Navarra follow-up (SUN) cohort. *Arch Gen Psychiatry*. 2009;66:1090-8.

Sanchez-Villegas A, Galbete C, Martinez-Gonzalez MA, Martinez JA, Razquin C, Salas-Salvado J, et al. The effect of the Mediterranean diet on plasma brain-derived neurotrophic factor (BDNF) levels: the PREDIMED-NAVARRA randomized trial. *Nutr Neurosci* 2011;14:195-201.

Santos FL, Esteves SS, da Costa Pereira A, Yancy Jr WS, Nunes JPL. Systematic review and meta-analysis of clinical trials of the effects of low carbohydrate diets on cardiovascular risk factors. *Obesity Reviews* 2012;13:1048-1066.

Saris WH, Astrup A, Prentice AM, Zunft HJ, Formiguera X, Verboeket-van de Venne WP, et al. Randomized controlled trial of changes in dietary carbohydrate/fat ratio and simple vs complex carbohydrates on body weight and blood lipids: the CARMEN study. The Carbohydrate Ratio Management in European National diets. *Int J Obes Relat Metab Disord* 2000;24:1310-8.

Saris WHM, Astrup A, Prentice AM, Zunft HJF, Formiguera X, Verboeket-Van De Venne WPHG, et al. Randomized controlled trial of changes in dietary carbohydrate/fat ratio and simple vs complex carbohydrates on body weight and blood lipids: The CARMEN study. *International Journal of Obesity* 2000;24:1310-1318.

Sarwer DB, Wadden TA, Moore RH, Baker AW, Gibbons LM, Raper SE, et al. Preoperative eating behavior, postoperative dietary adherence, and weight loss after gastric bypass surgery. *Surg Obes Relat Dis* 2008;4:640-6.

Sarzi-Puttini P, Comi D, Boccassini L, Muzzupappa S, Turiel M, Panni B, et al. Diet therapy for rheumatoid arthritis. A controlled double-blind study of two different dietary regimens. *Scand J Rheumatol* 2000;29:302-7.

Sauvaget C, Nagano J, Hayashi M, Yamada M. Animal protein, animal fat, and cholesterol intakes and risk of cerebral infarction mortality in the adult health study. *Stroke* 2004;35:1531-7.

Savoie M, Nowicka P, Shaw M, Yu S, Dziura J, Chavent G, et al. Long-term results of an obesity program in an ethnically diverse pediatric population. *Pediatrics* 2011;127:402-10.

Scavone G, Manto A, Pitocco D, Gagliardi L, Caputo S, Mancini L, et al. Effect of carbohydrate counting and medical nutritional therapy on glycaemic control in Type 1 diabetic subjects: A pilot study. *Diabetic Medicine* 2010;27:477-479.

Schlundt DG, Hill JO, Pope-Cordle J, Arnold D, Virts KL, Katahn M. Randomized evaluation of a low fat ad libitum carbohydrate diet for weight reduction. *Int J Obes Relat Metab Disord* 1993;17:623-9.

Schulz M, Nothlings U, Hoffmann K, Bergmann MM, Boeing H. Identification of a food pattern characterized by high-fiber and low-fat food choices associated with low prospective weight change in the EPIC-Potsdam cohort. *J Nutr* 2005;135:1183-9.

Schusdziarra V, Hausmann M, Wiedemann C, Hess J, Barth C, Wagenpfeil S, et al. Successful weight loss and maintenance in everyday clinical practice with an individually tailored change of eating

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

habits on the basis of food energy density. *Eur J Nutr* 2011;50:351-61.

Schwarzfuchs D, Golan R, Shai I. Four-year follow-up after two-year dietary interventions. *N Engl J Med* 2012;367:1373-4.

Sedlacek SM, Playdon MC, Wolfe P, McGinley JN, Wisthoff MR, Daeninck EA, et al. Effect of a low fat versus a low carbohydrate weight loss dietary intervention on biomarkers of long term survival in breast cancer patients ('CHOICE'): Study protocol. *SOURCE BMC Cancer* (2011) 11 Article Number: 287. Date of Publication: 6 Jul 2011.

Seshadri P, Iqbal N, Stern L, Williams M, Chicano KL, Daily DA, et al. A randomized study comparing the effects of a low-carbohydrate diet and a conventional diet on lipoprotein subfractions and C-reactive protein levels in patients with severe obesity. *Am J Med* 2004;117:398-405.

Seshadri P, Samaha FF, Stern L, Ahima RS, Daily D, Iqbal N. Adipocytokine changes caused by low-carbohydrate compared to conventional diets in obesity. *Metabolic Syndrome and Related Disorders* 2005;3:66-74.

Seshadri P, Samaha FF, Stern L, Chicano KL, Daily DA, Iqbal N. Free fatty acids, insulin resistance, and corrected qt intervals in morbid obesity: effect of weight loss during 6 months with differing dietary interventions. *Endocr Pract* 2005;11:234-9.

Shah M, Jeffery RW, Laing B, Savre SG, Van Natta M, Strickland D. Hypertension Prevention Trial (HPT): food pattern changes resulting from intervention on sodium, potassium, and energy intake. Hypertension Prevention Trial Research Group. *J Am Diet Assoc* 1990;90:69-76.

Shai I, Schwarzfuchs D, Henkin Y, Shahar DR, Witkow S, Greenberg I, et al. Weight loss with a low-carbohydrate, Mediterranean, or low-fat diet. *N Engl J Med* 2008;359:229-41.

Shaw DI, Tierney AC, McCarthy S, Upritchard J, Vermunt S, Gulseth HL, et al. LIPGENE food-exchange model for alteration of dietary fat quantity and quality in free-living participants from eight European countries. *British Journal of Nutrition* 2009;101:750-759.

Shea MK, Houston DK, Nicklas BJ, Messier SP, Davis CC, Miller ME, et al. The effect of randomization to weight loss on total mortality in older overweight and obese adults: the ADAPT Study. *J Gerontol A Biol Sci Med Sci* 2010;65:519-25.

Shekelle RB, Shryock AM, Paul O, Lepper M, Stamler J, Liu S, et al. Diet, serum cholesterol, and death from coronary heart disease. The Western Electric study. *N Engl J Med* 1981;304:65-70.

Sheppard L, Kristal AR, Kushi LH. Weight loss in women participating in a randomized trial of low-fat diets. *Am J Clin Nutr* 1991;54:821-8.

Shintani T, Beckham S, Tang J, O'Connor HK, Hughes C. Waianae Diet Program: long-term follow-up. *Hawaii Med J* 1999;58:117-22.

Sichieri R, Moura AS, Genelhu V, Hu F, Willett WC. An 18-mo randomized trial of a low-glycemic-index diet and weight change in Brazilian women. *Am J Clin Nutr* 2007;86:707-13.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Sichieri R. Is fat intake important in the public health control of obesity? *Am J Clin Nutr* 2000;72:203-6.

Sieri S, Krogh V, Pala V, Muti P, Micheli A, Evangelista A, et al. Dietary patterns and risk of breast cancer in the ORDET cohort. *Cancer Epidemiol Biomarkers Prev* 2004;13:567-72.

Sieri S, Pala V, Brighenti F, Pellegrini N, Muti P, Micheli A, et al. Dietary glycemic index, glycemic load, and the risk of breast cancer in an Italian prospective cohort study. *Am J Clin Nutr*. 2007;86:1160-6.

Silvera SA, Jain M, Howe GR, Miller AB, Rohan TE. Dietary carbohydrates and breast cancer risk: a prospective study of the roles of overall glycemic index and glycemic load. *Int J Cancer*. 2005;114:653-8.

Silvera SA, Jain M, Howe GR, Miller AB, Rohan TE. Energy balance and breast cancer risk: a prospective cohort study. *Breast Cancer Res Treat*. 2006;97:97-106.

Silvera SA, Rohan TE, Jain M, Terry PD, Howe GR, Miller AB. Glycaemic index, glycaemic load and risk of endometrial cancer: a prospective cohort study. *Public Health Nutr* 2005;8:912-9.

Simell O, Niinikoski H, Ronnema T, Lapinleimu H, Routi T, Lagstrom H, et al. Special Turku Coronary Risk Factor Intervention Project for Babies (STRIP). *Am J Clin Nutr* 2000;72:1316S-1331S.

Simila ME, Valsta LM, Kontto JP, Albanes D, Virtamo J. Low-, medium- and high-glycaemic index carbohydrates and risk of type 2 diabetes in men. *Br J Nutr*. 2011;105:1258-64.

Simmons D, Rush E, Crook N. Development and piloting of a community health worker-based intervention for the prevention of diabetes among New Zealand Maori in Te Wai o Rona: Diabetes Prevention Strategy. *Public Health Nutr* 2008;11:1318-25.

Simon MS, Heilbrun LK, Boomer A, Kresge C, Depper J, Kim PN, et al. A randomized trial of a low-fat dietary intervention in women at high risk for breast cancer. *Nutr Cancer* 1997;27:136-42.

Sinclair AJ, Burdon MA, Nightingale PG, Ball AK, Good P, Matthews TD, et al. Low energy diet and intracranial pressure in women with idiopathic intracranial hypertension: Prospective cohort study. *BMJ* 2010;341:138.

Singh RB, Rastogi SS, Sircar AR, Mehta PJ, Sharma KK. Dietary strategies for risk-factor modification to prevent cardiovascular diseases. *Nutrition* 1991;7:210-4.

Singh RB, Rastogi SS, Verma R, Laxmi B, Singh R, Ghosh S, et al. Randomised controlled trial of cardioprotective diet in patients with recent acute myocardial infarction: results of one year follow up. *BMJ* 1992;304:1015-9.

Singh RB, Rastogi V, Rastogi SS, Niaz MA, Beegom R. Effect of diet and moderate exercise on central obesity and associated disturbances, myocardial infarction and mortality in patients with and without coronary artery disease. *Journal of the American College of Nutrition* 1996;15:592-601.

Skov AR, Haulrik N, Toubro S, Molgaard C, Astrup A. Effect of protein intake on bone mineralization during weight loss: a 6-month trial. *Obes Res* 2002;10:432-8.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Skov AR, Toubro S, Bulow J, Krabbe K, Parving HH, Astrup A. Changes in renal function during weight loss induced by high vs low-protein low-fat diets in overweight subjects. *Int J Obes Relat Metab Disord* 1999;23:1170-7.

Skov AR, Toubro S, Ronn B, Holm L, Astrup A. Randomized trial on protein vs carbohydrate in ad libitum fat reduced diet for the treatment of obesity. *Int J Obes Relat Metab Disord* 1999;23:528-36.

Sloth B, Due A, Larsen TM, Holst JJ, Hedning A, Astrup A. The effect of a high-MUFA, low-glycaemic index diet and a low-fat diet on appetite and glucose metabolism during a 6-month weight maintenance period. *Br J Nutr* 2009;101:1846-58.

Sluijs I, Beulens JW, van der AD, Spijkerman AM, Grobbee DE, van der Schouw YT. Dietary intake of total, animal, and vegetable protein and risk of type 2 diabetes in the European Prospective Investigation into Cancer and Nutrition (EPIC)-NL study. *Diabetes Care* 2010;33:43-8.

Smit E, Garcia-Palmieri MR, Figueroa NR, McGee DL, Messina M, Freudenheim JL, et al. Protein and legume intake and prostate cancer mortality in Puerto Rican men. *Nutr Cancer* 2007;58:146-52.

Smith DE, Wing RR. Diminished weight loss and behavioral compliance during repeated diets in obese patients with type II diabetes. *Health Psychol* 1991;10:378-83.

Sofer S, Eliraz A, Kaplan S, Voet H, Fink G, Kima T, et al. Greater weight loss and hormonal changes after 6 months diet with carbohydrates eaten mostly at dinner. *Obesity (Silver Spring)* 2011;19:2006-14.

Sofer S, Eliraz A, Kaplan S, Voet H, Fink G, Kima T, et al. Greater weight loss and hormonal changes after 6 months diet with carbohydrates eaten mostly at dinner. *Obesity (Silver Spring)* 2011;19:2006-14.

Sonnenberg L, Pencina M, Kimokoti R, Quatromoni P, Nam BH, D'Agostino R, et al. Dietary patterns and the metabolic syndrome in obese and non-obese Framingham women. *Obes Res* 2005;13:153-62.

Sothorn MS, von Almen TK, Schumacher H, Zelman M, Farris RP, Carlisle L, et al. An effective multidisciplinary approach to weight reduction in youth. *Ann N Y Acad Sci* 1993;699:292-4.

Stamler R, Stamler J, Gosch FC, McDonald AM. Primary prevention of hypertension--a randomized controlled trial. *Ann Clin Res* 1984;16 Suppl 43:136-42.

Stamler R, Stamler J, Grimm R. Nutritional therapy for high blood pressure. Final report of a four-year randomized controlled trial - The Hypertension Control Program. *Journal of the American Medical Association* 1987;257:1484-1491.

Stoll BA. Nutrition and breast cancer risk: can an effect via insulin resistance be demonstrated? *Breast Cancer Res Treat* 1996;38:239-46.

Streppel MT, Ocke MC, Boshuizen HC, Kok FJ, Kromhout D. Dietary fiber intake in relation to coronary heart disease and all-cause mortality over 40 y: the Zutphen Study. *Am J Clin Nutr* 2008;88:1119-25.

Strychar I. Diet in the management of weight loss. *CMAJ* 2006;174:56-63.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Sukumar D, Ambia-Sobhan H, Zurfluh R, Schlussek Y, Stahl T, Gordon C, et al. Areal and volumetric bone mineral density and geometry at two levels of protein intake during caloric restriction: a randomized controlled trial. *J Bone Miner Res* 2010.

Summer SS, Brehm BJ, Benoit SC, D'Alessio DA. Adiponectin changes in relation to the macronutrient composition of a weight-loss diet. *Obesity (Silver Spring)* 2011;19:2198-204.

Summerbell Carolyn D, Cameron C, Glasziou Paul P. Advice on low-fat diets for obesity. *Cochrane Database of Systematic Reviews* 2008.

Svetkey LP, Erlinger TP, Vollmer WM, Feldstein A, Cooper LS, Appel LJ, et al. Effect of lifestyle modifications on blood pressure by race, sex, hypertension status, and age. *J Hum Hypertens* 2005;19:21-31.

Swinburn BA, Metcalf PA, Ley SJ. Long-term (5-year) effects of a reduced-fat diet intervention in individuals with glucose intolerance. *Diabetes Care* 2001;24:619-24.

Swinburn BA, Woollard GA, Chang EC, Wilson MR. Effects of reduced-fat diets consumed ad libitum on intake of nutrients, particularly antioxidant vitamins. *J Am Diet Assoc* 1999;99:1400-5.

Tay J, Brinkworth GD, Noakes M, Keogh J, Clifton PM. Metabolic effects of weight loss on a very-low-carbohydrate diet compared with an isocaloric high-carbohydrate diet in abdominally obese subjects. *J Am Coll Cardiol* 2008;51:59-67.

ter Bogt NCW, Bemelmans WJE, Beltman FW, Broer J, Smit AJ, van der Meer K. Preventing weight gain: One-year results of a randomized lifestyle intervention. *American Journal of Preventive Medicine* 2009;37:270-277.

Terry P, Giovannucci E, Michels KB, Bergkvist L, Hansen H, Holmberg L, et al. Fruit, vegetables, dietary fiber, and risk of colorectal cancer. *J Natl Cancer Inst* 2001;93:525-33.

Tershakovec AM, Jawad AF, Stallings VA, Zemel BS, McKenzie JM, Stolley PD, et al. Growth of hypercholesterolemic children completing physician-initiated low-fat dietary intervention. *J Pediatr* 1998;133:28-34.

Thomas DE, Elliott EJ, Baur L. Low glycaemic index or low glycaemic load diets for overweight and obesity. *SOURCE Cochrane Database of Systematic Reviews (2007)* :3 Article Number: CD005105. Date of Publication: 2007 2007.

Thorpe MP, Jacobson EH, Layman DK, He X, Kris-Etherton PM, Evans EM. A diet high in protein, dairy, and calcium attenuates bone loss over twelve months of weight loss and maintenance relative to a conventional high-carbohydrate diet in adults. *J Nutr* 2008;138:1096-100.

Thorsteinsson R, Johannesson A, Jonsson H, Thorhallsson T, Sigurdsson JA. Effects of dietary intervention on serum lipids in factory workers. *Scand J Prim Health Care* 1994;12:93-9.

Togo P, Osler M, Sorensen TI, Heitmann BL. Food intake patterns and body mass index in observational studies. *Int J Obes Relat Metab Disord* 2001;25:1741-51.

Toobert DJ, Glasgow RE, Strycker LA, Barrera M, Jr., Radcliffe JL, Wander RC, et al. Biologic and quality-of-life outcomes from the Mediterranean Lifestyle Program: a randomized clinical trial.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Diabetes Care 2003;26:2288-93.

Torgerson JS, Agren L, Sjostrom L. Effects on body weight of strict or liberal adherence to an initial period of VLCD treatment. A randomised, one-year clinical trial of obese subjects. *Int J Obes Relat Metab Disord* 1999;23:190-7.

Trapp CB, Barnard ND. Usefulness of vegetarian and vegan diets for treating type 2 diabetes. *Current Diabetes Reports* 2010;10:152-158.

Trichopoulou A, Gnardellis C, Benetou V, Lagiou P, Bamia C, Trichopoulos D. Lipid, protein and carbohydrate intake in relation to body mass index. *Eur J Clin Nutr*. 2002;56:37-43.

Trichopoulou A, Gnardellis C, Lagiou A, Benetou V, Naska A, Trichopoulos D. Physical activity and energy intake selectively predict the waist-to-hip ratio in men but not in women. *Am J Clin Nutr*. 2001;74:574-8.

Trichopoulou A, Naska A, Orfanos P, Trichopoulos D. Mediterranean diet in relation to body mass index and waist-to-hip ratio: the Greek European Prospective Investigation into Cancer and Nutrition Study. *Am J Clin Nutr* 2005;82:935-40.

Trichopoulou A, Psaltopoulou T, Orfanos P, Hsieh CC, Trichopoulos D. Low-carbohydrate-high-protein diet and long-term survival in a general population cohort. *Eur J Clin Nutr* 2007;61:575-81.

Trichopoulou A, Psaltopoulou T, Orfanos P, Trichopoulos D. Diet and physical activity in relation to overall mortality amongst adult diabetics in a general population cohort. *J Intern Med* 2006;259:583-91.

Truby H, Baic S, deLooy A, Fox KR, Livingstone MB, Logan CM, et al. Randomised controlled trial of four commercial weight loss programmes in the UK: initial findings from the BBC "diet trials". *BMJ* 2006;332:1309-14.

Tsai AG, Glick HA, Shera D, Stern L, Samaha FF. Cost-effectiveness of a low-carbohydrate diet and a standard diet in severe obesity. *Obes Res* 2005;13:1834-40.

Tuomilehto H, Gylling H, Peltonen M, Martikainen T, Sahlman J, Kokkarinen J, et al. Sustained improvement in mild obstructive sleep apnea after a diet- and physical activity-based lifestyle intervention: postinterventional follow-up. *Am J Clin Nutr* 2010;92:688-96.

Turner-McGrievy GM, Barnard ND, Cohen J, Jenkins DJ, Gloede L, Green AA. Changes in nutrient intake and dietary quality among participants with type 2 diabetes following a low-fat vegan diet or a conventional diabetes diet for 22 weeks. *J Am Diet Assoc* 2008;108:1636-45.

Turner-McGrievy GM, Barnard ND, Scialli AR. A two-year randomized weight loss trial comparing a vegan diet to a more moderate low-fat diet. *Obesity (Silver Spring)* 2007;15:2276-81.

Turner-McGrievy GM, Jenkins DJ, Barnard ND, Cohen J, Gloede L, Green AA. Decreases in dietary glycemic index are related to weight loss among individuals following therapeutic diets for type 2 diabetes. *J Nutr* 2011;141:1469-74.

Tzima N, Pitsavos C, Panagiotakos DB, Skoumas J, Zampelas A, Chrysohoou C, et al. Mediterranean diet and insulin sensitivity, lipid profile and blood pressure levels, in overweight and obese people;

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

the Attica study. *Lipids Health Dis* 2007;6:22.

Wadden TA, Foster GD, Sarwer DB, Anderson DA, Gladis M, Sanderson RS, et al. Dieting and the development of eating disorders in obese women: results of a randomized controlled trial. *Am J Clin Nutr* 2004;80:560-8.

Wadden TA, Frey DL. A multicenter evaluation of a proprietary weight loss program for the treatment of marked obesity: a five-year follow-up. *Int J Eat Disord* 1997;22:203-12.

Wal JS, McBurney MI, Moellering N, Marth J, Dhurandhar NV. Moderate-carbohydrate low-fat versus low-carbohydrate high-fat meal replacements for weight loss. *Int J Food Sci Nutr* 2007;58:321-9.

Walker KZ, O'Dea K, Gomez M, Girgis S, Colagiuri R. Diet and exercise in the prevention of diabetes. *J Hum Nutr Diet*. 2010;23:344-52.

van Bakel MM, Kaaks R, Feskens EJ, Rohrmann S, Welch AA, Pala V, et al. Dietary glycaemic index and glycaemic load in the European Prospective Investigation into Cancer and Nutrition. *Eur J Clin Nutr*. 2009;63 Suppl 4:S188-205.

van Dam RM, Rimm EB, Willett WC, Stampfer MJ, Hu FB. Dietary patterns and risk for type 2 diabetes mellitus in U.S. men. *Ann Intern Med*. 2002;136:201-9.

van Dam RM, Willett WC, Rimm EB, Stampfer MJ, Hu FB. Dietary fat and meat intake in relation to risk of type 2 diabetes in men. *Diabetes Care* 2002;25:417-24.

van Gils CH, Peeters PH, Bueno-de-Mesquita HB, Boshuizen HC, Lahmann PH, Clavel-Chapelon F, et al. Consumption of vegetables and fruits and risk of breast cancer. *JAMA* 2005;293:183-93.

Van Loan M. The role of dairy foods and dietary calcium in weight management. *Journal of the American College of Nutrition* 2009;28:120S-129S.

van Strien T, van de Laar FA, van Leeuwe JF, Lucassen PL, van den Hoogen HJ, Rutten GE, et al. The dieting dilemma in patients with newly diagnosed type 2 diabetes: does dietary restraint predict weight gain 4 years after diagnosis? *Health Psychol*. 2007;26:105-12.

Van Strien T, Van de Laar FA. Intake of energy is best predicted by overeating tendency and consumption of fat is best predicted by dietary restraint: a 4-year follow-up of patients with newly diagnosed Type 2 diabetes. *Appetite*. 2008;50:544-7.

Wang L, Manson JE, Forman JP, Gaziano JM, Buring JE, Sesso HD. Dietary fatty acids and the risk of hypertension in middle-aged and older women. *Hypertension* 2010;56:598-604.

Wang P, Holst C, Astrup A, Bouwman FG, van Otterdijk S, Wodzig WK, et al. Blood profiling of proteins and steroids during weight maintenance with manipulation of dietary protein level and glycaemic index. *Br J Nutr* 2012;107:106-19.

Wang YF, Yancy WS, Jr., Yu D, Champagne C, Appel LJ, Lin PH. The relationship between dietary protein intake and blood pressure: results from the PREMIER study. *J Hum Hypertens*. 2008;22:745-54.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Warziski MT, Sereika SM, Styn MA, Music E, Burke LE. Changes in self-efficacy and dietary adherence: the impact on weight loss in the PREFER study. *J Behav Med* 2008;31:81-92.

Vasilaras TH, Astrup A, Raben A. Micronutrient intake in overweight subjects is not deficient on an ad libitum fat-reduced, high-simple carbohydrate diet. *Eur J Clin Nutr* 2004;58:326-36.

Vazquez JA, Adibi SA. Protein sparing during treatment of obesity: ketogenic versus nonketogenic very low calorie diet. *Metabolism* 1992;41:406-14.

Webb PM, Byrne C, Schnitt SJ, Connolly JL, Jacobs TW, Baer HJ, et al. A prospective study of diet and benign breast disease. *Cancer Epidemiol Biomarkers Prev.* 2004;13:1106-13.

Weijenberg MP, Mullie PF, Brants HA, Heinen MM, Goldbohm RA, van den Brandt PA. Dietary glycemic load, glycemic index and colorectal cancer risk: results from the Netherlands Cohort Study. *Int J Cancer.* 2008;122:620-9.

Weinsier RL, Johnston MH, Doleys DM, Bacon JA. Dietary management of obesity: evaluation of the time-energy displacement diet in terms of its efficacy and nutritional adequacy for long-term weight control. *Br J Nutr* 1982;47:367-79.

Weinsier RL, Seeman A, Herrera MG, Assal JP, Soeldner JS, Gleason RE. High- and low-carbohydrate diets in diabetes mellitus. Study of effects on diabetic control, insulin secretion, and blood lipids. *Ann Intern Med* 1974;80:332-41.

Velasquez MT, Bhatena SJ. Role of dietary soy protein in obesity. *International Journal of Medical Sciences* 2007;4:72-82.

Velie EM, Schairer C, Flood A, He JP, Khattree R, Schatzkin A. Empirically derived dietary patterns and risk of postmenopausal breast cancer in a large prospective cohort study. *Am J Clin Nutr.* 2005;82:1308-19.

Welty FK, Nasca MM, Lew NS, Gregoire S, Ruan Y. Effect of onsite dietitian counseling on weight loss and lipid levels in an outpatient physician office. *Am J Cardiol* 2007;100:73-5.

Westerterp-Plantenga MS, Kempen KP, Saris WH. Determinants of weight maintenance in women after diet-induced weight reduction. *Int J Obes Relat Metab Disord* 1998;22:1-6.

Westerterp-Plantenga MS, Lejeune MP, Nijs I, van Ooijen M, Kovacs EM. High protein intake sustains weight maintenance after body weight loss in humans. *Int J Obes Relat Metab Disord* 2004;28:57-64.

Westman EC, Samaha FF. Trial finds low carb diets reduce weight more than low fat diets in the short term, but have more minor adverse effects. *Evidence-based Cardiovascular Medicine* 2004;8:374-377.

Westman EC, Yancy WS, Jr., Olsen MK, Dudley T, Guyton JR. Effect of a low-carbohydrate, ketogenic diet program compared to a low-fat diet on fasting lipoprotein subclasses. *Int J Cardiol* 2006;110:212-6.

Vetter ML, Iqbal N, Dalton-Bakes C, Volger S, Wadden TA. Long-term effects of low-carbohydrate versus low-fat diets in obese persons. *Ann Intern Med* 2010;152:334-5.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Whelton PK, Appel LJ, Espeland MA, Applegate WB, Ettinger WH, Jr., Kostis JB, et al. Sodium reduction and weight loss in the treatment of hypertension in older persons: a randomized controlled trial of nonpharmacologic interventions in the elderly (TONE). TONE Collaborative Research Group. *JAMA* 1998;279:839-46.

Whelton PK, Kumanyika SK, Cook NR, Cutler JA, Borhani NO, Hennekens CH, et al. Efficacy of nonpharmacologic interventions in adults with high-normal blood pressure: results from phase 1 of the Trials of Hypertension Prevention. Trials of Hypertension Prevention Collaborative Research Group. *Am J Clin Nutr* 1997;65:652S-660S.

White C, Drummond S, De Looy A. Comparing advice to decrease both dietary fat and sucrose, or dietary fat only, on weight loss, weight maintenance and perceived quality of life. *Int J Food Sci Nutr* 2010;61:282-94.

Vignolo M, Rossi F, Bardazza G, Pistorio A, Parodi A, Spigno S, et al. Five-year follow-up of a cognitive-behavioural lifestyle multidisciplinary programme for childhood obesity outpatient treatment. *Eur J Clin Nutr* 2008;62:1047-57.

Wilfley DE, Tibbs TL, Van Buren D, Reach KP, Walker MS, Epstein LH. Lifestyle interventions in the treatment of childhood overweight: A meta-analytic review of randomized controlled trials. *Health Psychology* 2007;26:521-532.

Villareal R. A randomized, open-label, intervention study to evaluate the effect of diet composition on weight gain in obese, type 2 diabetes patients receiving intensive insulin therapy. *Journal* 2008:364.

Willett WC. Diet and breast cancer. *J Intern Med*. 2001;249:395-411.

Willett WC. Dietary fat plays a major role in obesity: No. *Obesity Reviews* 2002;3:59-68.

Williams CL, Strobino BA. Childhood diet, overweight, and CVD risk factors: the Healthy Start project. *Prev Cardiol* 2008;11:11-20.

Williamson DA, Anton SD, Han H, Champagne CM, Allen R, LeBlanc E, et al. Adherence is a multi-dimensional construct in the POUNDS LOST trial. *J Behav Med* 2010;33:35-46.

Williamson DA, Martin CK, Anton SD, York-Crowe E, Han H, Redman L, et al. Is caloric restriction associated with development of eating-disorder symptoms? Results from the CALERIE trial. *Health Psychol* 2008;27:S32-42.

Wing RR, Venditti E, Jakicic JM, Polley BA, Lang W. Lifestyle intervention in overweight individuals with a family history of diabetes. *Diabetes Care* 1998;21:350-9.

Wirfält E, Hedblad B, Gullberg B, Mattisson I, Andr n C, Rosander U, et al. Food patterns and components of the metabolic syndrome in men and women: a cross-sectional study within the Malm  Diet and Cancer cohort. *American Journal of Epidemiology* 2001;154:1150-1159.

Wolever TM, Gibbs AL, Mehling C, Chiasson JL, Connelly PW, Josse RG, et al. The Canadian Trial of Carbohydrates in Diabetes (CCD), a 1-y controlled trial of low-glycemic-index dietary carbohydrate in type 2 diabetes: no effect on glycated hemoglobin but reduction in C-reactive protein. *American*

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Journal of Clinical Nutrition 2008;87:114-125.

Wolfe AR, Arroyo C, Tedders SH, Li Y, Dai Q, Zhang J. Dietary protein and protein-rich food in relation to severely depressed mood: A 10 year follow-up of a national cohort. *Prog Neuropsychopharmacol Biol Psychiatry*. 2011;35:232-8.

von Ruesten A, Illner AK, Buijsse B, Heidemann C, Boeing H. Adherence to recommendations of the German food pyramid and risk of chronic diseases: results from the EPIC-Potsdam study. *Eur J Clin Nutr* 2010;64:1251-9.

Wood RJ, Volek JS, Liu Y, Shachter NS, Contois JH, Fernandez ML. Carbohydrate restriction alters lipoprotein metabolism by modifying VLDL, LDL, and HDL subfraction distribution and size in overweight men. *J Nutr* 2006;136:384-9.

Wu K, Hu FB, Willett WC, Giovannucci E. Dietary patterns and risk of prostate cancer in U.S. men. *Cancer Epidemiol Biomarkers Prev*. 2006;15:167-71.

Wu T, Gao X, Chen M, van Dam RM. Long-term effectiveness of diet-plus-exercise interventions vs. diet-only interventions for weight loss: a meta-analysis. *Obes Rev* 2009;10:313-23.

Wylie-Rosett J, Wassertheil-Smoller S, Blaufox MD, Davis BR, Langford HG, Oberman A, et al. Trial of antihypertensive intervention and management: greater efficacy with weight reduction than with a sodium-potassium intervention. *J Am Diet Assoc* 1993;93:408-15.

Yamagishi K, Iso H, Date C, Fukui M, Wakai K, Kikuchi S, et al. Fish, omega-3 polyunsaturated fatty acids, and mortality from cardiovascular diseases in a nationwide community-based cohort of Japanese men and women the JACC (Japan Collaborative Cohort Study for Evaluation of Cancer Risk) Study. *J Am Coll Cardiol* 2008;52:988-96.

Yamagishi K, Iso H, Yatsuya H, Tanabe N, Date C, Kikuchi S, et al. Dietary intake of saturated fatty acids and mortality from cardiovascular disease in Japanese: the Japan Collaborative Cohort Study for Evaluation of Cancer Risk (JACC) Study. *Am J Clin Nutr* 2010;92:759-65.

Yancy WS, Jr., Almirall D, Maciejewski ML, Kolotkin RL, McDuffie JR, Westman EC. Effects of two weight-loss diets on health-related quality of life. *Qual Life Res* 2009;18:281-9.

Yancy WS, Jr., Olsen MK, Dudley T, Westman EC. Acid-base analysis of individuals following two weight loss diets. *Eur J Clin Nutr* 2007;61:1416-22.

Yancy WS, Jr., Olsen MK, Guyton JR, Bakst RP, Westman EC. A low-carbohydrate, ketogenic diet versus a low-fat diet to treat obesity and hyperlipidemia: a randomized, controlled trial. *Ann Intern Med* 2004;140:769-77.

Yancy WS, Jr., Westman EC, McDuffie JR, Grambow SC, Jeffreys AS, Bolton J, et al. A randomized trial of a low-carbohydrate diet vs orlistat plus a low-fat diet for weight loss. *Arch Intern Med* 2010;170:136-45.

Yaskin J, Toner RW, Goldfarb N. Obesity management interventions: A review of the evidence. *Population Health Management* 2009;12:305-316.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Yatsuya H, Jeffery RW, Langer SL, Mitchell N, Flood AP, Welsh EM, et al. Changes in C-reactive protein during weight loss and the association with changes in anthropometric variables in men and women: LIFE Study. *Int J Obes (Lond)* 2011;35:684-91.

Yu CC, Sung RY, So RC, Lui KC, Lau W, Lam PK, et al. Effects of strength training on body composition and bone mineral content in children who are obese. *J Strength Cond Res* 2005;19:667-72.

Zambon A, Sartore G, Passera D, Francini-Pesenti F, Bassi A, Basso C, et al. Effects of hypocaloric dietary treatment enriched in oleic acid on LDL and HDL subclass distribution in mildly obese women. *J Intern Med* 1999;246:191-201.

Zemel MB, Thompson W, Zemel P, Nocton AM, Milstead A, Morris K, et al. Dietary calcium and dairy products accelerate weight and fat loss during energy restriction in obese adults. *American Journal of Clinical Nutrition* 2002;342s-343s.

Zeybek C, Aktuglu-Zeybek C, Onal H, Altay S, Erdem A, Celebi A. Right ventricular subclinical diastolic dysfunction in obese children: the effect of weight reduction with a low-carbohydrate diet. *Pediatr Cardiol* 2009;30:946-53.

Zhang S, Hunter DJ, Rosner BA, Colditz GA, Fuchs CS, Speizer FE, et al. Dietary fat and protein in relation to risk of non-Hodgkin's lymphoma among women. *J Natl Cancer Inst.* 1999;91:1751-8.

Kapitel 3.2

The National Diet-Heart Study Final Report. *Circulation* 1968;37:11-428.

Mortality from coronary heart disease is inversely related to fish consumption in The Netherlands. *Nutr Rev* 1985;43:271-3.

The Hypertension Prevention Trial: three-year effects of dietary changes on blood pressure. Hypertension Prevention Trial Research Group. *Arch Intern Med* 1990;150:153-62.

UK Prospective Diabetes Study (UKPDS). X. Urinary albumin excretion over 3 years in diet-treated type 2, (non-insulin-dependent) diabetic patients, and association with hypertension, hyperglycaemia and hypertriglyceridaemia. *Diabetologia*. 1993;36:1021-9.

Efficacy and safety of lowering dietary intake of fat and cholesterol in children with elevated low-density lipoprotein cholesterol. The Dietary Intervention Study in Children (DISC). The Writing Group for the DISC Collaborative Research Group. *JAMA*. 1995;273:1429-35.

Multicenter ophthalmic and nutritional age-related macular degeneration study--part 1: design, subjects and procedures. Age-related Macular Degeneration Study Group. *J Am Optom Assoc*. 1996;67:12-29.

Effects of weight loss and sodium reduction intervention on blood pressure and hypertension incidence in overweight people with high-normal blood pressure. The Trials of Hypertension Prevention, phase II. The Trials of Hypertension Prevention Collaborative Research Group. *Arch Intern Med* 1997;157:657-67.

All diets work equally poorly. *Journal of the National Medical Association* 2009;101:743.

Insulin-like growth factor 1 (IGF1), IGF binding protein 3 (IGFBP3), and breast cancer risk: Pooled individual data analysis of 17 prospective studies. *The Lancet Oncology* 2010;11:530-542.

Aatola H, Koivisto T, Hutri-Kahonen N, Juonala M, Mikkilä V, Lehtimäki T, et al. Lifetime fruit and vegetable consumption and arterial pulse wave velocity in adulthood: the Cardiovascular Risk in Young Finns Study. *Circulation* 2010;122:2521-8.

Abbott RD, Curb JD, Rodriguez BL, Sharp DS, Burchfiel CM, Yano K. Effect of dietary calcium and milk consumption on risk of thromboembolic stroke in older middle-aged men. The Honolulu Heart Program. *Stroke*. 1996;27:813-8.

Abramson JH, Hopp C. The control of cardiovascular risk factors in the elderly. *Prev Med* 1976;5:32-47.

Acharya S, Maskarinec G, Williams AE, Oshiro C, Hebshi S, Murphy SP. Nutritional changes among premenopausal women undertaking a soya based dietary intervention study in Hawaii. *Journal of Human Nutrition and Dietetics* 2004;17:413-419.

Adam O, Beringer C, Kless T, Lemmen C, Adam A, Wiseman M, et al. Anti-inflammatory effects of a low arachidonic acid diet and fish oil in patients with rheumatoid arthritis. *Rheumatol Int*. 2003;23:27-36.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Agudo A, Cabrera L, Amiano P, Ardanaz E, Barricarte A, Berenguer T, et al. Fruit and vegetable intakes, dietary antioxidant nutrients, and total mortality in Spanish adults: findings from the Spanish cohort of the European Prospective Investigation into Cancer and Nutrition (EPIC-Spain). *Am J Clin Nutr* 2007;85:1634-42.

Agurs-Collins T, Rosenberg L, Makambi K, Palmer JR, Adams-Campbell L. Dietary patterns and breast cancer risk in women participating in the Black Women's Health Study. *Am J Clin Nutr*. 2009;90:621-8.

Akbaraly TN, Brunner EJ, Ferrie JE, Marmot MG, Kivimaki M, Singh-Manoux A. Dietary pattern and depressive symptoms in middle age. *Br J Psychiatry* 2009;195:408-13.

Akbaraly TN, Singh-Manoux A, Tabak AG, Jokela M, Virtanen M, Ferrie JE, et al. Overall diet history and reversibility of the metabolic syndrome over 5 years: The Whitehall II prospective cohort study. *Diabetes Care* 2010;33:2339-2341.

Akesson A, Weismayer C, Newby PK, Wolk A. Combined effect of low-risk dietary and lifestyle behaviors in primary prevention of myocardial infarction in women. *Arch Intern Med* 2007;167:2122-7.

Akhter M, Inoue M, Kurahashi N, Iwasaki M, Sasazuki S, Tsugane S. Dietary soy and isoflavone intake and risk of colorectal cancer in the Japan public health center-based prospective study. *Cancer Epidemiol Biomarkers Prev* 2008;17:2128-35.

Albert CM, Gaziano JM, Willett WC, Manson JE. Nut consumption and decreased risk of sudden cardiac death in the Physicians' Health Study. *Arch Intern Med* 2002;162:1382-7.

Albert CM, Oh K, Whang W, Manson JE, Chae CU, Stampfer MJ, et al. Dietary alpha-linolenic acid intake and risk of sudden cardiac death and coronary heart disease. *Circulation* 2005;112:3232-8.

Albertson AM, Thompson D, Franko DL, Holschuh NM, Bauserman R, Barton BA. Prospective associations among cereal intake in childhood and adiposity, lipid levels, and physical activity during late adolescence. *J Am Diet Assoc* 2009;109:1775-80.

Alexy U, Libuda L, Mersmann S, Kersting M. Convenience foods in children's diet and association with dietary quality and body weight status. *Eur J Clin Nutr*. 2011;65:160-6.

Alexy U, Sichert-Hellert W, Kersting M, Schultze-Pawlitschko V. Pattern of long-term fat intake and BMI during childhood and adolescence--results of the DONALD Study. *Int J Obes Relat Metab Disord* 2004;28:1203-9.

Alonso A, Beunza JJ, Delgado-Rodriguez M, Martinez JA, Martinez-Gonzalez MA. Low-fat dairy consumption and reduced risk of hypertension: the Seguimiento Universidad de Navarra (SUN) cohort. *Am J Clin Nutr* 2005;82:972-9.

Alonso A, de la Fuente C, Martin-Arnau AM, de Irala J, Martinez JA, Martinez-Gonzalez MA. Fruit and vegetable consumption is inversely associated with blood pressure in a Mediterranean population with a high vegetable-fat intake: the Seguimiento Universidad de Navarra (SUN) Study. *Br J Nutr* 2004;92:311-9.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Altorf-van der Kuil W, Engberink MF, van Rooij FJ, Hofman A, van't Veer P, Witteman JC, et al. Dietary protein and risk of hypertension in a Dutch older population: the Rotterdam study. *J Hypertens*. 2010;28:2394-400.

Ambrosini GL, Huang RC, Mori TA, Hands BP, O'Sullivan TA, de Klerk NH, et al. Dietary patterns and markers for the metabolic syndrome in Australian adolescents. *Nutr Metab Cardiovasc Dis*. 2010;20:274-83.

Andersen P, Nilsen DW, Lyberg Beckmann S, Holme I, Hjermann I. Increased fibrinolytic potential after diet intervention in healthy coronary high-risk individuals. *Acta Med Scand* 1988;223:499-506.

Anderson R, Anderson D, Hurst C. Modeling factors that influence exercise and dietary change among midlife Australian women: results from the Healthy Aging of Women Study. *Maturitas*. 2010;67:151-8.

Appleby P, Roddam A, Allen N, Key T. Comparative fracture risk in vegetarians and nonvegetarians in EPIC-Oxford. *Eur J Clin Nutr* 2007;61:1400-6.

Appleby PN, Davey GK, Key TJ. Hypertension and blood pressure among meat eaters, fish eaters, vegetarians and vegans in EPIC-Oxford. *Public Health Nutr*. 2002;5:645-54.

Appleby PN, Key TJ, Burr ML, Thorogood M. Mortality and fresh fruit consumption. *IARC Sci Publ* 2002;156:131-3.

Appleby PN, Key TJ, Thorogood M, Burr ML, Mann J. Mortality in British vegetarians. *Public Health Nutr* 2002;5:29-36.

Appleby PN, Thorogood M, Mann JI, Key TJ. The Oxford Vegetarian Study: an overview. *Am J Clin Nutr* 1999;70:525S-531S.

Arntzenius AC, Kromhout D, Barth JD. Diet, lipoproteins, and the progression of coronary atherosclerosis. The Leiden Intervention Trial. *New England Journal of Medicine* 1985;312:805-811.

Arts IC, Jacobs DR, Jr., Gross M, Harnack LJ, Folsom AR. Dietary catechins and cancer incidence among postmenopausal women: the Iowa Women's Health Study (United States). *Cancer Causes Control* 2002;13:373-82.

Ascherio A, Rimm EB, Giovannucci EL, Colditz GA, Rosner B, Willett WC, et al. A prospective study of nutritional factors and hypertension among US men. *Circulation*. 1992;86:1475-84.

Ash S, Koutsoukos M, Capra S, Yeo S, Kerr G. Intensive diet therapy improves weight and glycaemic control in overweight men with type II diabetes. *American Journal of Clinical Nutrition* 2002;382s.

Astorg P. Dietary N-6 and N-3 polyunsaturated fatty acids and prostate cancer risk: a review of epidemiological and experimental evidence. *Cancer Causes Control* 2004;15:367-86.

Astrup A, Chaput JP, Gilbert JA, Lorenzen JK. Dairy beverages and energy balance. *Physiology and Behavior* 2010;100:67-75.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Astrup A, Skov AR, Haulrik N, Toubro S, Molgaard C. The effect of protein intake on bone mineralisation: A randomised controlled 6-months trial in overweight subjects. *American Journal of Clinical Nutrition* 2002;34:3s.

Aubertin-Leheudre M, Adlercreutz H. Relationship between animal protein intake and muscle mass index in healthy women. *British Journal of Nutrition* 2009;102:1803-1810.

Augustsson K, Michaud DS, Rimm EB, Leitzmann MF, Stampfer MJ, Willett WC, et al. A prospective study of intake of fish and marine fatty acids and prostate cancer. *Cancer Epidemiol Biomarkers Prev* 2003;12:64-7.

Aune D, Chan DS, Lau R, Vieira R, Greenwood DC, Kampman E, et al. Dietary fibre, whole grains, and risk of colorectal cancer: systematic review and dose-response meta-analysis of prospective studies. *BMJ* 2011;343:d6617.

Aune D, Lau R, Chan DS, Vieira R, Greenwood DC, Kampman E, et al. Nonlinear reduction in risk for colorectal cancer by fruit and vegetable intake based on meta-analysis of prospective studies. *Gastroenterology* 2011;141:106-18.

Aune D, Ursin G, Veierod MB. Meat consumption and the risk of type 2 diabetes: a systematic review and meta-analysis of cohort studies. *Diabetologia* 2009;52:2277-87.

Azad KA, Alam MN, Haq SA, Nahar S, Chowdhury MA, Ali SM, et al. Vegetarian diet in the treatment of fibromyalgia. *Bangladesh Med Res Counc Bull.* 2000;26:41-7.

Azadbakht L, Fard NRP, Karimi M, Baghaei MH, Surkan PJ, Rahimi M, et al. Effects of the Dietary Approaches to Stop Hypertension (DASH) eating plan on cardiovascular risks among type 2 diabetic patients: A randomized crossover clinical trial. *Diabetes Care* 2011;34:55-57.

Azadbakht L, Mirmiran P, Esmailzadeh A, Azizi F. Dairy consumption is inversely associated with the prevalence of the metabolic syndrome in Tehranian adults. *Am J Clin Nutr* 2005;82:523-30.

Azadbakht L, Mirmiran P, Esmailzadeh A, Azizi T, Azizi F. Beneficial effects of a Dietary Approaches to Stop Hypertension eating plan on features of the metabolic syndrome. *Diabetes Care* 2005;28:2823-31.

Balkau B, Vierron E, Vernay M, Born C, Arondel D, Petrella A, et al. The impact of 3-year changes in lifestyle habits on metabolic syndrome parameters: the D.E.S.I.R study. *Eur J Cardiovasc Prev Rehabil.* 2006;13:334-40.

Ballard-Barbash R, Callaway CW. Marine fish oils: role in prevention of coronary artery disease. *Mayo Clin Proc* 1987;62:113-8.

Bandera EV, Gifkins DM, Moore DF, McCullough ML, Kushi LH. Antioxidant vitamins and the risk of endometrial cancer: a dose-response meta-analysis. *Cancer Causes Control.* 2009;20:699-711.

Bandera EV, Kushi LH, Moore DF, Gifkins DM, McCullough ML. Fruits and vegetables and endometrial cancer risk: a systematic literature review and meta-analysis. *Nutr Cancer* 2007;58:6-21.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Banegas JR, Leon-Munoz LM, Guallar-Castillon P, Graciani A, Rodriguez-Artalejo F. Self-reported adherence to nonpharmacological treatment and association with mortality over 6 years: Population-based study in older persons with hypercholesterolemia. *Journal of the American Geriatrics Society* 2009;57:2287-2292.

Baran D, Sorensen A, Grimes J, Lew R, Karellas A, Johnson B, et al. Dietary modification with dairy products for preventing vertebral bone loss in premenopausal women: a three-year prospective study. *J Clin Endocrinol Metab.* 1990;70:264-70.

Barcelo F, Perona JS, Prades J, Funari SS, Gomez-Gracia E, Conde M, et al. Mediterranean-style diet effect on the structural properties of the erythrocyte cell membrane of hypertensive patients: the Prevencion con Dieta Mediterranea Study. *Hypertension* 2009;54:1143-50.

Barclay AW, Flood VM, Rochtchina E, Mitchell P, Brand-Miller JC. Glycemic index, dietary fiber, and risk of type 2 diabetes in a cohort of older Australians. *Diabetes Care* 2007;30:2811-3.

Barclay AW, Petocz P, McMillan-Price J, Flood VM, Prvan T, Mitchell P, et al. Glycemic index, glycemic load, and chronic disease risk--a meta-analysis of observational studies. *Am J Clin Nutr.* 2008;87:627-37.

Barnard ND, Cohen J, Jenkins DJA, Turner-McGrievy G, Gloede L, Jaster B, et al. A low-fat vegan diet improves glycemic control and cardiovascular risk factors in a randomized clinical trial in individuals with type 2 diabetes. *Diabetes Care* 2006;29:1777-1783.

Barr SI, Janelle KC, Prior JC. Vegetarian vs nonvegetarian diets, dietary restraint, and subclinical ovulatory disturbances: prospective 6-mo study. *Am J Clin Nutr.* 1994;60:887-94.

Barr SI, Prior JC, Janelle KC, Lentle BC. Spinal bone mineral density in premenopausal vegetarian and nonvegetarian women: cross-sectional and prospective comparisons. *J Am Diet Assoc.* 1998;98:760-5.

Barton BA, Eldridge AL, Thompson D, Affenito SG, Striegel-Moore RH, Franko DL, et al. The relationship of breakfast and cereal consumption to nutrient intake and body mass index: the National Heart, Lung, and Blood Institute Growth and Health Study. *J Am Diet Assoc* 2005;105:1383-9.

Bassett DR, Abel M, Moellering RC, Jr., Rosenblatt G, Stokes J, 3rd. Coronary heart disease in Hawaii: dietary intake, depot fat, "stress", smoking, and energy balance in Hawaiian and Japanese men. *Am J Clin Nutr* 1969;22:1483-503.

Bazzano LA, He J, Ogden LG, Loria CM, Whelton PK. Dietary fiber intake and reduced risk of coronary heart disease in US men and women: the National Health and Nutrition Examination Survey I Epidemiologic Follow-up Study. *Arch Intern Med* 2003;163:1897-904.

Bazzano LA, Song Y, Bubes V, Good CK, Manson JE, Liu S. Dietary intake of whole and refined grain breakfast cereals and weight gain in men. *Obes Res* 2005;13:1952-60.

Beasley JM, Lacroix AZ, Neuhouser ML, Huang Y, Tinker L, Woods N, et al. Protein intake and incident frailty in the women's health initiative observational study. *Journal of the American Geriatrics Society* 2010;58:1063-1071.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Beilin LJ, Burke V, Puddey IB, Mori TA, Hodgson JM. Recent developments concerning diet and hypertension. *Clinical and Experimental Pharmacology and Physiology* 2001;28:1078-1082.

Belalcazar LM, Reboussin DM, Haffner SM, Reeves RS, Schwenke DC, Hoogeveen RC, et al. Marine (omega)-3 fatty acid intake: Associations with cardiometabolic risk and response to weight loss intervention in the Look AHEAD (Action for Health in Diabetes) study. *Diabetes Care* 2010;33:197-199.

Bellisle F, Altenburg de Assis MA, Fieux B, Preziosi P, Galan P, Guy-Grand B, et al. Use of 'light' foods and drinks in French adults: biological, anthropometric and nutritional correlates. *J Hum Nutr Diet.* 2001;14:191-206.

Bemelmans WJ, Broer J, Hulshof KF, Siero FW, May JF, Meyboom-de Jong B. Long-term effects of nutritional group education for persons at high cardiovascular risk. *Eur J Public Health* 2004;14:240-5.

Benassi-Evans B, Clifton PM, Noakes M, Keogh JB, Fenech M. High protein-high red meat versus high carbohydrate weight loss diets do not differ in effect on genome stability and cell death in lymphocytes of overweight men. *Mutagenesis* 2009;24:271-277.

Bendinelli B, Masala G, Saieva C, Salvini S, Calonico C, Sacerdote C, et al. Fruit, vegetables, and olive oil and risk of coronary heart disease in Italian women: the EPICOR Study. *Am J Clin Nutr* 2011;93:275-83.

Benetou V, Orfanos P, Zylis D, Sieri S, Contiero P, Tumino R, et al. Diet and hip fractures among elderly Europeans in the EPIC cohort. *Eur J Clin Nutr.* 2011;65:132-9.

Bennett W. Dietary treatments of obesity. *Ann N Y Acad Sci* 1987;499:250-63.

Beresford SA, Johnson KC, Ritenbaugh C, Lasser NL, Snetselaar LG, Black HR, et al. Low-fat dietary pattern and risk of colorectal cancer: the Women's Health Initiative Randomized Controlled Dietary Modification Trial. *JAMA* 2006;295:643-54.

Bergqvist AG, Schall JI, Stallings VA, Zemel BS. Progressive bone mineral content loss in children with intractable epilepsy treated with the ketogenic diet. *Am J Clin Nutr.* 2008;88:1678-84.

Berkey CS, Colditz GA, Rockett HR, Frazier AL, Willett WC. Dairy consumption and female height growth: prospective cohort study. *Cancer Epidemiol Biomarkers Prev.* 2009;18:1881-7.

Berkey CS, Rockett HR, Willett WC, Colditz GA. Milk, dairy fat, dietary calcium, and weight gain: a longitudinal study of adolescents. *Arch Pediatr Adolesc Med* 2005;159:543-50.

Bernstein AM, Rosner BA, Willett WC. Cereal fiber and coronary heart disease: a comparison of modeling approaches for repeated dietary measurements, intermediate outcomes, and long follow-up. *Eur J Epidemiol* 2011;26:877-86.

Bertoia M, Albanes D, Mayne ST, Mannisto S, Virtamo J, Wright ME. No association between fruit, vegetables, antioxidant nutrients and risk of renal cell carcinoma. *Int J Cancer* 2010;126:1504-12.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Bertrais S, Galan P, Renault N, Zarebska M, Preziosi P, Hercberg S. Consumption of soup and nutritional intake in French adults: consequences for nutritional status. *J Hum Nutr Diet*. 2001;14:121-8.

Bes-Rastrollo M, Martinez-Gonzalez MA, Sanchez-Villegas A, de la Fuente Arrillaga C, Martinez JA. Association of fiber intake and fruit/vegetable consumption with weight gain in a Mediterranean population. *Nutrition* 2006;22:504-11.

Bes-Rastrollo M, Sabate J, Gomez-Gracia E, Alonso A, Martinez JA, Martinez-Gonzalez MA. Nut consumption and weight gain in a Mediterranean cohort: The SUN study. *Obesity (Silver Spring)* 2007;15:107-16.

Bes-Rastrollo M, Sanchez-Villegas A, de la Fuente C, de Irala J, Martinez JA, Martinez-Gonzalez MA. Olive oil consumption and weight change: the SUN prospective cohort study. *Lipids* 2006;41:249-56.

Bes-Rastrollo M, van Dam RM, Martinez-Gonzalez MA, Li TY, Sampson LL, Hu FB. Prospective study of dietary energy density and weight gain in women. *Am J Clin Nutr* 2008;88:769-77.

Bes-Rastrollo M, Wedick NM, Martinez-Gonzalez MA, Li TY, Sampson L, Hu FB. Prospective study of nut consumption, long-term weight change, and obesity risk in women. *Am J Clin Nutr* 2009;89:1913-9.

Beulens JW, de Bruijne LM, Stolk RP, Peeters PH, Bots ML, Grobbee DE, et al. High dietary glycemic load and glycemic index increase risk of cardiovascular disease among middle-aged women: a population-based follow-up study. *J Am Coll Cardiol*. 2007;50:14-21.

Beunza JJ, Toledo E, Hu FB, Bes-Rastrollo M, Serrano-Martinez M, Sanchez-Villegas A, et al. Adherence to the Mediterranean diet, long-term weight change, and incident overweight or obesity: the Seguimiento Universidad de Navarra (SUN) cohort. *Am J Clin Nutr* 2010;92:1484-93.

Bhoopat L, Rojnuckarin P, Hiransuthikul N, Intragumtornchai T. Low vegetable intake is strongly associated with venous thromboembolism in Thai population. *Blood Coagul Fibrinolysis* 2010;21:758-63.

Biel RK, Friedenreich CM, Csizmadi I, Robson PJ, McLaren L, Faris P, et al. Case-control study of dietary patterns and endometrial cancer risk. *Nutr Cancer* 2011;63:673-86.

Bierenbaum ML, Fleischman AI, Green DP, Raichelson RI, Hayton T, Watson PB, et al. The 5-year experience of modified fat diets on younger men with coronary heart disease. *Circulation* 1970;42:943-52.

Bierenbaum ML, Fleischman AI, Raichelson RI, Hayton T, Watson PB. Ten-year experience of modified-fat diets on younger men with coronary heart-disease. *Lancet*. 1973;1:1404-7.

Bingham S, Luben R, Welch A, Tasevska N, Wareham N, Khaw KT. Epidemiologic assessment of sugars consumption using biomarkers: comparisons of obese and nonobese individuals in the European prospective investigation of cancer Norfolk. *Cancer Epidemiol Biomarkers Prev* 2007;16:1651-4.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Bingham SA, Day NE, Luben R, Ferrari P, Slimani N, Norat T, et al. Dietary fibre in food and protection against colorectal cancer in the European Prospective Investigation into Cancer and Nutrition (EPIC): an observational study. *Lancet* 2003;361:1496-501.

Bingham SA, Norat T, Moskal A, Ferrari P, Slimani N, Clavel-Chapelon F, et al. Is the association with fiber from foods in colorectal cancer confounded by folate intake? *Cancer Epidemiol Biomarkers Prev* 2005;14:1552-6.

Bjerregaard LJ, Joensen AM, Dethlefsen C, Jensen MK, Johnsen SP, Tjønneland A, et al. Fish intake and acute coronary syndrome. *Eur Heart J* 2010;31:29-34.

Black HS, Herd JA, Goldberg LH, Wolf Jr JE, Thornby JI, Rosen T, et al. Effect of a low-fat diet on the incidence of actinic keratosis. *New England Journal of Medicine* 1994;330:1272-1275.

Blatt AD, Roe LS, Rolls BJ. Hidden vegetables: an effective strategy to reduce energy intake and increase vegetable intake in adults. *Am J Clin Nutr* 2011;93:756-63.

Blomhoff R, Carlsen MH, Andersen LF, Jacobs DR, Jr. Health benefits of nuts: potential role of antioxidants. *Br J Nutr* 2006;96 Suppl 2:S52-60.

Blotman F, Maheu E, Wulwik A, Caspard H, Lopez A. Efficacy and safety of avocado/soybean unsaponifiables in the treatment of symptomatic osteoarthritis of the knee and hip. A prospective, multicenter, three-month, randomized, double-blind, placebo-controlled trial. *Rev Rhum Engl Ed.* 1997;64:825-34.

Boden-Albala B, Elkind MS, White H, Szumski A, Paik MC, Sacco RL. Dietary total fat intake and ischemic stroke risk: the Northern Manhattan Study. *Neuroepidemiology.* 2009;32:296-301.

Boeing H, Dietrich T, Hoffmann K, Pischon T, Ferrari P, Lahmann PH, et al. Intake of fruits and vegetables and risk of cancer of the upper aero-digestive tract: the prospective EPIC-study. *Cancer Causes Control* 2006;17:957-69.

Boffetta P, Couto E, Wichmann J, Ferrari P, Trichopoulos D, Bueno-de-Mesquita HB, et al. Fruit and vegetable intake and overall cancer risk in the European Prospective Investigation into Cancer and Nutrition (EPIC). *J Natl Cancer Inst* 2010;102:529-37.

Boggs DA, Palmer JR, Spiegelman D, Stampfer MJ, Adams-Campbell LL, Rosenberg L. Dietary patterns and 14-y weight gain in African American women. *Am J Clin Nutr* 2011;94:86-94.

Boniface DR, Tefft ME. Dietary fats and 16-year coronary heart disease mortality in a cohort of men and women in Great Britain. *Eur J Clin Nutr* 2002;56:786-92.

Borgquist S, Wirfalt E, Jirstrom K, Anagnostaki L, Gullberg B, Berglund G, et al. Diet and body constitution in relation to subgroups of breast cancer defined by tumour grade, proliferation and key cell cycle regulators. *Breast Cancer Res.* 2007;9:R11.

Bosetti C, Pelucchi C, La Vecchia C. Diet and cancer in Mediterranean countries: carbohydrates and fats. *Public Health Nutr* 2009;12:1595-600.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Bostick RM, Kushi LH, Wu Y, Meyer KA, Sellers TA, Folsom AR. Relation of calcium, vitamin D, and dairy food intake to ischemic heart disease mortality among postmenopausal women. *Am J Epidemiol* 1999;149:151-61.

Bostick RM, Potter JD, Kushi LH, Sellers TA, Steinmetz KA, McKenzie DR, et al. Sugar, meat, and fat intake, and non-dietary risk factors for colon cancer incidence in Iowa women (United States). *Cancer Causes Control*. 1994;5:38-52.

Bostick RM, Potter JD, Sellers TA, McKenzie DR, Kushi LH, Folsom AR. Relation of calcium, vitamin D, and dairy food intake to incidence of colon cancer among older women. The Iowa Women's Health Study. *Am J Epidemiol* 1993;137:1302-17.

Botterweck AA, van den Brandt PA, Goldbohm RA. Vitamins, carotenoids, dietary fiber, and the risk of gastric carcinoma: results from a prospective study after 6.3 years of follow-up. *Cancer* 2000;88:737-48.

Bourn DM, Mann JI, McSkimming BJ, Waldron MA, Wishart JD. Impaired glucose tolerance and NIDDM: Does a lifestyle intervention program have an effect? *Diabetes Care* 1994;17:1311-1319.

Bovbjerg VE, McCann BS, Retzlaff BM, Walden CE, Knopp RH. Effect of cholesterol-lowering diets on indices of depression and hostility. *Ann Behav Med*. 1999;21:98-101.

Boyapati SM, Shu XO, Ruan ZX, Dai Q, Cai Q, Gao YT, et al. Soyfood intake and breast cancer survival: a followup of the Shanghai Breast Cancer Study. *Breast Cancer Res Treat* 2005;92:11-7.

Boyd NF, Martin LJ, Beaton M, Cousins M, Kriukov V. Long-term effects of participation in a randomized trial of a low-fat, high-carbohydrate diet. *Cancer Epidemiol Biomarkers Prev* 1996;5:217-22.

Breslow RA, Graubard BI, Sinha R, Subar AF. Diet and lung cancer mortality: a 1987 National Health Interview Survey cohort study. *Cancer Causes Control* 2000;11:419-31.

Brouwer IA, Katan MB, Zock PL. Dietary alpha-linolenic acid is associated with reduced risk of fatal coronary heart disease, but increased prostate cancer risk: a meta-analysis. *J Nutr* 2004;134:919-22.

Brunner EJ, Mosdol A, Witte DR, Martikainen P, Stafford M, Shipley MJ, et al. Dietary patterns and 15-y risks of major coronary events, diabetes, and mortality. *Am J Clin Nutr* 2008;87:1414-21.

Buchner FL, Bueno-de-Mesquita HB, Linseisen J, Boshuizen HC, Kiemeneij LA, Ros MM, et al. Fruits and vegetables consumption and the risk of histological subtypes of lung cancer in the European Prospective Investigation into Cancer and Nutrition (EPIC). *Cancer Causes Control* 2010;21:357-71.

Buckland G, Bach A, Serra-Majem L. Obesity and the Mediterranean diet: a systematic review of observational and intervention studies. *Obes Rev* 2008;9:582-93.

Buckland G, Gonzalez CA, Agudo A, Vilardell M, Berenguer A, Amiano P, et al. Adherence to the Mediterranean diet and risk of coronary heart disease in the Spanish EPIC Cohort Study. *Am J Epidemiol*. 2009;170:1518-29.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Buijsse B, Feskens EJ, Kok FJ, Kromhout D. Cocoa intake, blood pressure, and cardiovascular mortality: the Zutphen Elderly Study. *Arch Intern Med* 2006;166:411-7.

Buitrago-Lopez A, Sanderson J, Johnson L, Warnakula S, Wood A, Di Angelantonio E, et al. Chocolate consumption and cardiometabolic disorders: systematic review and meta-analysis. *BMJ* 2011;343:d4488.

Burger KNJ, Beulens JWJ, van der Schouw YT, Sluijs I, Spijkerman AMW, Sluik D, et al. Dietary fiber, carbohydrate quality and quantity, and mortality risk of individuals with diabetes mellitus. *SOURCE PLoS ONE* (2012) 7:8 Article Number: e43127. Date of Publication: 23 Aug 2012 2012.

Burke LE, Choo J, Music E, Warziski M, Styn MA, Kim Y, et al. PREFER study: a randomized clinical trial testing treatment preference and two dietary options in behavioral weight management--rationale, design and baseline characteristics. *Contemp Clin Trials* 2006;27:34-48.

Burke LE, Hudson AG, Warziski MT, Styn MA, Music E, Elci OU, et al. Effects of a vegetarian diet and treatment preference on biochemical and dietary variables in overweight and obese adults: a randomized clinical trial. *Am J Clin Nutr* 2007;86:588-96.

Burke LE, Warziski M, Styn MA, Music E, Hudson AG, Sereika SM. A randomized clinical trial of a standard versus vegetarian diet for weight loss: the impact of treatment preference. *Int J Obes (Lond)* 2008;32:166-76.

Burke V, Lee AH, Hunter E, Spargo R, Smith R, Beilin LJ, et al. Alcohol intake and incidence of coronary disease in Australian aborigines. *Alcohol Alcohol* 2007;42:119-24.

Burr ML. Fish and ischaemic heart disease. *World Rev Nutr Diet* 1993;72:49-60.

Burr ML, Butland BK. Heart disease in British vegetarians. *Am J Clin Nutr* 1988;48:830-2.

Burr ML, Fehily AM, Gilbert JF, Rogers S, Holliday RM, Sweetnam PM, et al. Effects of changes in fat, fish, and fibre intakes on death and myocardial reinfarction: diet and reinfarction trial (DART). *Lancet* 1989;2:757-61.

Burr ML, Sweetnam PM. Vegetarianism, dietary fiber, and mortality. *Am J Clin Nutr* 1982;36:873-7.

Butler LM, Wu AH, Wang R, Koh WP, Yuan JM, Yu MC. A vegetable-fruit-soy dietary pattern protects against breast cancer among postmenopausal Singapore Chinese women. *Am J Clin Nutr* 2010;91:1013-9.

Butler TL, Fraser GE, Beeson WL, Knutsen SF, Herring RP, Chan J, et al. Cohort profile: The Adventist Health Study-2 (AHS-2). *Int J Epidemiol* 2008;37:260-5.

Buyken AE, Toeller M, Heitkamp G, Vitelli F, Stehle P, Scherbaum WA, et al. Relation of fibre intake to HbA1c and the prevalence of severe ketoacidosis and severe hypoglycaemia. *EURODIAB IDDM Complications Study Group. Diabetologia* 1998;41:882-90.

Cade JE, Burley VJ, Greenwood DC. The UK Women's Cohort Study: comparison of vegetarians, fish-eaters and meat-eaters. *Public Health Nutr.* 2004;7:871-8.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Cade JE, Burley VJ, Greenwood DC. Dietary fibre and risk of breast cancer in the UK Women's Cohort Study. *Int J Epidemiol* 2007;36:431-8.

Caggiula AW, Christakis G, Farrand M, Hulley SB, Johnson R, Lasser NL, et al. The multiple risk intervention trial (MRFIT). IV. Intervention on blood lipids. *Prev Med* 1981;10:443-75.

Cai H, Shu XO, Gao YT, Li H, Yang G, Zheng W. A prospective study of dietary patterns and mortality in Chinese women. *Epidemiology* 2007;18:393-401.

Campbell MK, Demark-Wahnefried W, Symons M, Kalsbeek WD, Dodds J, Cowan A, et al. Fruit and vegetable consumption and prevention of cancer: the Black Churches United for Better Health project. *Am J Public Health* 1999;89:1390-6.

Canfi A, Gepner Y, Schwarzfuchs D, Golan R, Shahar DR, Fraser D, et al. Effect of changes in the intake of weight of specific food groups on successful body weight loss during a multi-dietary strategy intervention trial. *J Am Coll Nutr* 2011;30:491-501.

Carayol M, Grosclaude P, Delpierre C. Prospective studies of dietary alpha-linolenic acid intake and prostate cancer risk: a meta-analysis. *Cancer Causes Control* 2010;21:347-55.

Carruth BR, Skinner JD. The role of dietary calcium and other nutrients in moderating body fat in preschool children. *Int J Obes Relat Metab Disord* 2001;25:559-66.

Carter P, Gray LJ, Troughton J, Khunti K, Davies MJ. Fruit and vegetable intake and incidence of type 2 diabetes mellitus: systematic review and meta-analysis. *BMJ* 2010;341:c4229.

Carty CL, Kooperberg C, Neuhauser ML, Tinker L, Howard B, Wactawski-Wende J, et al. Low-fat dietary pattern and change in body-composition traits in the Women's Health Initiative Dietary Modification Trial. *American Journal of Clinical Nutrition* 2011;93:516-524.

Cassidy A, O'Reilly EJ, Kay C, Sampson L, Franz M, Forman JP, et al. Habitual intake of flavonoid subclasses and incident hypertension in adults. *Am J Clin Nutr* 2011;93:338-47.

Caygill CP, Charlett A, Hill MJ. Fat, fish, fish oil and cancer. *Br J Cancer* 1996;74:159-64.

Cerhan JR, Saag KG, Merlino LA, Mikuls TR, Criswell LA. Antioxidant micronutrients and risk of rheumatoid arthritis in a cohort of older women. *Am J Epidemiol.* 2003;157:345-54.

Challier B, Perarnau JM, Viel JF. Garlic, onion and cereal fibre as protective factors for breast cancer: a French case-control study. *Eur J Epidemiol* 1998;14:737-47.

Champagne CM, Broyles ST, Moran LD, Cash KC, Levy EJ, Lin PH, et al. Dietary intakes associated with successful weight loss and maintenance during the Weight Loss Maintenance trial. *J Am Diet Assoc* 2011;111:1826-35.

Chan HLY, Wong VWS. Can dietetic intervention for obesity ever succeed in real life? *Journal of Gastroenterology and Hepatology* 2007;22:459-460.

Chan R, Woo J, Lau W, Leung J, Xu L, Zhao X, et al. Effects of lifestyle and diet on bone health in young adult Chinese women living in Hong Kong and Beijing. *Food Nutr Bull.* 2009;30:370-8.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Chan R, Woo J, Leung J. Effects of food groups and dietary nutrients on bone loss in elderly Chinese population. *J Nutr Health Aging*. 2011;15:287-94.

Chang SC, Ziegler RG, Dunn B, Stolzenberg-Solomon R, Lacey JV, Jr., Huang WY, et al. Association of energy intake and energy balance with postmenopausal breast cancer in the prostate, lung, colorectal, and ovarian cancer screening trial. *Cancer Epidemiol Biomarkers Prev*. 2006;15:334-41.

Chang-Claude J, Frentzel-Beyme R. Dietary and lifestyle determinants of mortality among German vegetarians. *Int J Epidemiol* 1993;22:228-36.

Chang-Claude J, Frentzel-Beyme R, Eilber U. Mortality pattern of German vegetarians after 11 years of follow-up. *Epidemiology* 1992;3:395-401.

Chang-Claude J, Hermann S, Eilber U, Steindorf K. Lifestyle determinants and mortality in German vegetarians and health-conscious persons: results of a 21-year follow-up. *Cancer Epidemiol Biomarkers Prev* 2005;14:963-8.

Chao A, Thun MJ, Connell CJ, McCullough ML, Jacobs EJ, Flanders WD, et al. Meat consumption and risk of colorectal cancer. *JAMA*. 2005;293:172-82.

Chavarro JE, Stampfer MJ, Hall MN, Sesso HD, Ma J. A 22-y prospective study of fish intake in relation to prostate cancer incidence and mortality. *Am J Clin Nutr* 2008;88:1297-303.

Chen G, Heilbrun LK, Venkatramanamoorthy R, Maranci V, Redd JN, Klurfeld DM, et al. Effects of low-fat and/or high-fruit-and-vegetable diets on plasma levels of 8-isoprostane-F2alpha in the Nutrition and Breast Health study. *Nutr Cancer* 2004;50:155-60.

Chen J, Gu D, Huang J, Rao DC, Jaquish CE, Hixson JE, et al. Metabolic syndrome and salt sensitivity of blood pressure in non-diabetic people in China: a dietary intervention study. *The Lancet* 2009;373:829-835.

Cheng G, Karaolis-Danckert N, Libuda L, Bolzenius K, Remer T, Buyken AE. Relation of dietary glycemic index, glycemic load, and fiber and whole-grain intakes during puberty to the concurrent development of percent body fat and body mass index. *Am J Epidemiol*. 2009;169:667-77.

Chiuve SE, Korngold EC, Januzzi JL, Jr., Gantzer ML, Albert CM. Plasma and dietary magnesium and risk of sudden cardiac death in women. *Am J Clin Nutr* 2011;93:253-60.

Cho E, Hung S, Willett WC, Spiegelman D, Rimm EB, Seddon JM, et al. Prospective study of dietary fat and the risk of age-related macular degeneration. *American Journal of Clinical Nutrition* 2001;73:209-218.

Cho E, Seddon JM, Rosner B, Willett WC, Hankinson SE. Prospective study of intake of fruits, vegetables, vitamins, and carotenoids and risk of age-related maculopathy. *Arch Ophthalmol*. 2004;122:883-92.

Cho E, Smith-Warner SA, Spiegelman D, Beeson WL, van den Brandt PA, Colditz GA, et al. Dairy foods, calcium, and colorectal cancer: a pooled analysis of 10 cohort studies. *J Natl Cancer Inst* 2004;96:1015-22.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Choi HK. Dietary risk factors for rheumatic diseases. *Curr Opin Rheumatol*. 2005;17:141-6.

Choi HK, Atkinson K, Karlson EW, Willett W, Curhan G. Purine-rich foods, dairy and protein intake, and the risk of gout in men. *N Engl J Med*. 2004;350:1093-103.

Choi HK, Gao X, Curhan G. Vitamin C intake and the risk of gout in men: a prospective study. *Arch Intern Med*. 2009;169:502-7.

Choquette S, Riesco E, Cormier E, Dion T, Aubertin-Leheudre M, Dionne JJ. Effects of soya isoflavones and exercise on body composition and clinical risk factors of cardiovascular diseases in overweight postmenopausal women: a 6-month double-blind controlled trial. *Br J Nutr* 2011;105:1199-209.

Chow CK. Meat consumption and prospective weight change. *American Journal of Clinical Nutrition* 2011;93:663-664.

Chow WH, Schuman LM, McLaughlin JK, Bjelke E, Gridley G, Wacholder S, et al. A cohort study of tobacco use, diet, occupation, and lung cancer mortality. *Cancer Causes Control* 1992;3:247-54.

Chrysohoou C, Panagiotakos DB, Aggelopoulos P, Kastorini CM, Kehagia I, Pitsavos C, et al. The Mediterranean diet contributes to the preservation of left ventricular systolic function and to the long-term favorable prognosis of patients who have had an acute coronary event. *Am J Clin Nutr* 2010;92:47-54.

Chrysohoou C, Panagiotakos DB, Pitsavos C, Skoumas J, Krinos X, Chloptsios Y, et al. Long-term fish consumption is associated with protection against arrhythmia in healthy persons in a Mediterranean region--the ATTICA study. *Am J Clin Nutr* 2007;85:1385-91.

Clifford CK, Butrum RR, Greenwald P, Yates JW. Clinical trials of low fat diets and breast cancer prevention. *Prog Clin Biol Res* 1986;222:93-115.

Colangelo LA, He K, Whooley MA, Daviglius ML, Liu K. Higher dietary intake of long-chain omega-3 polyunsaturated fatty acids is inversely associated with depressive symptoms in women. *Nutrition* 2009;25:1011-9.

Colditz GA, Branch LG, Lipnick RJ, Willett WC, Rosner B, Posner BM, et al. Increased green and yellow vegetable intake and lowered cancer deaths in an elderly population. *Am J Clin Nutr* 1985;41:32-6.

Colditz GA, Manson JE, Stampfer MJ, Rosner B, Willett WC, Speizer FE. Diet and risk of clinical diabetes in women. *Am J Clin Nutr* 1992;55:1018-23.

Colditz GA, Willett WC, Stampfer MJ, London SJ, Segal MR, Speizer FE. Patterns of weight change and their relation to diet in a cohort of healthy women. *Am J Clin Nutr* 1990;51:1100-5.

Colin Ramirez E, Castillo Martinez L, Orea Tejada A, Rebollar Gonzalez V, Narvaez David R, Asensio Lafuente E. Effects of a nutritional intervention on body composition, clinical status, and quality of life in patients with heart failure. *Nutrition* 2004;20:890-5.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Collins CE, Warren JM, Neve M, McCoy P, Stokes B. Systematic review of interventions in the management of overweight and obese children which include a dietary component. *International Journal of Evidence-Based Healthcare* 2007;5:2-53.

Cook NR, Kumanyika SK, Cutler JA, Whelton PK. Dose-response of sodium excretion and blood pressure change among overweight, nonhypertensive adults in a 3-year dietary intervention study. *J Hum Hypertens* 2005;19:47-54.

Cooper AJ, Sharp SJ, Lentjes MA, Luben RN, Khaw KT, Wareham NJ, et al. A prospective study of the association between quantity and variety of fruit and vegetable intake and incident type 2 diabetes. *Diabetes Care* 2012;35:1293-300.

Coppell KJ, Kataoka M, Williams SM, Chisholm AW, Vorgers SM, Mann JI. Nutritional intervention in patients with type 2 diabetes who are hyperglycaemic despite optimised drug treatment - Lifestyle over and above drugs in diabetes (LOADD) study: Randomised controlled trial. *BMJ* 2010;341:237.

Corbett Jr EC. Altering the glycaemic index or carbohydrate intake did not affect glycaemic control in type 2 diabetes: Commentary. *Evidence-Based Medicine* 2008;13:107.

Corbett Jr EC. 6-Month treatment with a low-glycaemic diet was better than a high-fibre diet for glycaemic control in type 2 diabetes. *Evidence-Based Medicine* 2009;14:72.

Corle DK, Sharbaugh C, Mateski DJ, Coyne T, Paskett ED, Cahill J, et al. Self-rated quality of life measures: effect of change to a low-fat, high-fiber, fruit and vegetable enriched diet. *Ann Behav Med* 2001;23:198-207.

Couch SC, Schwarzman F, Carroll J, Koenigsberger D, Nordli DR, Deckelbaum RJ, et al. Growth and nutritional outcomes of children treated with the ketogenic diet. *J Am Diet Assoc.* 1999;99:1573-5.

Cowin IS, Emmett PM. Associations between dietary intakes and blood cholesterol concentrations at 31 months. *Eur J Clin Nutr* 2001;55:39-49.

Cox RH, Parker GG, Watson AC, Robinson SH, Simonson CJ, Elledge JC, et al. Dietary cancer risk of low-income women and change with intervention. *Journal of the American Dietetic Association* 1995;95:1031-1034.

Crawford P, Paden SL. What is the dietary treatment for low HDL cholesterol? *Journal of Family Practice* 2006;55:1076-1078.

Crawley HF, While D. The diet and body weight of British teenage smokers at 16-17 years. *Eur J Clin Nutr.* 1995;49:904-14.

Crichton GE, Murphy KJ, Bryan J. Dairy intake and cognitive health in middle-aged South Australians. *Asia Pac J Clin Nutr* 2010;19:161-71.

Crowe FL, Appleby PN, Allen NE, Key TJ. Diet and risk of diverticular disease in Oxford cohort of European Prospective Investigation into Cancer and Nutrition (EPIC): prospective study of British vegetarians and non-vegetarians. *BMJ* 2011;343:d4131.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Crowe FL, Key TJ, Appleby PN, Travis RC, Overvad K, Jakobsen MU, et al. Dietary fat intake and risk of prostate cancer in the European Prospective Investigation into Cancer and Nutrition. *American Journal of Clinical Nutrition* 2008;87:1405-1413.

Crowe FL, Roddam AW, Key TJ, Appleby PN, Overvad K, Jakobsen MU, et al. Fruit and vegetable intake and mortality from ischaemic heart disease: results from the European Prospective Investigation into Cancer and Nutrition (EPIC)-Heart study. *Eur Heart J* 2011;32:1235-43.

Cui R, Iso H, Date C, Kikuchi S, Tamakoshi A. Dietary folate and vitamin b6 and B12 intake in relation to mortality from cardiovascular diseases: Japan collaborative cohort study. *Stroke* 2010;41:1285-9.

Cutler GJ, Flood A, Hannan PJ, Slavin JL, Neumark-Sztainer D. Association between major patterns of dietary intake and weight status in adolescents. *Br J Nutr* 2012;108:349-56.

Cutler JA, Stamler J. Chapter 1. Introduction and summary of the dietary and nutritional methods and findings in the Multiple Risk Factor Intervention Trial. *American Journal of Clinical Nutrition* 1997;65:184S-190S.

Dale KS, McAuley KA, Taylor RW, Williams SM, Farmer VL, Hansen P, et al. Determining optimal approaches for weight maintenance: a randomized controlled trial. *CMAJ* 2009;180:E39-46.

Daniel CR, Cross AJ, Graubard BI, Hollenbeck AR, Park Y, Sinha R. Prospective investigation of poultry and fish intake in relation to cancer risk. *Cancer Prev Res (Phila)* 2011;4:1903-11.

Dargent-Molina P, Sabia S, Touvier M, Kesse E, Breart G, Clavel-Chapelon F, et al. Proteins, dietary acid load, and calcium and risk of postmenopausal fractures in the E3N French women prospective study. *J Bone Miner Res* 2008;23:1915-22.

Das SK, Saltzman E, Gilhooly CH, DeLany JP, Golden JK, Pittas AG, et al. Low or moderate dietary energy restriction for long-term weight loss: what works best? *Obesity (Silver Spring)* 2009;17:2019-24.

Dauchet L, Amouyel P, Dallongeville J. Fruit and vegetable consumption and risk of stroke: a meta-analysis of cohort studies. *Neurology* 2005;65:1193-7.

Daviglus ML, Stamler J, Orenca AJ, Dyer AR, Liu K, Greenland P, et al. Fish consumption and the 30-year risk of fatal myocardial infarction. *N Engl J Med* 1997;336:1046-53.

Davis JN, Alexander KE, Ventura EE, Toledo-Corral CM, Goran MI. Inverse relation between dietary fiber intake and visceral adiposity in overweight Latino youth. *Am J Clin Nutr* 2009;90:1160-6.

Day RS, Fulton JE, Dai S, Mihalopoulos NL, Barradas DT. Nutrient intake, physical activity, and CVD risk factors in children: Project HeartBeat! *Am J Prev Med*. 2009;37:S25-33.

de Castro MB, Kac G, de Leon AP, Sichieri R. High-protein diet promotes a moderate postpartum weight loss in a prospective cohort of Brazilian women. *Nutrition*. 2009;25:1120-8.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

de Goede J, Geleijnse JM, Boer JM, Kromhout D, Verschuren WM. Marine (n-3) fatty acids, fish consumption, and the 10-year risk of fatal and nonfatal coronary heart disease in a large population of Dutch adults with low fish intake. *J Nutr* 2010;140:1023-8.

de Lorgeril M, Renaud S, Mamelle N, Salen P, Martin JL, Monjaud I, et al. Mediterranean alpha-linolenic acid-rich diet in secondary prevention of coronary heart disease. *Lancet* 1994;343:1454-9.

de Lorgeril M, Salen P. Mediterranean type of diet for the prevention of coronary heart disease. A global perspective from the seven countries study to the most recent dietary trials. *Int J Vitam Nutr Res* 2001;71:166-72.

de Lorgeril M, Salen P, Martin JL, Monjaud I, Boucher P, Mamelle N. Mediterranean dietary pattern in a randomized trial: prolonged survival and possible reduced cancer rate. *Arch Intern Med* 1998;158:1181-7.

de Lorgeril M, Salen P, Martin JL, Monjaud I, Delaye J, Mamelle N. Mediterranean diet, traditional risk factors, and the rate of cardiovascular complications after myocardial infarction: final report of the Lyon Diet Heart Study. *Circulation* 1999;99:779-85.

de Munter JS, Hu FB, Spiegelman D, Franz M, van Dam RM. Whole grain, bran, and germ intake and risk of type 2 diabetes: a prospective cohort study and systematic review. *PLoS Med*. 2007;4:e261.

Deheeger M, Rolland-Cachera MF, Fontvieille AM. Physical activity and body composition in 10 year old French children: linkages with nutritional intake? *Int J Obes Relat Metab Disord*. 1997;21:372-9.

deKoning L, Anand SS. Adherence to a Mediterranean diet and survival in a Greek population. Trichopoulos A, Costacou T, Bamia C, Trichopoulos D. *N Engl J Med* 2003; 348: 2599-608. *Vasc Med* 2004;9:145-6.

Del Rio D, Agnoli C, Pellegrini N, Krogh V, Brighenti F, Mazzeo T, et al. Total antioxidant capacity of the diet is associated with lower risk of ischemic stroke in a large Italian cohort. *J Nutr* 2011;141:118-23.

D'Elia L, Barba G, Cappuccio FP, Strazzullo P. Potassium intake, stroke, and cardiovascular disease: A meta-analysis of prospective studies. *Journal of the American College of Cardiology* 2011;57:1210-1219.

Dijkstra SC, Brouwer IA, van Rooij FJ, Hofman A, Witteman JC, Geleijnse JM. Intake of very long chain n-3 fatty acids from fish and the incidence of heart failure: the Rotterdam Study. *Eur J Heart Fail* 2009;11:922-8.

Djousse L, Gaziano JM. Egg consumption in relation to cardiovascular disease and mortality: the Physicians' Health Study. *Am J Clin Nutr* 2008;87:964-9.

Djousse L, Gaziano JM, Buring JE, Lee IM. Egg consumption and risk of type 2 diabetes in men and women. *Diabetes Care* 2009;32:295-300.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Djousse L, Gaziano JM, Kase CS, Kurth T. Nut consumption and risk of stroke in US male physicians. *Clin Nutr.* 2010;29:605-9.

Djousse L, Levy D, Murabito JM, Cupples LA, Ellison RC. Alcohol consumption and risk of intermittent claudication in the Framingham Heart Study. *Circulation* 2000;102:3092-7.

Djousse L, Rudich T, Gaziano JM. Nut consumption and risk of heart failure in the Physicians' Health Study I. *Am J Clin Nutr* 2008;88:930-3.

Djousse L, Rudich T, Gaziano JM. Nut consumption and risk of hypertension in US male physicians. *Clin Nutr* 2009;28:10-4.

Dodson PM, Pacy PJ, Cox EV. Long-term follow-up of the treatment of essential hypertension with a high-fibre, low-fat and low-sodium dietary regimen. *Hum Nutr Clin Nutr.* 1985;39:213-20.

Dolecek TA, Johnson RL, Grandits GA, Farrand-Zukel M, Caggiula AW. Chapter 9. Nutritional adequacy of diets reported at baseline and during trial years 1-6 by the special intervention and usual care groups in the Multiple Risk Factor Intervention Trial. *American Journal of Clinical Nutrition* 1997;65:305S-313.

Dorant E, van den Brandt PA, Goldbohm RA, Sturmans F. Consumption of onions and a reduced risk of stomach carcinoma. *Gastroenterology* 1996;110:12-20.

Dorgan JF, Liu L, Klifa C, Hylton N, Shepherd JA, Stanczyk FZ, et al. Adolescent diet and subsequent serum hormones, breast density, and bone mineral density in young women: results of the Dietary Intervention Study in Children follow-up study. *Cancer Epidemiol Biomarkers Prev* 2010;19:1545-56.

Drenick EJ, Johnson D. Weight reduction by fasting and semistarvation in morbid obesity: long-term follow-up. *Int J Obes* 1978;2:123-32.

Drogan D, Hoffmann K, Schulz M, Bergmann MM, Boeing H, Weikert C. A food pattern predicting prospective weight change is associated with risk of fatal but not with nonfatal cardiovascular disease. *J Nutr* 2007;137:1961-7.

Drogan D, Klipstein-Grobusch K, Dierkes J, Weikert C, Boeing H. Dietary intake of folate equivalents and risk of myocardial infarction in the European Prospective Investigation into Cancer and Nutrition (EPIC)--Potsdam study. *Public Health Nutr.* 2006;9:465-71.

Du H, van der AD, Boshuizen HC, Forouhi NG, Wareham NJ, Halkjaer J, et al. Dietary fiber and subsequent changes in body weight and waist circumference in European men and women. *Am J Clin Nutr* 2010;91:329-36.

Duffey KJ, Gordon-Larsen P, Steffen LM, Jacobs DR, Jr., Popkin BM. Regular consumption from fast food establishments relative to other restaurants is differentially associated with metabolic outcomes in young adults. *J Nutr.* 2009;139:2113-8.

Dwyer JT. Dietary fat and breast cancer: testing interventions to reduce risks. *Adv Exp Med Biol* 1992;322:155-83.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Dwyer T, Calvert GD, Baghurst KI, Leitch DR. Diet, other lifestyle factors and HDL cholesterol in a population of Australian male service recruits. *Am J Epidemiol* 1981;114:683-96.

Eagan MS, Lyle RM, Gunther CW, Peacock M, Teegarden D. Effect of 1-year dairy product intervention on fat mass in young women: 6-month follow-up. *Obesity (Silver Spring)*. 2006;14:2242-8.

Earnshaw SA, Worley A, Hosking DJ. Current diet does not relate to bone mineral density after the menopause. The Nottingham Early Postmenopausal Intervention Cohort (EPIC) Study Group. *Br J Nutr*. 1997;78:65-72.

Eckel RH. Egg consumption in relation to cardiovascular disease and mortality: the story gets more complex. *Am J Clin Nutr* 2008;87:799-800.

Economos CD, Hyatt RR, Goldberg JP, Must A, Naumova EN, Collins JJ, et al. A community intervention reduces BMI z-score in children: Shape up Somerville first year results. *Obesity* 2007;15:1325-1336.

Egeberg R, Frederiksen K, Olsen A, Johnsen NF, Loft S, Overvad K, et al. Intake of wholegrain products is associated with dietary, lifestyle, anthropometric and socio-economic factors in Denmark. *Public Health Nutr*. 2009;12:1519-30.

Eichholzer M, Stahelin HB, Gey KF. Inverse correlation between essential antioxidants in plasma and subsequent risk to develop cancer, ischemic heart disease and stroke respectively: 12-year follow-up of the Prospective Basel Study. *EXS*. 1992;62:398-410.

Elias MC, Parise ER, Carvalho LD, Szejnfeld D, Netto JP. Effect of 6-month nutritional intervention on non-alcoholic fatty liver disease. *Nutrition* 2010;26:1094-1099.

Elkan AC, Sjoberg B, Kolsrud B, Ringertz B, Hafstrom I, Frostegard J. Gluten-free vegan diet induces decreased LDL and oxidized LDL levels and raised atheroprotective natural antibodies against phosphorylcholine in patients with rheumatoid arthritis: A randomized study. *SOURCE Arthritis Research and Therapy* (2008) 10:2 Article Number: R34. Date of Publication: 18 Mar 2008 2008.

Ellingsen I, Hjerkin EM, Arnesen H, Seljeflot I, Hjerkmann I, Tonstad S. Follow-up of diet and cardiovascular risk factors 20 years after cessation of intervention in the Oslo Diet and Antismoking Study. *Eur J Clin Nutr* 2006;60:378-85.

Ellingsen I, Hjerkmann I, Abdelnoor M, Hjerkin EM, Tonstad S. Dietary and antismoking advice and ischemic heart disease mortality in men with normal or high fasting triacylglycerol concentrations: a 23-y follow-up study. *Am J Clin Nutr* 2003;78:935-40.

Ellis RW, Darga LL, Lucas CP. The short- and long-term effects of a low-fat, cholesterol-free, hypocaloric diet on serum triglyceride and cholesterol distribution in severely obese humans. *Int J Obes* 1987;11:29-40.

Ellsworth JL, Kushi LH, Folsom AR. Frequent nut intake and risk of death from coronary heart disease and all causes in postmenopausal women: the Iowa Women's Health Study. *Nutr Metab Cardiovasc Dis* 2001;11:372-7.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Elmstahl S, Gullberg B, Janzon L, Johnell O, Elmstahl B. Increased incidence of fractures in middle-aged and elderly men with low intakes of phosphorus and zinc. *Osteoporos Int* 1998;8:333-40.

Elwood P, Hughes J, Fehily A. Milk, heart disease and obesity: An examination of the evidence. *British Journal of Cardiology* 2005;12:283-290.

Elwood PC, Givens DI, Beswick AD, Fehily AM, Pickering JE, Gallacher J. The survival advantage of milk and dairy consumption: an overview of evidence from cohort studies of vascular diseases, diabetes and cancer. *J Am Coll Nutr* 2008;27:723S-34S.

Engeset D, Alsaker E, Ciampi A, Lund E. Dietary patterns and lifestyle factors in the Norwegian EPIC cohort: the Norwegian Women and Cancer (NOWAC) study. *Eur J Clin Nutr*. 2005;59:675-84.

Epstein LH, Gordy CC, Raynor HA, Beddome M, Kilanowski CK, Paluch R. Increasing fruit and vegetable intake and decreasing fat and sugar intake in families at risk for childhood obesity. *Obes Res* 2001;9:171-8.

Epstein MM, Kasperzyk JL, Andren O, Giovannucci EL, Wolk A, Hakansson N, et al. Dietary zinc and prostate cancer survival in a Swedish cohort. *Am J Clin Nutr*. 2011;93:586-93.

Erkkila AT, Herrington DM, Mozaffarian D, Lichtenstein AH. Cereal fiber and whole-grain intake are associated with reduced progression of coronary-artery atherosclerosis in postmenopausal women with coronary artery disease. *Am Heart J* 2005;150:94-101.

Erkkila AT, Lehto S, Pyorala K, Uusitupa MI. n-3 Fatty acids and 5-y risks of death and cardiovascular disease events in patients with coronary artery disease. *Am J Clin Nutr* 2003;78:65-71.

Eshak ES, Iso H, Date C, Kikuchi S, Watanabe Y, Wada Y, et al. Dietary fiber intake is associated with reduced risk of mortality from cardiovascular disease among Japanese men and women. *J Nutr* 2010;140:1445-53.

Eshak ES, Iso H, Date C, Yamagishi K, Kikuchi S, Watanabe Y, et al. Rice intake is associated with reduced risk of mortality from cardiovascular disease in Japanese men but not women. *J Nutr*. 2011;141:595-602.

Esmailzadeh A, Azadbakht L. Dietary flavonoid intake and cardiovascular mortality. *Br J Nutr* 2008;100:695-7.

Esmailzadeh A, Mirmiran P, Azizi F. Whole-grain consumption and the metabolic syndrome: a favorable association in Tehranian adults. *Eur J Clin Nutr*. 2005;59:353-62.

Esposito K, Giugliano F, Maiorino MI, Giugliano D. Dietary Factors, Mediterranean Diet and Erectile Dysfunction. *Journal of Sexual Medicine* 2010;7:2338-2345.

Esposito K, Kastorini CM, Panagiotakos DB, Giugliano D. Mediterranean diet and weight loss: meta-analysis of randomized controlled trials. *Metab Syndr Relat Disord* 2011;9:1-12.

Esposito K, Maiorino MI, Ciotola M, Di Palo C, Scognamiglio P, Gicchino M, et al. Effects of a Mediterranean-style diet on the need for antihyperglycemic drug therapy in patients with newly diagnosed type 2 diabetes: a randomized trial. *Ann Intern Med* 2009;151:306-14.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Esposito K, Marfella R, Ciotola M, Di Palo C, Giugliano F, Giugliano G, et al. Effect of a Mediterranean-style diet on endothelial dysfunction and markers of vascular inflammation in the metabolic syndrome: A randomized trial. *Journal of the American Medical Association* 2004;292:1440-1446.

Esrey KL, Joseph L, Grover SA. Relationship between dietary intake and coronary heart disease mortality: lipid research clinics prevalence follow-up study. *J Clin Epidemiol* 1996;49:211-6.

Estaquio C, Druesne-Pecollo N, Latino-Martel P, Dauchet L, Hercberg S, Bertrais S. Socioeconomic differences in fruit and vegetable consumption among middle-aged French adults: adherence to the 5 A Day recommendation. *J Am Diet Assoc.* 2008;108:2021-30.

Estruch R, Martinez-Gonzalez MA, Corella D, Salas-Salvado J, Ruiz-Gutierrez V, Covas MI, et al. Effects of a Mediterranean-style diet on cardiovascular risk factors: a randomized trial. *Ann Intern Med* 2006;145:1-11.

Etminan M, Takkouche B, Caamano-Isorna F. The role of tomato products and lycopene in the prevention of prostate cancer: a meta-analysis of observational studies. *Cancer Epidemiol Biomarkers Prev* 2004;13:340-5.

Fagerberg B, Andersson OK, Isaksson B, Bjorntorp P. Blood pressure control during weight reduction in obese hypertensive men: separate effects of sodium and energy restriction. *Br Med J (Clin Res Ed)* 1984;288:11-4.

Farchi G, Fidanza F, Grossi P, Lancia A, Mariotti S, Menotti A. Relationship between eating patterns meeting recommendations and subsequent mortality in 20 years. *Eur J Clin Nutr* 1995;49:408-19.

Farchi G, Fidanza F, Mariotti S, Menotti A. Is diet an independent risk factor for mortality? 20 year mortality in the Italian rural cohorts of the Seven Countries Study. *Eur J Clin Nutr* 1994;48:19-29.

Faria SL, de Oliveira Kelly E, Lins RD, Faria OP. Nutritional management of weight regain after bariatric surgery. *Obes Surg* 2010;20:135-9.

Faria SL, Faria OP, Lopes TC, Galvao MV, de Oliveira Kelly E, Ito MK. Relation between carbohydrate intake and weight loss after bariatric surgery. *Obes Surg* 2009;19:708-16.

Fawzi W, Herrera MG, Nestel P. Tomato intake in relation to mortality and morbidity among Sudanese children. *J Nutr* 2000;130:2537-42.

Fawzi WW, Herrera MG, Willett WC, Nestel P, el Amin A, Lipsitz S, et al. Dietary vitamin A intake and the risk of mortality among children. *Am J Clin Nutr* 1994;59:401-8.

Feart C, Samieri C, Rondeau V, Amieva H, Portet F, Dartigues JF, et al. Adherence to a Mediterranean diet, cognitive decline, and risk of dementia. *JAMA.* 2009;302:638-48.

Fehily AM, Coles RJ, Evans WD, Elwood PC. Factors affecting bone density in young adults. *Am J Clin Nutr.* 1992;56:579-86.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Fehily AM, Yarnell JW, Sweetnam PM, Elwood PC. Diet and incident ischaemic heart disease: the Caerphilly Study. *Br J Nutr*. 1993;69:303-14.

Fernandez E, Negri E, La Vecchia C, Franceschi S. Diet diversity and colorectal cancer. *Prev Med* 2000;31:11-4.

Feskanich D, Weber P, Willett WC, Rockett H, Booth SL, Colditz GA. Vitamin K intake and hip fractures in women: a prospective study. *Am J Clin Nutr* 1999;69:74-9.

Feskanich D, Willett WC, Colditz GA. Calcium, vitamin D, milk consumption, and hip fractures: a prospective study among postmenopausal women. *Am J Clin Nutr* 2003;77:504-11.

Feskanich D, Willett WC, Stampfer MJ, Colditz GA. Protein consumption and bone fractures in women. *Am J Epidemiol* 1996;143:472-9.

Feskanich D, Willett WC, Stampfer MJ, Colditz GA. Milk, dietary calcium, and bone fractures in women: a 12-year prospective study. *Am J Public Health*. 1997;87:992-7.

Feskanich D, Ziegler RG, Michaud DS, Giovannucci EL, Speizer FE, Willett WC, et al. Prospective study of fruit and vegetable consumption and risk of lung cancer among men and women. *J Natl Cancer Inst* 2000;92:1812-23.

Feskens EJ, Bowles CH, Kromhout D. Association between fish intake and coronary heart disease mortality. Differences in normoglycemic and glucose intolerant elderly subjects. *Diabetes Care* 1993;16:1029-34.

Feskens EJ, Virtanen SM, Rasanen L, Tuomilehto J, Stengard J, Pekkanen J, et al. Dietary factors determining diabetes and impaired glucose tolerance. A 20-year follow-up of the Finnish and Dutch cohorts of the Seven Countries Study. *Diabetes Care* 1995;18:1104-12.

Fidanza F. Diet and incidence of coronary heart disease. *Bibl Nutr Dieta*. 1973:93-107.

Fidanza F, Alberti A, Lanti M, Menotti A. Mediterranean diet score: correlation with 25-year mortality from coronary heart disease in the Seven Countries Study. *Nutr Metab Cardiovasc Dis* 2004;14:397.

Fidanza F, Alberti A, Lanti M, Menotti A. Mediterranean Adequacy Index: correlation with 25-year mortality from coronary heart disease in the Seven Countries Study. *Nutr Metab Cardiovasc Dis* 2004;14:254-8.

Field AE, Aneja P, Austin SB, Shrier LA, de Moor C, Gordon-Larsen P. Race and gender differences in the association of dieting and gains in BMI among young adults. *Obesity (Silver Spring)* 2007;15:456-64.

Field AE, Gillman MW, Rosner B, Rockett HR, Colditz GA. Association between fruit and vegetable intake and change in body mass index among a large sample of children and adolescents in the United States. *Int J Obes Relat Metab Disord*. 2003;27:821-6.

Field AE, Willett WC, Lissner L, Colditz GA. Dietary fat and weight gain among women in the Nurses' Health Study. *Obesity (Silver Spring)* 2007;15:967-76.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Finkelstein B, Fryer BA. Meal frequency and weight reduction of young women. *Am J Clin Nutr* 1971;24:465-8.

Finley CE, Barlow CE, Halton TL, Haskell WL. Glycemic index, glycemic load, and prevalence of the metabolic syndrome in the cooper center longitudinal study. *J Am Diet Assoc* 2010;110:1820-9.

Fitzgibbon ML, Stolley MR, Schiffer L, Sharp LK, Singh V, Dyer A. Obesity Reduction Black Intervention Trial (ORBIT): 18-month results. *Obesity* 2010;18:2317-2325.

Fleischauer AT, Arab L. Garlic and cancer: a critical review of the epidemiologic literature. *J Nutr* 2001;131:1032S-40S.

Fleming DJ, Tucker KL, Jacques PF, Dallal GE, Wilson PW, Wood RJ. Dietary factors associated with the risk of high iron stores in the elderly Framingham Heart Study cohort. *Am J Clin Nutr* 2002;76:1375-84.

Flood A, Caprario L, Chatterjee N, Lacey JV, Jr., Schairer C, Schatzkin A. Folate, methionine, alcohol, and colorectal cancer in a prospective study of women in the United States. *Cancer Causes Control* 2002;13:551-61.

Flood A, Mitchell N, Jaeb M, Finch EA, Laqua PS, Welsh EM, et al. Energy density and weight change in a long-term weight-loss trial. *The International Journal of Behavioral Nutrition and Physical Activity* 2009;6.

Flood A, Velie EM, Chatterjee N, Subar AF, Thompson FE, Lacey JV, Jr., et al. Fruit and vegetable intakes and the risk of colorectal cancer in the Breast Cancer Detection Demonstration Project follow-up cohort. *Am J Clin Nutr* 2002;75:936-43.

Flood V, Smith W, Wang JJ, Manzi F, Webb K, Mitchell P. Dietary antioxidant intake and incidence of early age-related maculopathy: the Blue Mountains Eye Study. *Ophthalmology* 2002;109:2272-8.

Flores-Mateo G, Navas-Acien A, Pastor-Barriuso R, Guallar E. Selenium and coronary heart disease: a meta-analysis. *Am J Clin Nutr* 2006;84:762-73.

Flynn MA, Nolph GB, Baker AS, Krause G. Aging in humans: a continuous 20-year study of physiologic and dietary parameters. *J Am Coll Nutr* 1992;11:660-72.

Folsom AR, Demissie Z. Fish intake, marine omega-3 fatty acids, and mortality in a cohort of postmenopausal women. *Am J Epidemiol* 2004;160:1005-10.

Folsom AR, Parker ED, Harnack LJ. Degree of concordance with DASH diet guidelines and incidence of hypertension and fatal cardiovascular disease. *Am J Hypertens* 2007;20:225-32.

Fontaine KR, Yang D, Gadbury GL, Heshka S, Schwartz LG, Murugesan R, et al. Results of soy-based meal replacement formula on weight, anthropometry, serum lipids & blood pressure during a 40-week clinical weight loss trial. *SOURCE Nutrition Journal* (2003) 2 (1-7). Date of Publication: 18 Nov 2003 2003.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Ford AL, Bergh C, Södersten P, Sabin MA, Hollinghurst S, Hunt LP, et al. Treatment of childhood obesity by retraining eating behaviour: Randomised controlled trial. *BMJ: British Medical Journal* 2010;340.

Ford ES, Mokdad AH. Fruit and vegetable consumption and diabetes mellitus incidence among U.S. adults. *Prev Med* 2001;32:33-9.

Forlani G, Lorusso C, Moscatiello S, Ridolfi V, Melchionda N, Di Domizio S, et al. Are behavioural approaches feasible and effective in the treatment of type 2 diabetes? A propensity score analysis vs. prescriptive diet. *Nutr Metab Cardiovasc Dis.* 2009;19:313-20.

Forouhi NG, Sharp SJ, Du H, van der AD, Halkjaer J, Schulze MB, et al. Dietary fat intake and subsequent weight change in adults: results from the European Prospective Investigation into Cancer and Nutrition cohorts. *Am J Clin Nutr* 2009;90:1632-41.

Forrest J, Campbell P, Kreiger N, Sloan M. Salivary gland cancer: an exploratory analysis of dietary factors. *Nutr Cancer* 2008;60:469-73.

Foster GD, Wyatt HR, Hill JO, Makris AP, Rosenbaum DL, Brill C, et al. Weight and metabolic outcomes after 2 years on a low-carbohydrate versus low-fat diet. *Obstetrical and Gynecological Survey* 2010;65:769-770.

Fowke JH, Shu XO, Dai Q, Shintani A, Conaway CC, Chung FL, et al. Urinary isothiocyanate excretion, brassica consumption, and gene polymorphisms among women living in Shanghai, China. *Cancer Epidemiol Biomarkers Prev* 2003;12:1536-9.

Franceschi S. Micronutrients and breast cancer. *Eur J Cancer Prev* 1997;6:535-9.

Franceschi S, Dal Maso L, Augustin L, Negri E, Parpinel M, Boyle P, et al. Dietary glycemic load and colorectal cancer risk. *Ann Oncol* 2001;12:173-8.

Franceschi S, Schinella D, Bidoli E, Dal Maso L, La Vecchia C, Parazzini F, et al. The influence of body size, smoking, and diet on bone density in pre- and postmenopausal women. *Epidemiology.* 1996;7:411-4.

Frankel S, Gunnell DJ, Peters TJ, Maynard M, Davey Smith G. Childhood energy intake and adult mortality from cancer: the Boyd Orr Cohort Study. *BMJ* 1998;316:499-504.

Franko DL, Albertson AM, Thompson DR, Barton BA. Cereal consumption and indicators of cardiovascular risk in adolescent girls. *Public Health Nutr.* 2011;14:584-90.

Fraser GE. Diet as primordial prevention in Seventh-Day Adventists. *Prev Med* 1999;29:S18-23.

Fraser GE. Vegetarian diets: what do we know of their effects on common chronic diseases? *Am J Clin Nutr* 2009;89:1607S-1612S.

Freeland KR, Wilson C, Wolever TMS. Adaptation of colonic fermentation and glucagon-like peptide-1 secretion with increased wheat fibre intake for 1 year in hyperinsulinaemic human subjects. *British Journal of Nutrition* 2010;103:82-90.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Frentzel-Beyme R, Chang-Claude J. Vegetarian diets and colon cancer: The German experience. *American Journal of Clinical Nutrition* 1994;59:1143S-1152S.

Fretts AM, Howard BV, McKnight B, Duncan GE, Beresford SA, Mete M, et al. Associations of processed meat and unprocessed red meat intake with incident diabetes: the Strong Heart Family Study. *Am J Clin Nutr* 2012;95:752-8.

Frisch S, Zittermann A, Berthold HK, Gotting C, Kuhn J, Kleesiek K, et al. A randomized controlled trial on the efficacy of carbohydrate-reduced or fat-reduced diets in patients attending a telemedically guided weight loss program. *SOURCE Cardiovascular Diabetology* (2009) 8 Article Number: 36. Date of Publication: 18 Jul 2009 2009.

Frost L, Vestergaard P. n-3 Fatty acids consumed from fish and risk of atrial fibrillation or flutter: the Danish Diet, Cancer, and Health Study. *Am J Clin Nutr* 2005;81:50-4.

Fuchs CS, Giovannucci EL, Colditz GA, Hunter DJ, Stampfer MJ, Rosner B, et al. Dietary fiber and the risk of colorectal cancer and adenoma in women. *N Engl J Med* 1999;340:169-76.

Fuller NR, Lau NS, Denyer G, Caterson ID. A 12-month, randomised, controlled trial to examine the efficacy of the Korean diet in an Australian overweight and obese population - A follow up analysis. *Obesity Research and Clinical Practice* 2012;6:e298-e307.

Fulton JE, Dai S, Steffen LM, Grunbaum JA, Shah SM, Labarthe DR. Physical activity, energy intake, sedentary behavior, and adiposity in youth. *Am J Prev Med*. 2009;37:S40-9.

Fumeron F, Lamri A, Emery N, Bellili N, Jaziri R, Porchay-Balderelli I, et al. Dairy products and the metabolic syndrome in a prospective study, DESIR. *J Am Coll Nutr* 2011;30:454S-63S.

Fung TT, Chiuve SE, McCullough ML, Rexrode KM, Logroscino G, Hu FB. Adherence to a DASH-style diet and risk of coronary heart disease and stroke in women. *Arch Intern Med* 2008;168:713-20.

Fung TT, Hu FB, Wu K, Chiuve SE, Fuchs CS, Giovannucci E. The Mediterranean and Dietary Approaches to Stop Hypertension (DASH) diets and colorectal cancer. *Am J Clin Nutr* 2010;92:1429-35.

Fung TT, Hunter DJ, Spiegelman D, Colditz GA, Speizer FE, Willett WC. Vitamins and carotenoids intake and the risk of basal cell carcinoma of the skin in women (United States). *Cancer Causes Control* 2002;13:221-30.

Fung TT, McCullough M, van Dam RM, Hu FB. A prospective study of overall diet quality and risk of type 2 diabetes in women. *Diabetes Care* 2007;30:1753-7.

Fung TT, Rexrode KM, Mantzoros CS, Manson JE, Willett WC, Hu FB. Mediterranean diet and incidence of and mortality from coronary heart disease and stroke in women. *Circulation* 2009;119:1093-100.

Fung TT, Schulze M, Manson JE, Willett WC, Hu FB. Dietary patterns, meat intake, and the risk of type 2 diabetes in women. *Arch Intern Med* 2004;164:2235-40.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Fung TT, van Dam RM, Hankinson SE, Stampfer M, Willett WC, Hu FB. Low-carbohydrate diets and all-cause and cause-specific mortality: two cohort studies. *Ann Intern Med* 2010;153:289-98.

Furberg AS, Thune I. Energy intake, energy expenditure and BMI influence the risk of endometrial cancer in a prospective study in Norway. *IARC Sci Publ.* 2002;156:247-8.

Gallus S, Bosetti C, La Vecchia C. Mediterranean diet and cancer risk. *Eur J Cancer Prev* 2004;13:447-52.

Gandini S, Merzenich H, Robertson C, Boyle P. Meta-analysis of studies on breast cancer risk and diet: the role of fruit and vegetable consumption and the intake of associated micronutrients. *Eur J Cancer* 2000;36:636-46.

Garrow JS. Treatment of morbid obesity by nonsurgical means: diet, drugs, behavior modification, exercise. *Gastroenterol Clin North Am* 1987;16:443-9.

Gartside PS, Glueck CJ. The important role of modifiable dietary and behavioral characteristics in the causation and prevention of coronary heart disease hospitalization and mortality: the prospective NHANES I follow-up study. *J Am Coll Nutr* 1995;14:71-9.

Gast GC, de Roos NM, Sluijs I, Bots ML, Beulens JW, Geleijnse JM, et al. A high menaquinone intake reduces the incidence of coronary heart disease. *Nutr Metab Cardiovasc Dis* 2009;19:504-10.

Gates MA, Tworoger SS, Hecht JL, De Vivo I, Rosner B, Hankinson SE. A prospective study of dietary flavonoid intake and incidence of epithelial ovarian cancer. *Int J Cancer* 2007;121:2225-32.

Gaziano JM, Manson JE, Branch LG, Colditz GA, Willett WC, Buring JE. A prospective study of consumption of carotenoids in fruits and vegetables and decreased cardiovascular mortality in the elderly. *Ann Epidemiol* 1995;5:255-60.

Gaziano JM, Manson JE, Buring JE, Hennekens CH. Dietary antioxidants and cardiovascular disease. *Ann N Y Acad Sci* 1992;669:249-58; discussion 258-9.

Genkinger JM, Hunter DJ, Spiegelman D, Anderson KE, Buring JE, Freudenheim JL, et al. Alcohol intake and ovarian cancer risk: a pooled analysis of 10 cohort studies. *Br J Cancer*. 2006;94:757-62.

Genkinger JM, Platz EA, Hoffman SC, Comstock GW, Helzlsouer KJ. Fruit, vegetable, and antioxidant intake and all-cause, cancer, and cardiovascular disease mortality in a community-dwelling population in Washington County, Maryland. *Am J Epidemiol* 2004;160:1223-33.

George SM, Mayne ST, Leitzmann MF, Park Y, Schatzkin A, Flood A, et al. Dietary glycemic index, glycemic load, and risk of cancer: a prospective cohort study. *Am J Epidemiol.* 2009;169:462-72.

Gibbs KE. The dangers of broccoli. *Surg Obes Relat Dis* 2010;6:439-40.

Gilbertson HR, Brand-Miller JC, Thorburn AW, Evans S, Chondros P, Werther GA. The effect of flexible low glycemic index dietary advice versus measured carbohydrate exchange diets on glycemic control in children with type 1 diabetes. *Diabetes Care* 2001;24:1137-43.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Giles GG, Simpson JA, English DR, Hodge AM, Gertig DM, Macinnis RJ, et al. Dietary carbohydrate, fibre, glycaemic index, glycaemic load and the risk of postmenopausal breast cancer. *Int J Cancer* 2006;118:1843-7.

Gilhooly CH, Das SK, Golden JK, McCrory MA, Rochon J, DeLany JP, et al. Use of cereal fiber to facilitate adherence to a human caloric restriction program. *Aging Clin Exp Res* 2008;20:513-20.

Gillen LJ, Tapsell LC, Patch CS, Owen A, Batterham M. Structured dietary advice incorporating walnuts achieves optimal fat and energy balance in patients with type 2 diabetes mellitus. *J Am Diet Assoc* 2005;105:1087-96.

Gillman MW, Cupples LA, Gagnon D, Posner BM, Ellison RC, Castelli WP, et al. Protective effect of fruits and vegetables on development of stroke in men. *JAMA* 1995;273:1113-7.

Gillum RF, Mussolino ME, Madans JH. Fish consumption and hypertension incidence in African Americans and whites: the NHANES I Epidemiologic Follow-up Study. *J Natl Med Assoc* 2001;93:124-8.

Gilsing AM, Weijenberg MP, Hughes LA, Ambergen T, Dagnelie PC, Goldbohm RA, et al. Longitudinal changes in BMI in older adults are associated with meat consumption differentially, by type of meat consumed. *J Nutr* 2012;142:340-9.

Giordano G, Guarini P, Ferrari P, Biondi-Zoccai G, Schiavone B, Giordano A. Beneficial impact on cardiovascular risk profile of water buffalo meat consumption. *European Journal of Clinical Nutrition* 2010;64:1000-1006.

Giovannucci E, Rimm EB, Liu Y, Stampfer MJ, Willett WC. A prospective study of tomato products, lycopene, and prostate cancer risk. *J Natl Cancer Inst* 2002;94:391-8.

Giovannucci E, Rimm EB, Liu Y, Stampfer MJ, Willett WC. A prospective study of cruciferous vegetables and prostate cancer. *Cancer Epidemiol Biomarkers Prev* 2003;12:1403-9.

Giugliano D, Ceriello A, Esposito K. The Effects of Diet on Inflammation. Emphasis on the Metabolic Syndrome. *Journal of the American College of Cardiology* 2006;48:677-685.

Giugliano D, Ceriello A, Esposito K. Mediterranean diet inversely associated with the incidence of metabolic syndrome: The sun prospective cohort Response to tortosa et al. *Diabetes Care* 2008;31:E36.

Giugliano F, Maiorino MI, Di Palo C, Autorino R, De Sio M, Giugliano D, et al. Adherence to mediterranean diet and sexual function in women with type 2 diabetes. *Journal of Sexual Medicine* 2010;7:1883-1890.

Goldbohm RA, Chorus AM, Galindo Garre F, Schouten LJ, van den Brandt PA. Dairy consumption and 10-y total and cardiovascular mortality: a prospective cohort study in the Netherlands. *Am J Clin Nutr*. 2011;93:615-27.

Golley RK, Magarey AM, Daniels LA. Children's food and activity patterns following a six-month child weight management program. *Int J Pediatr Obes* 2011;6:409-14.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Gong Q, Gregg EW, Wang J, An Y, Zhang P, Yang W, et al. Long-term effects of a randomised trial of a 6-year lifestyle intervention in impaired glucose tolerance on diabetes-related microvascular complications: The China da Qing Diabetes Prevention Outcome Study. *Diabetologia* 2011;54:300-307.

Gong Z, Holly EA, Wang F, Chan JM, Bracci PM. Intake of fatty acids and antioxidants and pancreatic cancer in a large population-based case-control study in the San Francisco Bay Area. *Int J Cancer*. 2010;127:1893-904.

Gonzalez A, Peters U, Lampe JW, White E. Boron intake and prostate cancer risk. *Cancer Causes Control* 2007;18:1131-40.

Gonzalez CA, Riboli E. Diet and cancer prevention: where we are, where we are going. *Nutr Cancer* 2006;56:225-31.

Gonzalez CA, Riboli E. Diet and cancer prevention: Contributions from the European Prospective Investigation into Cancer and Nutrition (EPIC) study. *Eur J Cancer* 2010;46:2555-62.

Gonzalez S, Huerta JM, Fernandez S, Patterson AM, Lasheras C. Differences in overall mortality in the elderly may be explained by diet. *Gerontology* 2008;54:232-7.

Goodwin JS, Brodwick M. Diet, aging, and cancer. *Clin Geriatr Med* 1995;11:577-89.

Goodwin PJ, Ennis M, Pritchard KI, Koo J, Trudeau ME, Hood N. Diet and breast cancer: evidence that extremes in diet are associated with poor survival. *J Clin Oncol*. 2003;21:2500-7.

Goulet J, Lapointe A, Lamarche B, Lemieux S. Effect of a nutritional intervention promoting the Mediterranean food pattern on anthropometric profile in healthy women from the Quebec city metropolitan area. *European Journal of Clinical Nutrition* 2007;61:1293-1300.

Goyenechea E, Holst C, Saris WH, Jebb S, Kafatos A, Pfeiffer A, et al. Effects of different protein content and glycemic index of ad libitum diets on diabetes risk factors in overweight adults: the DIOGenes multicentre, randomised, dietary intervention trial. *Diabetes Metab Res Rev* 2011.

Graham S, Zielezny M, Marshall J, Priore R, Freudenheim J, Brasure J, et al. Diet in the epidemiology of postmenopausal breast cancer in the New York State Cohort. *Am J Epidemiol*. 1992;136:1327-37.

Greenwald P, Sondik E, Lynch BS. Diet and chemoprevention in NCI's research strategy to achieve national cancer control objectives. *Annu Rev Public Health* 1986;7:267-91.

Greenwood DC, Cade JE, Draper A, Barrett JH, Calvert C, Greenhalgh A. Seven unique food consumption patterns identified among women in the UK Women's Cohort Study. *Eur J Clin Nutr*. 2000;54:314-20.

Griffiths RA, Holliday J. An evaluation and follow-up investigation of a behavioural group treatment programme for obesity. *Psychother Psychosom* 1987;48:157-64.

Gu Y, Nieves JW, Stern Y, Luchsinger JA, Scarmeas N. Food combination and Alzheimer disease risk: a protective diet. *Arch Neurol*. 2010;67:699-706.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Gunnell DJ, Frankel S, Nanchahal K, Braddon FE, Smith GD. Lifecourse exposure and later disease: a follow-up study based on a survey of family diet and health in pre-war Britain (1937-1939). *Public Health* 1996;110:85-94.

Gunther AL, Buyken AE, Kroke A. The influence of habitual protein intake in early childhood on BMI and age at adiposity rebound: results from the DONALD Study. *Int J Obes (Lond)*. 2006;30:1072-9.

Gunther AL, Remer T, Kroke A, Buyken AE. Early protein intake and later obesity risk: which protein sources at which time points throughout infancy and childhood are important for body mass index and body fat percentage at 7 y of age? *Am J Clin Nutr* 2007;86:1765-72.

Gunther CW, Legowski PA, Lyle RM, McCabe GP, Eagan MS, Peacock M, et al. Dairy products do not lead to alterations in body weight or fat mass in young women in a 1-y intervention. *American Journal of Clinical Nutrition* 2005;81:751-756.

Hadden DR, Blair ALT, Wilson EA. Natural history of diabetes presenting age 40-69 years: A prospective study of the influence of intensive dietary therapy. *Quarterly Journal of Medicine* 1986;59:579-598.

Hailu A, Knutsen SF, Fraser GE. Associations between meat consumption and the prevalence of degenerative arthritis and soft tissue disorders in the adventist health study, California U.S.A. *J Nutr Health Aging*. 2006;10:7-14.

Haimoto H, Iwata M, Wakai K, Umegaki H. Long-term effects of a diet loosely restricting carbohydrates on HbA1c levels, BMI and tapering of sulfonylureas in type 2 diabetes: a 2-year follow-up study. *Diabetes Res Clin Pract*. 2008;79:350-6.

Halton TL, Liu S, Manson JE, Hu FB. Low-carbohydrate-diet score and risk of type 2 diabetes in women. *Am J Clin Nutr* 2008;87:339-46.

Halton TL, Willett WC, Liu S, Manson JE, Albert CM, Rexrode K, et al. Low-carbohydrate-diet score and the risk of coronary heart disease in women. *N Engl J Med* 2006;355:1991-2002.

Hamer M, Chida Y. Intake of fruit, vegetables, and antioxidants and risk of type 2 diabetes: systematic review and meta-analysis. *J Hypertens* 2007;25:2361-9.

Handjieva-Darlenska T, Handjiev S, Larsen TM, van Baak MA, Lindroos A, Papadaki A, et al. Predictors of weight loss maintenance and attrition during a 6-month dietary intervention period: Results from the diogenes study. *Clinical Obesity* 2011;1:62-68.

Hankin JH. Role of nutrition in women's health: diet and breast cancer. *J Am Diet Assoc* 1993;93:994-9.

Hannan MT, Tucker KL, Dawson-Hughes B, Cupples LA, Felson DT, Kiel DP. Effect of dietary protein on bone loss in elderly men and women: the Framingham Osteoporosis Study. *J Bone Miner Res*. 2000;15:2504-12.

Hansen L, Dragsted LO, Olsen A, Christensen J, Tjønneland A, Schmidt EB, et al. Fruit and vegetable intake and risk of acute coronary syndrome. *Br J Nutr* 2010;104:248-55.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Hardcastle AC, Aucott L, Fraser WD, Reid DM, Macdonald HM. Dietary patterns, bone resorption and bone mineral density in early post-menopausal Scottish women. *Eur J Clin Nutr*. 2011;65:378-85.

Harding AH, Day NE, Khaw KT, Bingham S, Luben R, Welsh A, et al. Dietary fat and the risk of clinical type 2 diabetes: the European prospective investigation of Cancer-Norfolk study. *Am J Epidemiol* 2004;159:73-82.

Hardy DS, Hoelscher DM, Aragaki C, Stevens J, Steffen LM, Pankow JS, et al. Association of glycemic index and glycemic load with risk of incident coronary heart disease among Whites and African Americans with and without type 2 diabetes: the Atherosclerosis Risk in Communities study. *Ann Epidemiol* 2010;20:610-6.

Harriss LR, English DR, Powles J, Giles GG, Tonkin AM, Hodge AM, et al. Dietary patterns and cardiovascular mortality in the Melbourne Collaborative Cohort Study. *Am J Clin Nutr* 2007;86:221-9.

Hartman TJ, Albert PS, Zhang Z, Bagshaw D, Kris-Etherton PM, Ulbrecht J, et al. Consumption of a legume-enriched, low-glycemic index diet is associated with biomarkers of insulin resistance and inflammation among men at risk for colorectal cancer. *Journal of Nutrition* 2010;140:60-67.

Harvey-Berino J, Gold BC, Lauber R, Starinski A. The impact of calcium and dairy product consumption on weight loss. *Obes Res* 2005;13:1720-6.

Haufe S, Engeli S, Kast P, Bohnke J, Utz W, Haas V, et al. Randomized comparison of reduced fat and reduced carbohydrate hypocaloric diets on intrahepatic fat in overweight and obese human subjects. *Hepatology* 2011;53:1504-14.

Haugen MA, Kjeldsen-Kragh J, Bjerve KS, Hostmark AT, Forre O. Changes in plasma phospholipid fatty acids and their relationship to disease activity in rheumatoid arthritis patients treated with a vegetarian diet. *Br J Nutr*. 1994;72:555-66.

Havas S, Anliker J, Damron D, Langenberg P, Ballesteros M, Feldman R. Final results of the Maryland WIC 5-A-Day Promotion Program. *Am J Public Health* 1998;88:1161-7.

Havas S, Anliker J, Greenberg D, Block G, Block T, Blik C, et al. Final results of the Maryland WIC Food for Life Program. *Prev Med* 2003;37:406-16.

Haveman-Nies A, de Groot LP, Burema J, Cruz JA, Osler M, van Staveren WA. Dietary quality and lifestyle factors in relation to 10-year mortality in older Europeans: the SENECA study. *Am J Epidemiol* 2002;156:962-8.

He FJ, Nowson CA, Lucas M, MacGregor GA. Increased consumption of fruit and vegetables is related to a reduced risk of coronary heart disease: meta-analysis of cohort studies. *J Hum Hypertens* 2007;21:717-28.

He FJ, Nowson CA, MacGregor GA. Fruit and vegetable consumption and stroke: meta-analysis of cohort studies. *Lancet* 2006;367:320-6.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

He K, Hu FB, Colditz GA, Manson JE, Willett WC, Liu S. Changes in intake of fruits and vegetables in relation to risk of obesity and weight gain among middle-aged women. *Int J Obes Relat Metab Disord* 2004;28:1569-74.

He K, Liu K, Daviglus ML, Jenny NS, Mayer-Davis E, Jiang R, et al. Associations of dietary long-chain n-3 polyunsaturated fatty acids and fish with biomarkers of inflammation and endothelial activation (from the Multi-Ethnic Study of Atherosclerosis). *Am J Cardiol*. 2009;103:1238-43.

He K, Song Y, Daviglus ML, Liu K, Van Horn L, Dyer AR, et al. Accumulated evidence on fish consumption and coronary heart disease mortality: a meta-analysis of cohort studies. *Circulation* 2004;109:2705-11.

He M, van Dam RM, Rimm E, Hu FB, Qi L. Whole-grain, cereal fiber, bran, and germ intake and the risks of all-cause and cardiovascular disease-specific mortality among women with type 2 diabetes mellitus. *Circulation* 2010;121:2162-8.

Hebden L, Chey T, Allman-Farinelli M. Lifestyle intervention for preventing weight gain in young adults: a systematic review and meta-analysis of RCTs. *Obes Rev* 2012;13:692-710.

Heidemann C, Hoffmann K, Spranger J, Klipstein-Grobusch K, Mohlig M, Pfeiffer AF, et al. A dietary pattern protective against type 2 diabetes in the European Prospective Investigation into Cancer and Nutrition (EPIC)--Potsdam Study cohort. *Diabetologia* 2005;48:1126-34.

Heidemann C, Schulze MB, Franco OH, van Dam RM, Mantzoros CS, Hu FB. Dietary patterns and risk of mortality from cardiovascular disease, cancer, and all causes in a prospective cohort of women. *Circulation* 2008;118:230-7.

Heinen MM, Verhage BA, Goldbohm RA, van den Brandt PA. Intake of vegetables, fruits, carotenoids and vitamins C and E and pancreatic cancer risk in The Netherlands Cohort Study. *Int J Cancer* 2012;130:147-58.

Heitmann BL, Lissner L, Sorensen TI, Bengtsson C. Dietary fat intake and weight gain in women genetically predisposed for obesity. *Am J Clin Nutr* 1995;61:1213-7.

Helmersson J, Arnlov J, Larsson A, Basu S. Low dietary intake of beta-carotene, alpha-tocopherol and ascorbic acid is associated with increased inflammatory and oxidative stress status in a Swedish cohort. *Br J Nutr*. 2009;101:1775-82.

Henes ST, Collier DN, Morrissey SL, Cummings DM, Kolasa KM. Medical nutrition therapy for overweight youth in their medical home: the KIDPOWER experience. *Patient Educ Couns* 2010;81:43-6.

Heo M, Kim RS, Wylie-Rosett J, Allison DB, Heymsfield SB, Faith MS. Inverse association between fruit and vegetable intake and BMI even after controlling for demographic, socioeconomic and lifestyle factors. *Obes Facts* 2011;4:449-55.

Heroux M, Janssen I, Lam M, Lee DC, Hebert JR, Sui X, et al. Dietary patterns and the risk of mortality: impact of cardiorespiratory fitness. *Int J Epidemiol*. 2010;39:197-209.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Hertog MG, Bueno-de-Mesquita HB, Fehily AM, Sweetnam PM, Elwood PC, Kromhout D. Fruit and vegetable consumption and cancer mortality in the Caerphilly Study. *Cancer Epidemiol Biomarkers Prev* 1996;5:673-7.

Hertog MG, Feskens EJ, Hollman PC, Katan MB, Kromhout D. Dietary antioxidant flavonoids and risk of coronary heart disease: the Zutphen Elderly Study. *Lancet* 1993;342:1007-11.

Hertog MG, Feskens EJ, Hollman PC, Katan MB, Kromhout D. Dietary flavonoids and cancer risk in the Zutphen Elderly Study. *Nutr Cancer* 1994;22:175-84.

Hertog MG, Kromhout D, Aravanis C, Blackburn H, Buzina R, Fidanza F, et al. Flavonoid intake and long-term risk of coronary heart disease and cancer in the seven countries study. *Arch Intern Med* 1995;155:381-6.

Hill HA, Austin H. Nutrition and endometrial cancer. *Cancer Causes Control* 1996;7:19-32.

Hinton PS, Rector RS, Donnelly JE, Smith BK, Bailey B. Total body bone mineral content and density during weight loss and maintenance on a low- or recommended-dairy weight-maintenance diet in obese men and women. *Eur J Clin Nutr* 2010;64:392-9.

Hirose K, Matsuo K, Iwata H, Tajima K. Dietary patterns and the risk of breast cancer in Japanese women. *Cancer Sci* 2007;98:1431-8.

Hjartaker A, Laake P, Lund E. Childhood and adult milk consumption and risk of premenopausal breast cancer in a cohort of 48,844 women - the Norwegian women and cancer study. *Int J Cancer* 2001;93:888-93.

Hjerkinn EM, Sandvik L, Hjerermann I, Arnesen H. Effect of diet intervention on long-term mortality in healthy middle-aged men with combined hyperlipidaemia. *Journal of Internal Medicine* 2004;255:68-73.

Ho SC, Chan SG, Yi Q, Wong E, Leung PC. Soy intake and the maintenance of peak bone mass in Hong Kong Chinese women. *J Bone Miner Res.* 2001;16:1363-9.

Hodge AM, English DR, O'Dea K, Giles GG. Glycemic index and dietary fiber and the risk of type 2 diabetes. *Diabetes Care* 2004;27:2701-6.

Hodge AM, English DR, O'Dea K, Giles GG. Dietary patterns and diabetes incidence in the Melbourne Collaborative Cohort Study. *Am J Epidemiol* 2007;165:603-10.

Hodgson JM, Devine A, Burke V, Dick IM, Prince RL. Chocolate consumption and bone density in older women. *Am J Clin Nutr.* 2008;87:175-80.

Hodson L, Harnden KE, Roberts R, Dennis AL, Frayn KN. Does the DASH diet lower blood pressure by altering peripheral vascular function. *Journal of Human Hypertension* 2010;24:312-319.

Hooper L. Dietetic guidelines: Diet in secondary prevention of cardiovascular disease. *Journal of Human Nutrition and Dietetics* 2001;14:297-305.

Hooper L, Thompson RL, Harrison RA, Summerbell CD, Ness AR, Moore HJ, et al. Risks and benefits of omega 3 fats for mortality, cardiovascular disease, and cancer: systematic review. *BMJ* 2006;332:752-60.

Hopping BN, Erber E, Grandinetti A, Verheus M, Kolonel LN, Maskarinec G. Dietary fiber, magnesium, and glycemic load alter risk of type 2 diabetes in a multiethnic cohort in Hawaii. *J Nutr* 2010;140:68-74.

Horrocks PM, Blackmore R, Wright AD. A long-term follow-up of dietary advice in maturity onset diabetes: the experience of one centre in the UK prospective study. *Diabet Med* 1987;4:241-4.

Houtsmuller AJ, Zahn KJ, Henkes HE. Unsaturated fats and progression of diabetic retinopathy. *Doc Ophthalmol* 1980;48:363-71.

Howard BV, Abbott WG, Swinburn BA. Evaluation of metabolic effects of substitution of complex carbohydrates for saturated fat in individuals with obesity and NIDDM. *Diabetes Care* 1991;14:786-95.

Howard BV, Van Horn L, Hsia J, Manson JE, Stefanick ML, Wassertheil-Smoller S, et al. Low-fat dietary pattern and risk of cardiovascular disease: the Women's Health Initiative Randomized Controlled Dietary Modification Trial. *JAMA* 2006;295:655-66.

Howard-Williams J, Patel P, Jelfs R, Carter RD, Awdry P, Bron A, et al. Polyunsaturated fatty acids and diabetic retinopathy. *Br J Ophthalmol* 1985;69:15-8.

Howarth NC, Murphy SP, Wilkens LR, Hankin JH, Kolonel LN. Dietary energy density is associated with overweight status among 5 ethnic groups in the multiethnic cohort study. *J Nutr* 2006;136:2243-8.

Hsieh LJ, Carter HB, Landis PK, Tucker KL, Metter EJ, Newschaffer CJ, et al. Association of energy intake with prostate cancer in a long-term aging study: Baltimore Longitudinal Study of Aging (United States). *Urology*. 2003;61:297-301.

Hu EA, Pan A, Malik V, Sun Q. White rice consumption and risk of type 2 diabetes: meta-analysis and systematic review. *BMJ* 2012;344:e1454.

Hu FB, Cho E, Rexrode KM, Albert CM, Manson JE. Fish and long-chain omega-3 fatty acid intake and risk of coronary heart disease and total mortality in diabetic women. *Circulation* 2003;107:1852-7.

Hu FB, Rimm EB, Stampfer MJ, Ascherio A, Spiegelman D, Willett WC. Prospective study of major dietary patterns and risk of coronary heart disease in men. *Am J Clin Nutr* 2000;72:912-21.

Hu J, La Vecchia C, DesMeules M, Negri E, Mery L. Nutrient and fiber intake and risk of renal cell carcinoma. *Nutr Cancer* 2008;60:720-8.

Huijbregts P, Feskens E, Rasanen L, Fidanza F, Nissinen A, Menotti A, et al. Dietary pattern and 20 year mortality in elderly men in Finland, Italy, and The Netherlands: longitudinal cohort study. *BMJ* 1997;315:13-7.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Hung HC, Merchant A, Willett W, Ascherio A, Rosner BA, Rimm E, et al. The association between fruit and vegetable consumption and peripheral arterial disease. *Epidemiology* 2003;14:659-65.

Hung T, Sievenpiper JL, Marchie A, Kendall CWC, Jenkins DJA. Fat versus carbohydrate in insulin resistance, obesity, diabetes and cardiovascular disease. *Current Opinion in Clinical Nutrition and Metabolic Care* 2003;6:165-176.

Hunninghake DB, Maki KC, Kwiterovich Jr PO, Davidson MH, Dicklin MR, Kafonek SD. Incorporation of lean red meat into a national cholesterol education program Step I diet: A long-term, randomized clinical trial in free-living persons with hypercholesterolemia. *Journal of the American College of Nutrition* 2000;19:351-360.

Hunter DJ, Willett WC. Diet, body size, and breast cancer. *Epidemiol Rev.* 1993;15:110-32.

Huntley AL, Ernst E. Soy for the treatment of perimenopausal symptoms - A systematic review. *Maturitas* 2004;47:1-9.

Huse DM, Branes LA, Colligan RC, Nelson RA, Palumbo PJ. The challenge of obesity in childhood. I. Incidence, prevalence, and staging. *Mayo Clin Proc* 1982;57:279-84.

Hwang JH, Kim MK, Lee JK. Dietary supplements reduce the risk of cervical intraepithelial neoplasia. *Int J Gynecol Cancer* 2010;20:398-403.

Hylander B, Rossner S. Three year's follow-up of members of a Swedish commercial weight-reducing club. *Acta Med Scand* 1981;210:485-8.

Iestra J, Knoops K, Kromhout D, de Groot L, Grobbee D, van Staveren W. Lifestyle, Mediterranean diet and survival in European post-myocardial infarction patients. *Eur J Cardiovasc Prev Rehabil* 2006;13:894-900.

Imamura F, Jacques PF, Herrington DM, Dallal GE, Lichtenstein AH. Adherence to 2005 Dietary Guidelines for Americans is associated with a reduced progression of coronary artery atherosclerosis in women with established coronary artery disease. *Am J Clin Nutr* 2009;90:193-201.

Iribarren C, Markovitz JH, Jacobs DR, Jr., Schreiner PJ, Daviglius M, Hibbeln JR. Dietary intake of n-3, n-6 fatty acids and fish: relationship with hostility in young adults--the CARDIA study. *Eur J Clin Nutr* 2004;58:24-31.

Iscovich JM, Iscovich RB, Howe G, Shiboski S, Kaldor JM. A case-control study of diet and breast cancer in Argentina. *Int J Cancer* 1989;44:770-6.

Islami F, Malekshah AF, Kimiagar M, Pourshams A, Wakefield J, Gogiani G, et al. Patterns of food and nutrient consumption in northern Iran, a high-risk area for esophageal cancer. *Nutr Cancer.* 2009;61:475-83.

Iso H, Kubota Y. Nutrition and disease in the Japan Collaborative Cohort Study for Evaluation of Cancer (JACC). *Asian Pac J Cancer Prev* 2007;8 Suppl:35-80.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Iwasaki M, Yoshihara A, Moynihan P, Watanabe R, Taylor GW, Miyazaki H. Longitudinal relationship between dietary (omega)-3 fatty acids and periodontal disease. *Nutrition* 2010;26:1105-1109.

Jacobs DR, Jr., Hohe C, Mursu J, Robien K, Folsom AR. Whole grain intake, incident hip fracture and presumed frailty in the Iowa Women's Health Study. *Br J Nutr* 2010;104:1537-43.

Jacobs DR, Jr., Meyer HE, Solvoll K. Reduced mortality among whole grain bread eaters in men and women in the Norwegian County Study. *Eur J Clin Nutr* 2001;55:137-43.

Jacobs DR, Jr., Meyer KA, Kushi LH, Folsom AR. Whole-grain intake may reduce the risk of ischemic heart disease death in postmenopausal women: the Iowa Women's Health Study. *Am J Clin Nutr* 1998;68:248-57.

Jacobs DR, Jr., Meyer KA, Kushi LH, Folsom AR. Is whole grain intake associated with reduced total and cause-specific death rates in older women? The Iowa Women's Health Study. *Am J Public Health* 1999;89:322-9.

Jacobs DR, Pereira MA, Meyer KA, Kushi LH. Fiber from whole grains, but not refined grains, is inversely associated with all-cause mortality in older women: the Iowa women's health study. *J Am Coll Nutr* 2000;19:326S-330S.

Jacobsen BK, Knutsen SF, Fraser GE. Does high soy milk intake reduce prostate cancer incidence? The Adventist Health Study (United States). *Cancer Causes Control* 1998;9:553-7.

Janicki-Deverts D, Cohen S, Matthews KA, Gross MD, Jacobs DR, Jr. Socioeconomic status, antioxidant micronutrients, and correlates of oxidative damage: the Coronary Artery Risk Development in Young Adults (CARDIA) study. *Psychosom Med*. 2009;71:541-8.

Jansen MC, Bueno-de-Mesquita HB, Rasanen L, Fidanza F, Menotti A, Nissinen A, et al. Consumption of plant foods and stomach cancer mortality in the seven countries study. Is grain consumption a risk factor? Seven Countries Study Research Group. *Nutr Cancer* 1999;34:49-55.

Jansen MC, Bueno-de-Mesquita HB, Rasanen L, Fidanza F, Nissinen AM, Menotti A, et al. Cohort analysis of fruit and vegetable consumption and lung cancer mortality in European men. *Int J Cancer* 2001;92:913-8.

Janszky I, Mukamal KJ, Ljung R, Ahnve S, Ahlbom A, Hallqvist J. Chocolate consumption and mortality following a first acute myocardial infarction: the Stockholm Heart Epidemiology Program. *J Intern Med* 2009;266:248-57.

Jarvandi S, Gougeon R, Bader A, Dasgupta K. Differences in food intake among obese and nonobese women and men with type 2 diabetes. *J Am Coll Nutr* 2011;30:225-32.

Jarvinen R, Knekt P, Rissanen H, Reunanen A. Intake of fish and long-chain n-3 fatty acids and the risk of coronary heart mortality in men and women. *Br J Nutr* 2006;95:824-9.

Jeffery RW, Hellerstedt WL, French SA, Baxter JE. A randomized trial of counseling for fat restriction versus calorie restriction in the treatment of obesity. *Int J Obes Relat Metab Disord* 1995;19:132-7.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Jenab M, Ferrari P, Slimani N, Norat T, Casagrande C, Overvad K, et al. Association of nut and seed intake with colorectal cancer risk in the European Prospective Investigation into Cancer and Nutrition. *Cancer Epidemiol Biomarkers Prev* 2004;13:1595-603.

Jenkins DJ, Jenkins AL, Rao AV, Thompson LU. Cancer risk: possible protective role of high carbohydrate high fiber diets. *Am J Gastroenterol* 1986;81:931-5.

Jenkins DJ, Srichaikul K, Kendall CW, Sievenpiper JL, Abdulnour S, Mirrahimi A, et al. The relation of low glycaemic index fruit consumption to glycaemic control and risk factors for coronary heart disease in type 2 diabetes. *Diabetologia* 2011;54:271-9.

Jenkins DJA, Kendall CWC, McKeown-Eyssen G, Josse RG, Silverberg J, Booth GL, et al. Effect of a low-glycemic index or a high-cereal fiber diet on type 2 diabetes: A randomized trial. *JAMA - Journal of the American Medical Association* 2008;300:2742-2753.

Jenkins DJA, Kendall CWC, Nguyen TH, Teitel J, Marchie A, Chiu M, et al. Effect on hematologic risk factors for coronary heart disease of a cholesterol reducing diet. *European Journal of Clinical Nutrition* 2007;61:483-492.

Jensen JD, Wing GJ, Dellavalle RP. Nutrition and melanoma prevention. *Clin Dermatol* 2010;28:644-9.

Jiao L, Flood A, Subar AF, Hollenbeck AR, Schatzkin A, Stolzenberg-Solomon R. Glycemic index, carbohydrates, glycemic load, and the risk of pancreatic cancer in a prospective cohort study. *Cancer Epidemiol Biomarkers Prev*. 2009;18:1144-51.

Jimenez-Cruz A, Jimenez AB, Pichardo-Osuna A, Chaudry T, Bacardi-Gascon M. Long term effect of Mediterranean diet on weight loss. *Nutr Hosp*. 2009;24:753-4.

Joensen AM, Schmidt EB, Dethlefsen C, Johnsen SP, Tjønneland A, Rasmussen LH, et al. Dietary intake of total marine n-3 polyunsaturated fatty acids, eicosapentaenoic acid, docosahexaenoic acid and docosapentaenoic acid and the risk of acute coronary syndrome - a cohort study. *Br J Nutr* 2010;103:602-7.

Johnsen SP, Overvad K, Stripp C, Tjønneland A, Husted SE, Sorensen HT. Intake of fruit and vegetables and the risk of ischemic stroke in a cohort of Danish men and women. *Am J Clin Nutr* 2003;78:57-64.

Johnson ST, Newton AS, Chopra M, Buckingham J, Huang TTK, Franks PW, et al. In search of quality evidence for lifestyle management and glycemic control in children and adolescents with type 2 diabetes: A systematic review. *SOURCE BMC Pediatrics* (2010) 10 Article Number: 97. Date of Publication: 23 Dec 2010 2010.

Jonas CR, McCullough ML, Teras LR, Walker-Thurmond KA, Thun MJ, Calle EE. Dietary glycemic index, glycemic load, and risk of incident breast cancer in postmenopausal women. *Cancer Epidemiol Biomarkers Prev*. 2003;12:573-7.

Jonsson T, Granfeldt Y, Ahren B, Branell UC, Palsson G, Hansson A, et al. Beneficial effects of a Paleolithic diet on cardiovascular risk factors in type 2 diabetes: A randomized cross-over pilot study. *SOURCE Cardiovascular Diabetology* (2009) 8 Article Number: 35. Date of Publication: 16 Jul 2009 2009.

Joshiyura KJ, Hu FB, Manson JE, Stampfer MJ, Rimm EB, Speizer FE, et al. The effect of fruit and vegetable intake on risk for coronary heart disease. *Ann Intern Med* 2001;134:1106-14.

Kabagambe EK, Baylin A, Campos H. Nonfatal acute myocardial infarction in Costa Rica: modifiable risk factors, population-attributable risks, and adherence to dietary guidelines. *Circulation* 2007;115:1075-81.

Kabat GC, Miller AB, Jain M, Rohan TE. Dietary iron and haem iron intake and risk of endometrial cancer: A prospective cohort study. *British Journal of Cancer* 2008;98:194-198.

Kahn HA, Phillips RL, Snowdon DA, Choi W. Association between reported diet and all-cause mortality. Twenty-one-year follow-up on 27,530 adult Seventh-Day Adventists. *Am J Epidemiol* 1984;119:775-87.

Kaitosaari T, Ronnema T, Raitakari O, Talvia S, Kallio K, Volanen I, et al. Effect of 7-year infancy-onset dietary intervention on serum lipoproteins and lipoprotein subclasses in healthy children in the prospective, randomized Special Turku Coronary Risk Factor Intervention Project for Children (STRIP) study. *Circulation* 2003;108:672-7.

Kalarchian MA, Levine MD, Arslanian SA, Ewing LJ, Houck PR, Cheng Y, et al. Family-based treatment of severe pediatric obesity: Randomized, controlled trial. *Pediatrics* 2009;124:1060-1068.

Kaluza J, Hakansson N, Brzozowska A, Wolk A. Diet quality and mortality: a population-based prospective study of men. *Eur J Clin Nutr* 2009;63:451-7.

Kamphuis MH, Geerlings MI, Tijhuis MA, Kalmijn S, Grobbee DE, Kromhout D. Depression and cardiovascular mortality: a role for n-3 fatty acids? *Am J Clin Nutr* 2006;84:1513-7.

Kampman E, Arts IC, Hollman PC. Plant foods versus compounds in carcinogenesis; observational versus experimental human studies. *Int J Vitam Nutr Res* 2003;73:70-8.

Kanis JA, Johansson H, Oden A, De Laet C, Johnell O, Eisman JA, et al. A meta-analysis of milk intake and fracture risk: low utility for case finding. *Osteoporos Int*. 2005;16:799-804.

Kant AK. Weight-loss attempts and reporting of foods and nutrients, and biomarkers in a national cohort. *Int J Obes Relat Metab Disord* 2002;26:1194-204.

Kant AK, Graubard BI, Schatzkin A. Dietary patterns predict mortality in a national cohort: the National Health Interview Surveys, 1987 and 1992. *J Nutr*. 2004;134:1793-9.

Kant AK, Leitzmann MF, Park Y, Hollenbeck A, Schatzkin A. Patterns of recommended dietary behaviors predict subsequent risk of mortality in a large cohort of men and women in the United States. *J Nutr*. 2009;139:1374-80.

Kant AK, Schatzkin A, Graubard BI, Schairer C. A prospective study of diet quality and mortality in women. *JAMA*. 2000;283:2109-15.

Kant AK, Schatzkin A, Ziegler RG. Dietary diversity and subsequent cause-specific mortality in the NHANES I epidemiologic follow-up study. *J Am Coll Nutr* 1995;14:233-8.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Karagianni V, Merikas E, Georgopoulos F, Gikas A, Athanasopoulos N, Malgarinos G, et al. Risk factors for colorectal polyps: findings from a Greek case-control study. *Rev Med Chir Soc Med Nat Iasi* 2010;114:662-70.

Kasim-Karakas SE, Almario RU, Mueller WM, Peerson J. Changes in plasma lipoproteins during low-fat, high-carbohydrate diets: effects of energy intake. *Am J Clin Nutr*. 2000;71:1439-47.

Kasperzyk JL, Fall K, Mucci LA, Hakansson N, Wolk A, Johansson JE, et al. One-carbon metabolism-related nutrients and prostate cancer survival. *Am J Clin Nutr* 2009;90:561-9.

Kastorini CM, Milionis HJ, Esposito K, Giugliano D, Goudevenos JA, Panagiotakos DB. The effect of mediterranean diet on metabolic syndrome and its components: A meta-analysis of 50 studies and 534,906 individuals. *Journal of the American College of Cardiology* 2011;57:1299-1313.

Kasum CM, Nicodemus K, Harnack LJ, Jacobs DR, Jr., Folsom AR. Whole grain intake and incident endometrial cancer: the Iowa Women's Health Study. *Nutr Cancer*. 2001;39:180-6.

Kaushik S, Wang JJ, Wong TY, Flood V, Barclay A, Brand-Miller J, et al. Glycemic index, retinal vascular caliber, and stroke mortality. *Stroke* 2009;40:206-12.

Kawashima H, Takase H, Yasunaga K, Wakaki Y, Katsuragi Y, Mori K, et al. One-year ad libitum consumption of diacylglycerol oil as part of a regular diet results in modest weight loss in comparison with consumption of a triacylglycerol control oil in overweight Japanese subjects. *J Am Diet Assoc* 2008;108:57-66.

Keim NL, Van Loan MD, Horn WF, Barbieri TF, Mayclin PL. Weight loss is greater with consumption of large morning meals and fat-free mass is preserved with large evening meals in women on a controlled weight reduction regimen. *Journal of Nutrition* 1997;127:75-82.

Kelemen LE, Kushi LH, Jacobs DR, Jr., Cerhan JR. Associations of dietary protein with disease and mortality in a prospective study of postmenopausal women. *Am J Epidemiol* 2005;161:239-49.

Keli SO, Hertog MG, Feskens EJ, Kromhout D. Dietary flavonoids, antioxidant vitamins, and incidence of stroke: the Zutphen study. *Arch Intern Med* 1996;156:637-42.

Kelishadi R, Zemel MB, Hashemipour M, Hosseini M, Mohammadifard N, Poursafa P. Can a dairy-rich diet be effective in long-term weight control of young children? *J Am Coll Nutr* 2009;28:601-10.

Kelly JH, Jr., Sabate J. Nuts and coronary heart disease: an epidemiological perspective. *Br J Nutr* 2006;96 Suppl 2:S61-7.

Keränen A-M, Savolainen MJ, Reponen AH, Kujari M-L, Lindeman SM, Bloigu RS, et al. The effect of eating behavior on weight loss and maintenance during a lifestyle intervention. *Preventive Medicine: An International Journal Devoted to Practice and Theory* 2009;49:32-38.

Keszei AP, Verhage BA, Heinen MM, Goldbohm RA, van den Brandt PA. Dietary folate and folate vitamers and the risk of pancreatic cancer in the Netherlands cohort study. *Cancer Epidemiol Biomarkers Prev*. 2009;18:1785-91.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Key TJ, Allen N, Appleby P, Overvad K, Tjonneland A, Miller A, et al. Fruits and vegetables and prostate cancer: no association among 1104 cases in a prospective study of 130544 men in the European Prospective Investigation into Cancer and Nutrition (EPIC). *Int J Cancer* 2004;109:119-24.

Key TJ, Appleby PN, Davey GK, Allen NE, Spencer EA, Travis RC. Mortality in British vegetarians: review and preliminary results from EPIC-Oxford. *Am J Clin Nutr* 2003;78:533S-538S.

Key TJ, Appleby PN, Spencer EA, Roddam AW, Neale RE, Allen NE. Calcium, diet and fracture risk: a prospective study of 1898 incident fractures among 34 696 British women and men. *Public Health Nutr* 2007;10:1314-20.

Key TJ, Appleby PN, Spencer EA, Travis RC, Roddam AW, Allen NE. Mortality in British vegetarians: results from the European Prospective Investigation into Cancer and Nutrition (EPIC-Oxford). *Am J Clin Nutr* 2009;89:1613S-1619S.

Key TJ, Fraser GE, Thorogood M, Appleby PN, Beral V, Reeves G, et al. Mortality in vegetarians and non-vegetarians: a collaborative analysis of 8300 deaths among 76,000 men and women in five prospective studies. *Public Health Nutr* 1998;1:33-41.

Key TJ, Fraser GE, Thorogood M, Appleby PN, Beral V, Reeves G, et al. Mortality in vegetarians and nonvegetarians: detailed findings from a collaborative analysis of 5 prospective studies. *Am J Clin Nutr* 1999;70:516S-524S.

Key TJ, Thorogood M, Appleby PN, Burr ML. Dietary habits and mortality in 11,000 vegetarians and health conscious people: results of a 17 year follow up. *BMJ* 1996;313:775-9.

Khan MM, Goto R, Kobayashi K, Suzumura S, Nagata Y, Sonoda T, et al. Dietary habits and cancer mortality among middle aged and older Japanese living in hokkaido, Japan by cancer site and sex. *Asian Pac J Cancer Prev* 2004;5:58-65.

Khaw KT, Barrett-Connor E. Dietary fiber and reduced ischemic heart disease mortality rates in men and women: a 12-year prospective study. *Am J Epidemiol* 1987;126:1093-102.

Kim JL, Winkvist A, Aberg MA, Aberg N, Sundberg R, Toren K, et al. Fish consumption and school grades in Swedish adolescents: a study of the large general population. *Acta Paediatr* 2010;99:72-7.

Kinjo Y, Beral V, Akiba S, Key T, Mizuno S, Appleby P, et al. Possible protective effect of milk, meat and fish for cerebrovascular disease mortality in Japan. *J Epidemiol* 1999;9:268-74.

Kinlen LJ, Hermon C, Smith PG. A proportionate study of cancer mortality among members of a vegetarian society. *Br J Cancer* 1983;48:355-61.

Kirii K, Iso H, Date C, Fukui M, Tamakoshi A. Magnesium intake and risk of self-reported type 2 diabetes among Japanese. *J Am Coll Nutr* 2010;29:99-106.

Kirsh VA, Mayne ST, Peters U, Chatterjee N, Leitzmann MF, Dixon LB, et al. A prospective study of lycopene and tomato product intake and risk of prostate cancer. *Cancer Epidemiol Biomarkers Prev* 2006;15:92-8.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Kitchen MS, Ransley JK, Greenwood DC, Clarke GP, Conner MT, Jupp J, et al. Study protocol: A cluster randomised controlled trial of a school based fruit and vegetable intervention Project Tomato. *SOURCE BMC Health Services Research* (2009) 9 Article Number: 101. Date of Publication: 2009 2009.

Kiyohara Y, Shinohara A, Kato I, Shiota T, Kubo M, Tanizaki Y, et al. Dietary factors and development of impaired glucose tolerance and diabetes in a general Japanese population: the hisayama study. *J Epidemiol* 2003;13:251-8.

Kjeldsen-Kragh J, Haugen M, Borchgrevink CF, Forre O. Vegetarian diet for patients with rheumatoid arthritis--status: two years after introduction of the diet. *Clin Rheumatol*. 1994;13:475-82.

Klassen AC, Garrett-Mayer E, Houts PS, Shankar S, Torio CM. The relationship of body size to participation and success in a fruits and vegetables intervention among low-income women. *Journal of Community Health* 2008;33:78-89.

Kleber M, Schaefer A, Winkel K, Hoffmann D, Wunsch R, Kersting M, et al. Lifestyle intervention "Obeldicks Mini" for obese children aged 4 to 7 years. *Klin Padiatr* 2009;221:290-4.

Klein S. Clinical Trial Experience with Fat-Restricted vs. Carbohydrate-Restricted Weight-Loss Diets. *Obesity Research* 2004;12:141S-144S.

Klesges RC, Isbell TR, Klesges LM. Relationship between dietary restraint, energy intake, physical activity, and body weight: a prospective analysis. *J Abnorm Psychol*. 1992;101:668-74.

Klesges RC, Klesges LM, Haddock CK, Eck LH. A longitudinal analysis of the impact of dietary intake and physical activity on weight change in adults. *Am J Clin Nutr* 1992;55:818-22.

Klipstein-Grobusch K, Geleijnse JM, den Breeijen JH, Boeing H, Hofman A, Grobbee DE, et al. Dietary antioxidants and risk of myocardial infarction in the elderly: the Rotterdam Study. *Am J Clin Nutr* 1999;69:261-6.

Knekt P, Jarvinen R, Reunanen A, Maatela J. Flavonoid intake and coronary mortality in Finland: a cohort study. *BMJ* 1996;312:478-81.

Knoops KT, de Groot LC, Kromhout D, Perrin AE, Moreiras-Varela O, Menotti A, et al. Mediterranean diet, lifestyle factors, and 10-year mortality in elderly European men and women: the HALE project. *JAMA* 2004;292:1433-9.

Knoops KT, Groot de LC, Fidanza F, Alberti-Fidanza A, Kromhout D, van Staveren WA. Comparison of three different dietary scores in relation to 10-year mortality in elderly European subjects: the HALE project. *Eur J Clin Nutr*. 2006;60:746-55.

Knowlden AP, Sharma M. Systematic review of family and home-based interventions targeting paediatric overweight and obesity. *Obes Rev* 2012;13:499-508.

Knowles HC, Jr. Long-term juvenile diabetes treated with unmeasured diet. *Trans Assoc Am Physicians* 1971;84:95-101.

Kochar J, Gaziano JM, Djousse L. Nut consumption and risk of type II diabetes in the Physicians' Health Study. *Eur J Clin Nutr* 2010;64:75-9.

Kochar J, Gaziano JM, Djousse L. Breakfast cereals and risk of hypertension in the Physicians' Health Study I. *Clinical Nutrition* 2012;31:89-92.

Koh WP, Wu AH, Wang R, Ang LW, Heng D, Yuan JM, et al. Gender-specific associations between soy and risk of hip fracture in the Singapore Chinese Health Study. *Am J Epidemiol* 2009;170:901-9.

Koh-Banerjee P, Franz M, Sampson L, Liu S, Jacobs DR, Jr., Spiegelman D, et al. Changes in whole-grain, bran, and cereal fiber consumption in relation to 8-y weight gain among men. *Am J Clin Nutr*. 2004;80:1237-45.

Kojima M, Wakai K, Tamakoshi K, Tokudome S, Toyoshima H, Watanabe Y, et al. Diet and colorectal cancer mortality: results from the Japan Collaborative Cohort Study. *Nutr Cancer* 2004;50:23-32.

Kolonel LN, Yoshizawa CN, Hankin JH. Diet and prostatic cancer: a case-control study in Hawaii. *Am J Epidemiol*. 1988;127:999-1012.

Konings EJ, Goldbohm RA, Brants HA, Saris WH, van den Brandt PA. Intake of dietary folate vitamers and risk of colorectal carcinoma: results from The Netherlands Cohort Study. *Cancer* 2002;95:1421-33.

Koppes LL, Boon N, Nooyens AC, van Mechelen W, Saris WH. Macronutrient distribution over a period of 23 years in relation to energy intake and body fatness. *Br J Nutr*. 2009;101:108-15.

Koppo K, Siklova-Vitkova M, Klimcakova E, Polak J, Marques MA, Berlan M, et al. Catecholamine and insulin control of lipolysis in subcutaneous adipose tissue during long-term diet-induced weight loss in obese women. *Am J Physiol Endocrinol Metab* 2012;302:E226-32.

Kossoff EH, Rowley H, Sinha SR, Vining EP. A prospective study of the modified Atkins diet for intractable epilepsy in adults. *Epilepsia*. 2008;49:316-9.

Kostis JB, Wilson AC, Shindler DM, Cosgrove NM, Lacy CR. Persistence of normotension after discontinuation of lifestyle intervention in the trial of TONE. *Trial of Nonpharmacologic Interventions in the Elderly*. *Am J Hypertens* 2002;15:732-4.

Kouris-Blazos A, Gnardellis C, Wahlqvist ML, Trichopoulos D, Lukito W, Trichopoulou A. Are the advantages of the Mediterranean diet transferable to other populations? A cohort study in Melbourne, Australia. *Br J Nutr* 1999;82:57-61.

Koushik A, Hunter DJ, Spiegelman D, Beeson WL, van den Brandt PA, Buring JE, et al. Fruits, vegetables, and colon cancer risk in a pooled analysis of 14 cohort studies. *J Natl Cancer Inst* 2007;99:1471-83.

Kratz M, Baars T, Guyenet S. The relationship between high-fat dairy consumption and obesity, cardiovascular, and metabolic disease. *Eur J Nutr* 2012.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Krebs JD, Browning LM, McLean NK, Rothwell JL, Mishra GD, Moore CS, et al. Additive benefits of long-chain n-3 polyunsaturated fatty acids and weight-loss in the management of cardiovascular disease risk in overweight hyperinsulinaemic women. *Int J Obes (Lond)* 2006;30:1535-44.

Kring SI, Heitmann BL. Fiber intake, not dietary energy density, is associated with subsequent change in BMI z-score among sub-groups of children. *Obes Facts* 2008;1:331-8.

Krishnan S, Rosenberg L, Singer M, Hu FB, Djousse L, Cupples LA, et al. Glycemic index, glycemic load, and cereal fiber intake and risk of type 2 diabetes in US black women. *Arch Intern Med* 2007;167:2304-9.

Krishnaswami J, Martinson M, Wakimoto P, Anglemeyer A. Community-engaged interventions on diet, activity, and weight outcomes in U.S. schools: a systematic review. *Am J Prev Med* 2012;43:81-91.

Kristal AR, White E, Shattuck AL, Curry S, Anderson GL, Fowler A, et al. Long-term maintenance of a low-fat diet: durability of fat-related dietary habits in the Women's Health Trial. *J Am Diet Assoc* 1992;92:553-9.

Kritchevsky SB. Dietary lipids and the low blood cholesterol-cancer association. *American Journal of Epidemiology* 1992;135:509-520.

Kritchevsky SB, Tell GS, Shimakawa T, Dennis B, Li R, Kohlmeier L, et al. Provitamin A carotenoid intake and carotid artery plaques: the Atherosclerosis Risk in Communities Study. *Am J Clin Nutr* 1998;68:726-33.

Kromhout D. Body weight, diet, and serum cholesterol in 871 middle-aged men during 10 years of follow-up (the Zutphen Study). *Am J Clin Nutr* 1983;38:591-8.

Kromhout D, Bloemberg B, Feskens E, Menotti A, Nissinen A. Saturated fat, vitamin C and smoking predict long-term population all-cause mortality rates in the Seven Countries Study. *Int J Epidemiol* 2000;29:260-5.

Kromhout D, Bosschieter EB, de Lezenne Coulander C. Dietary fibre and 10-year mortality from coronary heart disease, cancer, and all causes. The Zutphen study. *Lancet* 1982;2:518-22.

Kromhout D, Bosschieter EB, de Lezenne Coulander C. The inverse relation between fish consumption and 20-year mortality from coronary heart disease. *N Engl J Med* 1985;312:1205-9.

Kromhout D, de Lezenne Coulander C. Diet, prevalence and 10-year mortality from coronary heart disease in 871 middle-aged men. The Zutphen Study. *Am J Epidemiol.* 1984;119:733-41.

Kromhout D, Giltay EJ, Geleijnse JM. n-3 fatty acids and cardiovascular events after myocardial infarction. *N Engl J Med* 2010;363:2015-26.

Kromhout D, Menotti A, Bloemberg B, Aravanis C, Blackburn H, Buzina R, et al. Dietary saturated and trans fatty acids and cholesterol and 25-year mortality from coronary heart disease: the Seven Countries Study. *Prev Med* 1995;24:308-15.

Kromhout D, Saris WH, Horst CH. Energy intake, energy expenditure, and smoking in relation to body fatness: the Zutphen Study. *Am J Clin Nutr.* 1988;47:668-74.

Kuczmarski MF, Cremer Sees A, Hotchkiss L, Cotugna N, Evans MK, Zonderman AB. Higher Healthy Eating Index-2005 scores associated with reduced symptoms of depression in an urban population: findings from the Healthy Aging in Neighborhoods of Diversity Across the Life Span (HANDLS) study. *J Am Diet Assoc* 2010;110:383-9.

Kumagai S, Shibata H, Watanabe S, Suzuki T, Haga H. Effect of food intake pattern on all-cause mortality in the community elderly: a 7-year longitudinal study. *J Nutr Health Aging* 1999;3:29-33.

Kumanyika SK, Cook NR, Cutler JA, Belden L, Brewer A, Cohen JD, et al. Sodium reduction for hypertension prevention in overweight adults: further results from the Trials of Hypertension Prevention Phase II. *J Hum Hypertens* 2005;19:33-45.

Kurahashi N, Inoue M, Iwasaki M, Tanaka Y, Mizokami M, Tsugane S. Vegetable, fruit and antioxidant nutrient consumption and subsequent risk of hepatocellular carcinoma: a prospective cohort study in Japan. *Br J Cancer* 2009;100:181-4.

Kurosawa M, Kikuchi S, Xu J, Inaba Y. Highly salted food and mountain herbs elevate the risk for stomach cancer death in a rural area of Japan. *J Gastroenterol Hepatol* 2006;21:1681-6.

Kurumatani N, Kirita T, Zheng Y, Sugimura M, Yonemasu K. Time trends in the mortality rates for tobacco- and alcohol-related cancers within the oral cavity and pharynx in Japan, 1950-94. *J Epidemiol* 1999;9:46-52.

Kushi LH, Folsom AR, Prineas RJ, Mink PJ, Wu Y, Bostick RM. Dietary antioxidant vitamins and death from coronary heart disease in postmenopausal women. *N Engl J Med* 1996;334:1156-62.

Kushi LH, Lew RA, Stare FJ, Ellison CR, el Lozy M, Bourke G, et al. Diet and 20-year mortality from coronary heart disease. The Ireland-Boston Diet-Heart Study. *N Engl J Med* 1985;312:811-8.

Kwiterovich PO, Jr., Barton BA, McMahan RP, Obarzanek E, Hunsberger S, Simons-Morton D, et al. Effects of diet and sexual maturation on low-density lipoprotein cholesterol during puberty: the Dietary Intervention Study in Children (DISC). *Circulation*. 1997;96:2526-33.

Kyrozis A, Psaltopoulou T, Stathopoulos P, Trichopoulos D, Vassilopoulos D, Trichopoulou A. Dietary lipids and geriatric depression scale score among elders: the EPIC-Greece cohort. *J Psychiatr Res* 2009;43:763-9.

Laaksonen DE, Nyyssonen K, Niskanen L, Rissanen TH, Salonen JT. Prediction of cardiovascular mortality in middle-aged men by dietary and serum linoleic and polyunsaturated fatty acids. *Arch Intern Med* 2005;165:193-9.

Lackland DT, Egan BM. Dietary salt restriction and blood pressure in clinical trials. *Current Hypertension Reports* 2007;9:314-319.

Lagiou P, Sandin S, Weiderpass E, Lagiou A, Mucci L, Trichopoulos D, et al. Low carbohydrate-high protein diet and mortality in a cohort of Swedish women. *J Intern Med* 2007;261:366-74.

Lagiou P, Trichopoulos D, Sandin S, Lagiou A, Mucci L, Wolk A, et al. Mediterranean dietary pattern and mortality among young women: a cohort study in Sweden. *Br J Nutr* 2006;96:384-92.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Lajous M, Boutron-Ruault MC, Fabre A, Clavel-Chapelon F, Romieu I. Carbohydrate intake, glycemic index, glycemic load, and risk of postmenopausal breast cancer in a prospective study of French women. *Am J Clin Nutr* 2008;87:1384-91.

Lajous M, Tondeur L, Fagherazzi G, de Lauzon-Guillain B, Boutron-Ruault MC, Clavel-Chapelon F. Processed and unprocessed red meat consumption and incident type 2 diabetes among French women. *Diabetes Care* 2012;35:128-30.

Langsetmo L, Hanley DA, Prior JC, Barr SI, Anastassiades T, Towheed T, et al. Dietary patterns and incident low-trauma fractures in postmenopausal women and men aged \geq 50 y: a population-based cohort study. *Am J Clin Nutr* 2011;93:192-9.

Lanza E, Schatzkin A, Daston C, Corle D, Freedman L, Ballard-Barbash R, et al. Implementation of a 4-y, high-fiber, high-fruit-and-vegetable, low-fat dietary intervention: results of dietary changes in the Polyp Prevention Trial. *Am J Clin Nutr* 2001;74:387-401.

Lanza E, Yu B, Murphy G, Albert PS, Caan B, Marshall JR, et al. The polyp prevention trial continued follow-up study: no effect of a low-fat, high-fiber, high-fruit, and -vegetable diet on adenoma recurrence eight years after randomization. *Cancer Epidemiol Biomarkers Prev* 2007;16:1745-52.

Lapidus L, Andersson H, Bengtsson C, Bosaeus I. Dietary habits in relation to incidence of cardiovascular disease and death in women: a 12-year follow-up of participants in the population study of women in Gothenburg, Sweden. *Am J Clin Nutr* 1986;44:444-8.

Lapointe A, Weisnagel SJ, Provencher V, Begin C, Dufour-Bouchard AA, Trudeau C, et al. Using restrictive messages to limit high-fat foods or nonrestrictive messages to increase fruit and vegetable intake: what works better for postmenopausal women? *Eur J Clin Nutr* 2009;64:194-202.

Lappe JM, Rafferty KA, Davies KM, Lypaczewski G. Girls on a high-calcium diet gain weight at the same rate as girls on a normal diet: a pilot study. *J Am Diet Assoc* 2004;104:1361-7.

Larson-Meyer DE, Newcomer BR, Heilbronn LK, Volaufova J, Smith SR, Alfonso AJ, et al. Effect of 6-month calorie restriction and exercise on serum and liver lipids and markers of liver function. *Obesity (Silver Spring)* 2008;16:1355-62.

Larsson SC, Bergkvist L, Rutegard J, Giovannucci E, Wolk A. Calcium and dairy food intakes are inversely associated with colorectal cancer risk in the Cohort of Swedish Men. *Am J Clin Nutr* 2006;83:667-73; quiz 728-9.

Larsson SC, Bergkvist L, Wolk A. Milk and lactose intakes and ovarian cancer risk in the Swedish Mammography Cohort. *Am J Clin Nutr*. 2004;80:1353-7.

Larsson SC, Bergkvist L, Wolk A. Fruit and vegetable consumption and incidence of gastric cancer: a prospective study. *Cancer Epidemiol Biomarkers Prev* 2006;15:1998-2001.

Larsson SC, Bergkvist L, Wolk A. Glycemic load, glycemic index and breast cancer risk in a prospective cohort of Swedish women. *Int J Cancer*. 2009;125:153-7.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Larsson SC, Friberg E, Wolk A. Carbohydrate intake, glycemic index and glycemic load in relation to risk of endometrial cancer: A prospective study of Swedish women. *Int J Cancer* 2007;120:1103-7.

Larsson SC, Holmberg L, Wolk A. Fruit and vegetable consumption in relation to ovarian cancer incidence: the Swedish Mammography Cohort. *Br J Cancer* 2004;90:2167-70.

Larsson SC, Virtamo J, Wolk A. Fish consumption and risk of stroke in Swedish women. *Am J Clin Nutr*. 2011;93:487-93.

Larsson SC, Virtamo J, Wolk A. Chocolate consumption and risk of stroke in women. *J Am Coll Cardiol* 2011;58:1828-9.

Lasker DAW, Evans EM, Layman DK. Moderate carbohydrate, moderate protein weight loss diet reduces cardiovascular disease risk compared to high carbohydrate, low protein diet in obese adults: A randomized clinical trial. *SOURCE Nutrition and Metabolism* (2008) 5:1 Article Number: 30. Date of Publication: 2008 2008.

Law MR, Morris JK. By how much does fruit and vegetable consumption reduce the risk of ischaemic heart disease? *Eur J Clin Nutr* 1998;52:549-56.

Le Marchand L, Kolonel LN, Wilkens LR, Myers BC, Hirohata T. Animal fat consumption and prostate cancer: A prospective study in Hawaii. *Epidemiology* 1994;5:276-282.

Lean MEJ, Han TS, Prvan T, Richmond PR, Avenell A. Weight loss with high and low carbohydrate 1200 kcal diets in free living women. *European Journal of Clinical Nutrition* 1997;51:243-248.

Lee JE, Giovannucci E, Smith-Warner SA, Spiegelman D, Willett WC, Curhan GC. Intakes of fruits, vegetables, vitamins A, C, and E, and carotenoids and risk of renal cell cancer. *Cancer Epidemiol Biomarkers Prev*. 2006;15:2445-52.

Lee JE, Spiegelman D, Hunter DJ, Albanes D, Bernstein L, van den Brandt PA, et al. Fat, protein, and meat consumption and renal cell cancer risk: a pooled analysis of 13 prospective studies. *J Natl Cancer Inst*. 2008;100:1695-706.

Lee SA, Shu XO, Li H, Yang G, Cai H, Wen W, et al. Adolescent and adult soy food intake and breast cancer risk: results from the Shanghai Women's Health Study. *Am J Clin Nutr* 2009;89:1920-6.

Legrand P, Schmitt B, Mourot J, Catheline D, Chesneau G, Mireaux M, et al. The consumption of food products from linseed-fed animals maintains erythrocyte omega-3 fatty acids in obese humans. *Lipids* 2010;45:11-9.

Lehtonen HM, Suomela JP, Tahvonen R, Vaarno J, Venojärvi M, Viikari J, et al. Berry meals and risk factors associated with metabolic syndrome. *Eur J Clin Nutr* 2010;64:614-21.

Lemaitre RN, King IB, Mozaffarian D, Kuller LH, Tracy RP, Siscovick DS. n-3 Polyunsaturated fatty acids, fatal ischemic heart disease, and nonfatal myocardial infarction in older adults: the Cardiovascular Health Study. *Am J Clin Nutr* 2003;77:319-25.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Lengyel CO, Tate RB, Obirek Blatz AK. The relationships between food group consumption, self-rated health, and life satisfaction of community-dwelling canadian older men: the manitoba follow-up study. *J Nutr Elder* 2009;28:158-73.

Leosdottir M, Nilsson P, Nilsson JA, Mansson H, Berglund G. The association between total energy intake and early mortality: data from the Malmo Diet and Cancer Study. *J Intern Med* 2004;256:499-509.

Leosdottir M, Nilsson PM, Nilsson JA, Mansson H, Berglund G. Dietary fat intake and early mortality patterns--data from The Malmo Diet and Cancer Study. *J Intern Med* 2005;258:153-65.

Leren P. The effect of plasma cholesterol lowering diet in male survivors of myocardial infarction. A controlled clinical trial. *Acta Med Scand Suppl* 1966;466:1-92.

Leren P. The Oslo diet-heart study. Eleven-year report. *Circulation* 1970;42:935-42.

Leren P. Prevention of coronary heart disease by diet. *Postgrad Med J* 1975;51:suppl 44-6.

Leren P. Prevention of coronary heart disease: some results from the Oslo secondary and primary intervention studies. *J Am Coll Nutr* 1989;8:407-10.

Levitan EB, Mittleman MA, Hakansson N, Wolk A. Dietary glycemic index, dietary glycemic load, and cardiovascular disease in middle-aged and older Swedish men. *Am J Clin Nutr* 2007;85:1521-6.

Levitan EB, Mittleman MA, Wolk A. Dietary glycaemic index, dietary glycaemic load and incidence of myocardial infarction in women. *Br J Nutr* 2010;103:1049-55.

Levitan EB, Mittleman MA, Wolk A. Dietary glycemic index, dietary glycemic load, and incidence of heart failure events: a prospective study of middle-aged and elderly women. *J Am Coll Nutr* 2010;29:65-71.

Levitan EB, Wolk A, Mittleman MA. Relation of consistency with the dietary approaches to stop hypertension diet and incidence of heart failure in men aged 45 to 79 years. *Am J Cardiol* 2009;104:1416-20.

Levitan EB, Wolk A, Mittleman MA. Consistency with the DASH diet and incidence of heart failure. *Arch Intern Med* 2009;169:851-7.

Levitan EB, Wolk A, Mittleman MA. Fish consumption, marine omega-3 fatty acids, and incidence of heart failure: a population-based prospective study of middle-aged and elderly men. *Eur Heart J* 2009;30:1495-500.

Ley SJ, Metcalf PA, Scragg RKR, Swinburn BA. Long-term effects of a reduced fat diet intervention on cardiovascular disease risk factors in individuals with glucose intolerance. *Diabetes Research and Clinical Practice* 2004;63:103-112.

Li Y, Dai Q, Tedders SH, Arroyo C, Zhang J. Legume consumption and severe depressed mood, the modifying roles of gender and menopausal status. *Public Health Nutr* 2010;13:1198-206.

SBU. Mat vid fetma. En systematisk litteraturöversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Lien do TK, Nhung BT, Khan NC, Hop le T, Nga NT, Hung NT, et al. Impact of milk consumption on performance and health of primary school children in rural Vietnam. *Asia Pac J Clin Nutr*. 2009;18:326-34.

Lien LF, Brown AJ, Ard JD, Loria C, Erlinger TP, Feldstein AC, et al. Effects of PREMIER lifestyle modifications on participants with and without the metabolic syndrome. *Hypertension* 2007;50:609-16.

Liese AD, Weis KE, Schulz M, Toozee JA. Food intake patterns associated with incident type 2 diabetes: the Insulin Resistance Atherosclerosis Study. *Diabetes Care* 2009;32:263-8.

Lin J, Rexrode KM, Hu F, Albert CM, Chae CU, Rimm EB, et al. Dietary intakes of flavonols and flavones and coronary heart disease in US women. *Am J Epidemiol* 2007;165:1305-13.

Lin J, Zhang SM, Cook NR, Rexrode KM, Liu S, Manson JE, et al. Dietary intakes of fruit, vegetables, and fiber, and risk of colorectal cancer in a prospective cohort of women (United States). *Cancer Causes Control* 2005;16:225-33.

Lindstrom J, Ilanne-Parikka P, Peltonen M, Aunola S, Eriksson JG, Hemio K, et al. Sustained reduction in the incidence of type 2 diabetes by lifestyle intervention: follow-up of the Finnish Diabetes Prevention Study. *Lancet* 2006;368:1673-9.

Lindstrom J, Peltonen M, Eriksson JG, Louheranta A, Fogelholm M, Uusitupa M, et al. High-fibre, low-fat diet predicts long-term weight loss and decreased type 2 diabetes risk: the Finnish Diabetes Prevention Study. *Diabetologia* 2006;49:912-20.

Linos E, Holmes MD, Willett WC. Diet and breast cancer. *Current Oncology Reports* 2007;9:31-41.

Little P, Kelly J, Barnett J, Dorward M, Margetts B, Warm D. Randomised controlled factorial trial of dietary advice for patients with a single high blood pressure reading in primary care. *British Medical Journal* 2004;328:1054-1057.

Liu S, Lee IM, Ajani U, Cole SR, Buring JE, Manson JE. Intake of vegetables rich in carotenoids and risk of coronary heart disease in men: The Physicians' Health Study. *Int J Epidemiol*. 2001;30:130-5.

Liu S, Manson JE, Lee IM, Cole SR, Hennekens CH, Willett WC, et al. Fruit and vegetable intake and risk of cardiovascular disease: the Women's Health Study. *Am J Clin Nutr* 2000;72:922-8.

Liu S, Manson JE, Stampfer MJ, Holmes MD, Hu FB, Hankinson SE, et al. Dietary glycemic load assessed by food-frequency questionnaire in relation to plasma high-density-lipoprotein cholesterol and fasting plasma triacylglycerols in postmenopausal women. *Am J Clin Nutr*. 2001;73:560-6.

Liu S, Manson JE, Stampfer MJ, Hu FB, Giovannucci E, Colditz GA, et al. A prospective study of whole-grain intake and risk of type 2 diabetes mellitus in US women. *Am J Public Health* 2000;90:1409-15.

Liu S, Serdula M, Janket SJ, Cook NR, Sesso HD, Willett WC, et al. A prospective study of fruit and vegetable intake and the risk of type 2 diabetes in women. *Diabetes Care* 2004;27:2993-6.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Liu S, Sesso HD, Manson JE, Willett WC, Buring JE. Is intake of breakfast cereals related to total and cause-specific mortality in men? *Am J Clin Nutr* 2003;77:594-9.

Liu S, Stampfer MJ, Hu FB, Giovannucci E, Rimm E, Manson JE, et al. Whole-grain consumption and risk of coronary heart disease: results from the Nurses' Health Study. *Am J Clin Nutr* 1999;70:412-9.

Liu S, Willett WC, Manson JE, Hu FB, Rosner B, Colditz G. Relation between changes in intakes of dietary fiber and grain products and changes in weight and development of obesity among middle-aged women. *Am J Clin Nutr* 2003;78:920-7.

Liu S, Willett WC, Stampfer MJ, Hu FB, Franz M, Sampson L, et al. A prospective study of dietary glycemic load, carbohydrate intake, and risk of coronary heart disease in US women. *Am J Clin Nutr*. 2000;71:1455-61.

Liu Y, Sobue T, Otani T, Tsugane S. Vegetables, fruit consumption and risk of lung cancer among middle-aged Japanese men and women: JPHC study. *Cancer Causes Control* 2004;15:349-57.

Liu ZM, Ho SC, Chen YM, Ho YP. A mild favorable effect of soy protein with isoflavones on body composition: a 6-month double-blind randomized placebo-controlled trial among Chinese postmenopausal women. *International Journal of Obesity* 2010;34:309-318.

Llaneza P, Gonzalez C, Fernandez-Inarrea J, Alonso A, Diaz F, Arnott I, et al. Soy isoflavones, diet and physical exercise modify serum cytokines in healthy obese postmenopausal women. *Phytomedicine* 2011;18:245-250.

Logan KJ, Woodside JV, Young IS, McKinley MC, Perkins-Porras L, McKeown PP. Adoption and maintenance of a Mediterranean diet in patients with coronary heart disease from a Northern European population: a pilot randomised trial of different methods of delivering Mediterranean diet advice. *Journal of Human Nutrition & Dietetics* 2010;23:30-37.

London S, Willett W. Diet and the risk of breast cancer. *Hematol Oncol Clin North Am*. 1989;3:559-76.

Lopez CN, Martinez-Gonzalez MA, Sanchez-Villegas A, Alonso A, Pimenta AM, Bes-Rastrollo M. Costs of Mediterranean and western dietary patterns in a Spanish cohort and their relationship with prospective weight change. *Journal of Epidemiology and Community Health* 2009;63:920-927.

Lopez Plaza B, Bermejo LM, Zurita Rosa L, Rodriguez Duran D, Palma Milla S, Weber TK, et al. Regular consumption of functional oil enriched with omega 3 fatty acids from fish oil improves the blood lipid profile in obese people. *Clinical Nutrition, Supplement* 2012;7:217.

Lopez-Ridaura R, Willett WC, Rimm EB, Liu S, Stampfer MJ, Manson JE, et al. Magnesium intake and risk of type 2 diabetes in men and women. *Diabetes Care* 2004;27:134-40.

Lophatananon A, Archer J, Easton D, Pocock R, Dearnaley D, Guy M, et al. Dietary fat and early-onset prostate cancer risk. *Br J Nutr* 2010;103:1375-80.

Lu M, Cho E, Taylor A, Hankinson SE, Willett WC, Jacques PF. Prospective study of dietary fat and risk of cataract extraction among US women. *Am J Epidemiol*. 2005;161:948-59.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Lubans DR, Morgan PJ, Collins CE, Warren JM, Callister R. Exploring the mechanisms of weight loss in the SHED-IT intervention for overweight men: A mediation analysis. *The International Journal of Behavioral Nutrition and Physical Activity* 2009;6.

Ludwig DS, Pereira MA, Kroenke CH, Hilner JE, Van Horn L, Slattery ML, et al. Dietary fiber, weight gain, and cardiovascular disease risk factors in young adults. *JAMA* 1999;282:1539-46.

Ludwig DS, Peterson KE, Gortmaker SL. Relation between consumption of sugar-sweetened drinks and childhood obesity: a prospective, observational analysis. *Lancet* 2001;357:505-8.

Lund E, Bonna KH. Reduced breast cancer mortality among fishermen's wives in Norway. *Cancer Causes Control* 1993;4:283-7.

Lundgren H, Bengtsson C, Blohme G, Isaksson B, Lapidus L, Lenner RA, et al. Dietary habits and incidence of noninsulin-dependent diabetes mellitus in a population study of women in Gothenburg, Sweden. *Am J Clin Nutr.* 1989;49:708-12.

Lydeking-Olsen E, Beck-Jensen JE, Setchell KD, Holm-Jensen T. Soymilk or progesterone for prevention of bone loss--a 2 year randomized, placebo-controlled trial. *Eur J Nutr.* 2004;43:246-57.

M AM, Pera G, Agudo A, Bueno-de-Mesquita HB, Palli D, Boeing H, et al. Cereal fiber intake may reduce risk of gastric adenocarcinomas: the EPIC-EURGAST study. *Int J Cancer* 2007;121:1618-23.

Ma E, Sasazuki S, Inoue M, Iwasaki M, Sawada N, Takachi R, et al. High dietary intake of magnesium may decrease risk of colorectal cancer in Japanese men. *J Nutr* 2010;140:779-85.

Ma Y, Olendzki BC, Chiriboga D, Rosal M, Sinagra E, Crawford S, et al. PDA-assisted low glycemic index dietary intervention for type II diabetes: A pilot study. *European Journal of Clinical Nutrition* 2006;60:1235-1243.

Maclure M, Travis LB, Willett W, MacMahon B. A prospective cohort study of nutrient intake and age at menarche. *Am J Clin Nutr.* 1991;54:649-56.

Macquart-Moulin G, Riboli E, Cornee J, Charnay B, Berthezene P, Day N. Case-control study on colorectal cancer and diet in Marseilles. *Int J Cancer* 1986;38:183-91.

Madsen KA, Garber AK, Mietus-Snyder ML, Orrell-Valente JK, Tran CT, Wlasiuk L, et al. A clinic-based lifestyle intervention for pediatric obesity: efficacy and behavioral and biochemical predictors of response. *J Pediatr Endocrinol Metab* 2009;22:805-14.

Magarey AM, Daniels LA, Boulton TJ, Cockington RA. Does fat intake predict adiposity in healthy children and adolescents aged 2--15 y? A longitudinal analysis. *Eur J Clin Nutr* 2001;55:471-81.

Magni E, Bianchetti A, Rozzini R, Trabucchi M. Influence of nutritional intake on 6-year mortality in an Italian elderly population. *J Nutr Elder* 1994;13:25-34.

Maheu E, Mazieres B, Valat JP, Loyau G, Le Loet X, Bourgeois P, et al. Symptomatic efficacy of avocado/soybean unsaponifiables in the treatment of osteoarthritis of the knee and hip: a prospective, randomized, double-blind, placebo-controlled, multicenter clinical trial with a six-

month treatment period and a two-month followup demonstrating a persistent effect. *Arthritis Rheum.* 1998;41:81-91.

Mai V, Kant AK, Flood A, Lacey JV, Jr., Schairer C, Schatzkin A. Diet quality and subsequent cancer incidence and mortality in a prospective cohort of women. *Int J Epidemiol* 2005;34:54-60.

Mai XM, Becker AB, Sellers EA, Liem JJ, Kozyrskyj AL. Infrequent milk consumption plus being overweight may have great risk for asthma in girls. *Allergy* 2007;62:1295-301.

Malik VS, Sun Q, van Dam RM, Rimm EB, Willett WC, Rosner B, et al. Adolescent dairy product consumption and risk of type 2 diabetes in middle-aged women. *Am J Clin Nutr* 2011;94:854-61.

Manger MS, Strand E, Ebbing M, Seifert R, Refsum H, Nordrehaug JE, et al. Dietary intake of n-3 long-chain polyunsaturated fatty acids and coronary events in Norwegian patients with coronary artery disease. *Am J Clin Nutr* 2010;92:244-51.

Manios Y, Moschonis G, Koutsikas K, Papoutsou S, Petraki I, Bellou E, et al. Changes in body composition following a dietary and lifestyle intervention trial: the postmenopausal health study. *Maturitas.* 2009;62:58-65.

Mann JI, Appleby PN, Key TJ, Thorogood M. Dietary determinants of ischaemic heart disease in health conscious individuals. *Heart* 1997;78:450-5.

Mannisto S, Kontto J, Kataja-Tuomola M, Albanes D, Virtamo J. High processed meat consumption is a risk factor of type 2 diabetes in the Alpha-Tocopherol, Beta-Carotene Cancer Prevention study. *Br J Nutr* 2010;103:1817-22.

Marckmann P, Gronbaek M. Fish consumption and coronary heart disease mortality. A systematic review of prospective cohort studies. *Eur J Clin Nutr* 1999;53:585-90.

Mariotti R, Castrogiovanni F, Canale ML, Borelli G, Rondinini L. Weight loss and quality of life in chronic heart failure patients. *Journal of Cardiovascular Medicine* 2008;9:576-580.

Markovits N, Ben Amotz A, Levy Y. The effect of tomato-derived lycopene on low carotenoids and enhanced systemic inflammation and oxidation in severe obesity. *Isr Med Assoc J* 2009;11:598-601.

Marniemi J, Seppanen A, Hakala P. Long-term effects on lipid metabolism of weight reduction on lactovegetarian and mixed diet. *Int J Obes* 1990;14:113-25.

Martin LJ, Li Q, Melnichouk O, Greenberg C, Minkin S, Hislop G, et al. A randomized trial of dietary intervention for breast cancer prevention. *Cancer Research* 2011;71:123-133.

Martinez-Gonzalez MA, de la Fuente-Arrillaga C, Nunez-Cordoba JM, Basterra-Gortari FJ, Beunza JJ, Vazquez Z, et al. Adherence to Mediterranean diet and risk of developing diabetes: prospective cohort study. *BMJ* 2008;336:1348-51.

Martinez-Gonzalez MA, Fernandez-Garcia J, Sanchez-Izquierdo F, Lardelli-Claret P, Jimenez Moleon J, Galvez-Vargas R. Life-style factors associated with changes in serum lipids in a follow-up study of cardiovascular risk factors. *Eur J Epidemiol.* 1998;14:525-33.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Martinez-Gonzalez MA, Sanchez-Villegas A, De Irala J, Marti A, Martinez JA. Mediterranean diet and stroke: objectives and design of the SUN project. *Seguimiento Universidad de Navarra. Nutr Neurosci* 2002;5:65-73.

Martinez-Gonzalez MA, Tortosa A, Basterra-Gortari FJ, Bes-Rastrollo M. Mediterranean diet inversely associated with the incidence of metabolic syndrome: The sun prospective cohort response to giugliano, ceriello, and esposito. *Diabetes Care* 2008;31:E37.

Masala G, Ceroti M, Pala V, Krogh V, Vineis P, Sacerdote C, et al. A dietary pattern rich in olive oil and raw vegetables is associated with lower mortality in Italian elderly subjects. *Br J Nutr* 2007;98:406-15.

Maskarinec G, Steude JS, Franke AA, Cooney RV. Inflammatory markers in a 2-year soy intervention among premenopausal women. *SOURCE Journal of Inflammation* (2009) 6 Article Number: 9. Date of Publication: 2009 2009.

Massey LK. Dairy food consumption, blood pressure and stroke. *J Nutr* 2001;131:1875-8.

Matheson EM, Mainous AG, 3rd, Hill EG, Carnemolla MA. Shellfish consumption and risk of coronary heart disease. *J Am Diet Assoc.* 2009;109:1422-6.

Mattisson I, Wirfalt E, Johansson U, Gullberg B, Olsson H, Berglund G. Intakes of plant foods, fibre and fat and risk of breast cancer--a prospective study in the Malmo Diet and Cancer cohort. *Br J Cancer* 2004;90:122-7.

Maynard M, Gunnell D, Emmett P, Frankel S, Davey Smith G. Fruit, vegetables, and antioxidants in childhood and risk of adult cancer: the Boyd Orr cohort. *J Epidemiol Community Health* 2003;57:218-25.

McAlindon TE, Felson DT, Zhang Y, Hannan MT, Aliabadi P, Weissman B, et al. Relation of dietary intake and serum levels of vitamin D to progression of osteoarthritis of the knee among participants in the Framingham Study. *Ann Intern Med.* 1996;125:353-9.

McCarter DF. Low-carbohydrate diet effective for adults. *Journal of Family Practice* 2003;52:515-516.

McClure CW, Brusch CA. Treatment of oral syndrome obesity with non-traditional appetite control plan. *J Am Med Womens Assoc* 1973;28:239-48.

McCullough ML, Bandera EV, Patel R, Patel AV, Gansler T, Kushi LH, et al. A prospective study of fruits, vegetables, and risk of endometrial cancer. *Am J Epidemiol* 2007;166:902-11.

McCullough ML, Feskanich D, Rimm EB, Giovannucci EL, Ascherio A, Variyam JN, et al. Adherence to the Dietary Guidelines for Americans and risk of major chronic disease in men. *Am J Clin Nutr.* 2000;72:1223-31.

McCullough ML, Feskanich D, Stampfer MJ, Rosner BA, Hu FB, Hunter DJ, et al. Adherence to the Dietary Guidelines for Americans and risk of major chronic disease in women. *Am J Clin Nutr.* 2000;72:1214-22.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

McCullough ML, Robertson AS, Chao A, Jacobs EJ, Stampfer MJ, Jacobs DR, et al. A prospective study of whole grains, fruits, vegetables and colon cancer risk. *Cancer Causes Control* 2003;14:959-70.

McCullough ML, Robertson AS, Jacobs EJ, Chao A, Calle EE, Thun MJ. A prospective study of diet and stomach cancer mortality in United States men and women. *Cancer Epidemiol Biomarkers Prev* 2001;10:1201-5.

McCullough ML, Robertson AS, Rodriguez C, Jacobs EJ, Chao A, Carolyn J, et al. Calcium, vitamin D, dairy products, and risk of colorectal cancer in the Cancer Prevention Study II Nutrition Cohort (United States). *Cancer Causes Control* 2003;14:1-12.

McDoniel SO, Hammond RS. A 24-week randomised controlled trial comparing usual care and metabolic-based diet plans in obese adults. *Int J Clin Pract* 2010;64:1503-11.

McGlynn KA, Sakoda LC, Rubertone MV, Sesterhenn IA, Lyu C, Graubard BI, et al. Body size, dairy consumption, puberty, and risk of testicular germ cell tumors. *Am J Epidemiol.* 2007;165:355-63.

McGuire HL, Svetkey LP, Harsha DW, Elmer PJ, Appel LJ, Ard JD. Comprehensive lifestyle modification and blood pressure control: a review of the PREMIER trial. *J Clin Hypertens (Greenwich)* 2004;6:383-90.

McKeown NM, Meigs JB, Liu S, Rogers G, Yoshida M, Saltzman E, et al. Dietary carbohydrates and cardiovascular disease risk factors in the Framingham offspring cohort. *J Am Coll Nutr.* 2009;28:150-8.

McLaren BA. Nutritional control of overweight. *Can J Public Health* 1967;58:483-5.

McLaughlin JM, Olivo-Marston S, Vitolins MZ, Bittoni M, Reeves KW, Degraffinreid CR, et al. Effects of tomato- and soy-rich diets on the IGF-I hormonal network: a crossover study of postmenopausal women at high risk for breast cancer. *Cancer Prev Res (Phila)* 2011;4:702-10.

McTiernan A, Wactawski-Wende J, Wu L, Rodabough RJ, Watts NB, Tylavsky F, et al. Low-fat, increased fruit, vegetable, and grain dietary pattern, fractures, and bone mineral density: the Women's Health Initiative Dietary Modification Trial. *Am J Clin Nutr* 2009;89:1864-76.

Meinhold CL, Dodd KW, Jiao L, Flood A, Shikany JM, Genkinger JM, et al. Available carbohydrates, glycemic load, and pancreatic cancer: is there a link? *Am J Epidemiol.* 2010;171:1174-82.

Melanson EL, Astrup A, Donahoo WT. The relationship between dietary fat and fatty acid intake and body weight, diabetes, and the metabolic syndrome. *Annals of Nutrition and Metabolism* 2009;55:229-243.

Melanson KJ, Angelopoulos TJ, Nguyen VT, Martini M, Zukley L, Lowndes J, et al. Consumption of whole-grain cereals during weight loss: effects on dietary quality, dietary fiber, magnesium, vitamin B-6, and obesity. *J Am Diet Assoc* 2006;106:1380-8; quiz 1389-90.

Mellen PB, Liese AD, Toozee JA, Vitolins MZ, Wagenknecht LE, Herrington DM. Whole-grain intake and carotid artery atherosclerosis in a multiethnic cohort: the Insulin Resistance Atherosclerosis Study. *Am J Clin Nutr* 2007;85:1495-502.

Meng X, Zhu K, Devine A, Kerr DA, Binns CW, Prince RL. A 5-year cohort study of the effects of high protein intake on lean mass and BMC in elderly postmenopausal women. *J Bone Miner Res*. 2009;24:1827-34.

Messier SP. Knee strengthening exercises at home, but not dietary intervention, reduce knee pain and improve knee function in overweight and obese people over 2 years. *Evidence-Based Medicine* 2010;15:21-22.

Meyer BJ, van der Merwe M, Du Plessis DG, de Bruin EJ, Meyer AC. Some physiological effects of a mainly fruit diet in man. *S Afr Med J*. 1971;45:191-5.

Meyer F, Bairati I, Fradet Y, Moore L. Dietary energy and nutrients in relation to preclinical prostate cancer. *Nutr Cancer*. 1997;29:120-6.

Meyer F, Bairati I, Shadmani R, Fradet Y, Moore L. Dietary fat and prostate cancer survival. *Cancer Causes Control* 1999;10:245-51.

Meyer HE, Pedersen JI, Loken EB, Tverdal A. Dietary factors and the incidence of hip fracture in middle-aged Norwegians. A prospective study. *Am J Epidemiol* 1997;145:117-23.

Meyer KA, Kushi LH, Jacobs DR, Jr., Folsom AR. Dietary fat and incidence of type 2 diabetes in older Iowa women. *Diabetes Care* 2001;24:1528-35.

Meyer KA, Kushi LH, Jacobs DR, Jr., Slavin J, Sellers TA, Folsom AR. Carbohydrates, dietary fiber, and incident type 2 diabetes in older women. *Am J Clin Nutr* 2000;71:921-30.

Micha R, Mozaffarian D. Saturated fat and cardiometabolic risk factors, coronary heart disease, stroke, and diabetes: A fresh look at the evidence. *Lipids* 2010;45:893-905.

Michaelsson K, Holmberg L, Mallmin H, Sorensen S, Wolk A, Bergstrom R, et al. Diet and hip fracture risk: a case-control study. Study Group of the Multiple Risk Survey on Swedish Women for Eating Assessment. *Int J Epidemiol* 1995;24:771-82.

Michaud DS, Fuchs CS, Liu S, Willett WC, Colditz GA, Giovannucci E. Dietary glycemic load, carbohydrate, sugar, and colorectal cancer risk in men and women. *Cancer Epidemiol Biomarkers Prev*. 2005;14:138-47.

Michaud DS, Liu S, Giovannucci E, Willett WC, Colditz GA, Fuchs CS. Dietary sugar, glycemic load, and pancreatic cancer risk in a prospective study. *J Natl Cancer Inst*. 2002;94:1293-300.

Michaud DS, Spiegelman D, Clinton SK, Rimm EB, Willett WC, Giovannucci EL. Fruit and vegetable intake and incidence of bladder cancer in a male prospective cohort. *J Natl Cancer Inst* 1999;91:605-13.

Michels KB. Diet and cancer: Current knowledge, methodologic pitfalls and future directions. *International Journal of Cancer* 2005;116:665-666.

Michels KB, Fuchs CS, Giovannucci E, Colditz GA, Hunter DJ, Stampfer MJ, et al. Fiber intake and incidence of colorectal cancer among 76,947 women and 47,279 men. *Cancer Epidemiol Biomarkers Prev* 2005;14:842-9.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Michels KB, Giovannucci E, Chan AT, Singhanian R, Fuchs CS, Willett WC. Fruit and vegetable consumption and colorectal adenomas in the Nurses' Health Study. *Cancer Res* 2006;66:3942-53.

Michels KB, Mohllajee AP, Roset-Bahmanyar E, Beehler GP, Moysich KB. Diet and breast cancer: a review of the prospective observational studies. *Cancer* 2007;109:2712-49.

Michels KB, Wolk A. A prospective study of variety of healthy foods and mortality in women. *Int J Epidemiol* 2002;31:847-54.

Mihos C, Mariolis A, Manios Y, Naska A, Arapaki A, Mariolis-Sapsakos T, et al. Evaluation of a nutrition intervention in adolescents of an urban area in Greece: short- and long-term effects of the VYRONAS study. *Public Health Nutr* 2010;13:712-9.

Mikkila V, Rasanen L, Raitakari OT, Pietinen P, Viikari J. Longitudinal changes in diet from childhood into adulthood with respect to risk of cardiovascular diseases: The Cardiovascular Risk in Young Finns Study. *Eur J Clin Nutr*. 2004;58:1038-45.

Milder IE, Feskens EJ, Arts IC, Bueno-de-Mesquita HB, Hollman PC, Kromhout D. Intakes of 4 dietary lignans and cause-specific and all-cause mortality in the Zutphen Elderly Study. *Am J Clin Nutr* 2006;84:400-5.

Millen BE, Quatromoni PA. Nutritional research within the Framingham Heart Study. *J Nutr Health Aging* 2001;5:139-43.

Mills PK, Annegers JF, Phillips RL. Animal product consumption and subsequent fatal breast cancer risk among Seventh-day Adventists. *Am J Epidemiol* 1988;127:440-53.

Mirmiran P, Ramezankhani A, Azizi F. Combined effects of saturated fat and cholesterol intakes on serum lipids: Tehran Lipid and Glucose Study. *Nutrition*. 2009;25:526-31.

Missmer SA, Chavarro JE, Malspeis S, Bertone-Johnson ER, Hornstein MD, Spiegelman D, et al. A prospective study of dietary fat consumption and endometriosis risk. *Hum Reprod*. 2010;25:1528-35.

Mitrou PN, Kipnis V, Thiebaut AC, Reedy J, Subar AF, Wirfalt E, et al. Mediterranean dietary pattern and prediction of all-cause mortality in a US population: results from the NIH-AARP Diet and Health Study. *Arch Intern Med* 2007;167:2461-8.

Miura K, Greenland P, Stamler J, Liu K, Daviglius ML, Nakagawa H. Relation of vegetable, fruit, and meat intake to 7-year blood pressure change in middle-aged men: the Chicago Western Electric Study. *Am J Epidemiol* 2004;159:572-80.

Mommers M, Schouten LJ, Goldbohm RA, van den Brandt PA. Consumption of vegetables and fruits and risk of ovarian carcinoma. *Cancer* 2005;104:1512-9.

Monma Y, Niu K, Iwasaki K, Tomita N, Nakaya N, Hozawa A, et al. Dietary patterns associated with fall-related fracture in elderly Japanese: a population based prospective study. *BMC Geriatr* 2010;10:31.

Montonen J, Jarvinen R, Heliovaara M, Reunanen A, Aromaa A, Knekt P. Food consumption and the incidence of type II diabetes mellitus. *Eur J Clin Nutr* 2005;59:441-8.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Montonen J, Knekt P, Jarvinen R, Aromaa A, Reunanen A. Whole-grain and fiber intake and the incidence of type 2 diabetes. *Am J Clin Nutr* 2003;77:622-9.

Montonen J, Knekt P, Jarvinen R, Reunanen A. Dietary antioxidant intake and risk of type 2 diabetes. *Diabetes Care* 2004;27:362-6.

Moore C, Gitau R, Goff L, Lewis FJ, Griffin MD, Chatfield MD, et al. Successful manipulation of the quality and quantity of fat and carbohydrate consumed by free-living individuals using a food exchange model. *Journal of Nutrition* 2009;139:1534-1540.

Moore CS, Bryant SP, Mishra GD, Krebs JD, Browning LM, Miller GJ, et al. Oily fish reduces plasma triacylglycerols: a primary prevention study in overweight men and women. *Nutrition* 2006;22:1012-24.

Moore LL, Bradlee ML, Gao D, Singer MR. Low dairy intake in early childhood predicts excess body fat gain. *Obesity (Silver Spring)* 2006;14:1010-8.

Moore LL, Singer MR, Bradlee ML, Djousse L, Proctor MH, Cupples LA, et al. Intake of fruits, vegetables, and dairy products in early childhood and subsequent blood pressure change. *Epidemiology*. 2005;16:4-11.

Moore TJ, Alsabeeh N, Apovian CM, Murphy MC, Coffman GA, Cullum-Dugan D, et al. Weight, blood pressure, and dietary benefits after 12 months of a web-based nutrition education program (DASH for health): Longitudinal observational study. *SOURCE Journal of Medical Internet Research* (2008) 10:4 Article Number: e52. Date of Publication: October-December 2008 2008.

Morales E, Valero MA, Leon M, Hernandez E, Praga M. Beneficial effects of weight loss in overweight patients with chronic proteinuric nephropathies. *Am J Kidney Dis* 2003;41:319-27.

Moretti L, Canada T. A randomized study comparing the effects of a low-carbohydrate diet and a conventional diet on lipoprotein subfractions and C-reactive protein levels in patients with severe obesity: Commentary. *Nutrition in Clinical Practice* 2006;21:187-188.

Morgan LM, Griffin BA, Millward DJ, DeLooy A, Fox KR, Baic S, et al. Comparison of the effects of four commercially available weight-loss programmes on lipid-based cardiovascular risk factors. *Public Health Nutr* 2009;12:799-807.

Morgan PJ, Lubans DR, Collins CE, Warren JM, Callister R. 12-month outcomes and process evaluation of the SHED-IT RCT: an internet-based weight loss program targeting men. *Obesity (Silver Spring)* 2011;19:142-51.

Mosdol A, Witte DR, Frost G, Marmot MG, Brunner EJ. Dietary glycemic index and glycemic load are associated with high-density-lipoprotein cholesterol at baseline but not with increased risk of diabetes in the Whitehall II study. *Am J Clin Nutr* 2007;86:988-94.

Mozaffarian D, Cao H, King IB, Lemaitre RN, Song X, Siscovick DS, et al. Trans-palmitoleic acid, metabolic risk factors, and new-onset diabetes in U.S. adults: a cohort study. *Ann Intern Med* 2010;153:790-9.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Mozaffarian D, Hao T, Rimm EB, Willett WC, Hu FB. Changes in diet and lifestyle and long-term weight gain in women and men. *N Engl J Med* 2011;364:2392-404.

Mozaffarian D, Lemaitre RN, Kuller LH, Burke GL, Tracy RP, Siscovick DS. Cardiac benefits of fish consumption may depend on the type of fish meal consumed: the Cardiovascular Health Study. *Circulation* 2003;107:1372-7.

Mozaffarian D, Longstreth WT, Jr., Lemaitre RN, Manolio TA, Kuller LH, Burke GL, et al. Fish consumption and stroke risk in elderly individuals: the cardiovascular health study. *Arch Intern Med* 2005;165:200-6.

Munger RG, Cerhan JR, Chiu BC. Prospective study of dietary protein intake and risk of hip fracture in postmenopausal women. *Am J Clin Nutr* 1999;69:147-52.

Murakami K, Okubo H, Sasaki S. Effect of dietary factors on incidence of type 2 diabetes: a systematic review of cohort studies. *J Nutr Sci Vitaminol (Tokyo)* 2005;51:292-310.

Murakami K, Sasaki S. Dietary intake and depressive symptoms: a systematic review of observational studies. *Mol Nutr Food Res* 2010;54:471-88.

Murphy KJ, Meyer BJ, Mori TA, Burke V, Mansour J, Patch CS, et al. Impact of foods enriched with n-3 long-chain polyunsaturated fatty acids on erythrocyte n-3 levels and cardiovascular risk factors. *Br J Nutr* 2007;97:749-57.

Murtaugh MA, Jacobs DR, Jr., Jacob B, Steffen LM, Marquart L. Epidemiological support for the protection of whole grains against diabetes. *Proc Nutr Soc* 2003;62:143-9.

Must A, Lipman RD. Childhood energy intake and cancer mortality in adulthood. *Nutr Rev* 1999;57:21-4.

Musunuru K. Atherogenic dyslipidemia: Cardiovascular risk and dietary intervention. *Lipids* 2010;45:907-914.

Mutungi G, Ratliff J, Puglisi M, Torres-Gonzalez M, Vaishnav U, Leite JO, et al. Dietary cholesterol from eggs increases plasma HDL cholesterol in overweight men consuming a carbohydrate-restricted diet. *J Nutr* 2008;138:272-6.

Nader PR, Stone EJ, Lytle LA, Perry CL, Osganian SK, Kelder S, et al. Three-year maintenance of improved diet and physical activity: the CATCH cohort. *Child and Adolescent Trial for Cardiovascular Health. Arch Pediatr Adolesc Med.* 1999;153:695-704.

Nagata C, Takatsuka N, Kawakami N, Shimizu H. Association of diet with the onset of menopause in Japanese women. *Am J Epidemiol.* 2000;152:863-7.

Nagata C, Takatsuka N, Kawakami N, Shimizu H. A prospective cohort study of soy product intake and stomach cancer death. *Br J Cancer* 2002;87:31-6.

Nagura J, Iso H, Watanabe Y, Maruyama K, Date C, Toyoshima H, et al. Fruit, vegetable and bean intake and mortality from cardiovascular disease among Japanese men and women: the JACC Study. *Br J Nutr* 2009;102:285-92.

Nakamura K, Nagata C, Oba S, Takatsuka N, Shimizu H. Fruit and vegetable intake and mortality from cardiovascular disease are inversely associated in Japanese women but not in men. *J Nutr* 2008;138:1129-34.

Nakamura Y, Okamura T, Tamaki S, Kadowaki T, Hayakawa T, Kita Y, et al. Egg consumption, serum cholesterol, and cause-specific and all-cause mortality: the National Integrated Project for Prospective Observation of Non-communicable Disease and Its Trends in the Aged, 1980 (NIPPON DATA80). *Am J Clin Nutr* 2004;80:58-63.

Nanri A, Mizoue T, Noda M, Takahashi Y, Kirii K, Inoue M, et al. Magnesium intake and type II diabetes in Japanese men and women: the Japan Public Health Center-based Prospective Study. *Eur J Clin Nutr* 2010;64:1244-7.

Nathan I, Hackett AF, Kirby S. A longitudinal study of the growth of matched pairs of vegetarian and omnivorous children, aged 7-11 years, in the north-west of England. *Eur J Clin Nutr* 1997;51:20-5.

Nerfeldt P, Nilsson BY, Mayor L, Udden J, Friberg D. A two-year weight reduction program in obese sleep apnea patients. *J Clin Sleep Med* 2010;6:479-86.

Ness AR. The Avon Longitudinal Study of Parents and Children (ALSPAC)--a resource for the study of the environmental determinants of childhood obesity. *Eur J Endocrinol* 2004;151 Suppl 3:U141-9.

Ness AR, Ashfield-Watt PAL, Whiting JM, Smith GD, Hughes J, Burr ML. The long-term effect of dietary advice on the diet of men with angina: the diet and angina randomized trial. *Journal of Human Nutrition & Dietetics* 2004;17:117-119.

Ness AR, Gunnell D, Hughes J, Elwood PC, Smith GD, Burr ML. Height, body mass index, and survival in men with coronary disease: Follow up of the diet and reinfarction trial (DART). *Journal of Epidemiology and Community Health* 2002;56:218-219.

Ness AR, Powles JW. Fruit and vegetables, and cardiovascular disease: a review. *Int J Epidemiol* 1997;26:1-13.

Neuhouser ML, Barnett MJ, Kristal AR, Ambrosone CB, King I, Thornquist M, et al. (n-6) PUFA increase and dairy foods decrease prostate cancer risk in heavy smokers. *J Nutr* 2007;137:1821-7.

Neutzling MB, Taddei JA, Gigante DP. Risk factors of obesity among Brazilian adolescents: a case-control study. *Public Health Nutr* 2003;6:743-9.

Newby PK, Maras J, Bakun P, Muller D, Ferrucci L, Tucker KL. Intake of whole grains, refined grains, and cereal fiber measured with 7-d diet records and associations with risk factors for chronic disease. *Am J Clin Nutr* 2007;86:1745-53.

Newby PK, Muller D, Hallfrisch J, Qiao N, Andres R, Tucker KL. Dietary patterns and changes in body mass index and waist circumference in adults. *Am J Clin Nutr* 2003;77:1417-25.

Newby PK, Peterson KE, Berkey CS, Leppert J, Willett WC, Colditz GA. Dietary composition and weight change among low-income preschool children. *Arch Pediatr Adolesc Med* 2003;157:759-64.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Newby PK, Tucker KL, Wolk A. Risk of overweight and obesity among semivegetarian, lactovegetarian, and vegan women. *Am J Clin Nutr* 2005;81:1267-74.

Ngoan LT, Mizoue T, Fujino Y, Tokui N, Yoshimura T. Dietary factors and stomach cancer mortality. *Br J Cancer* 2002;87:37-42.

Nicklas TA, Dwyer J, Mitchell P, Zive M, Montgomery D, Lytle L, et al. Impact of fat reduction on micronutrient density of children's diets: the CATCH Study. *Prev Med* 1996;25:478-85.

Nicodemus KK, Sweeney C, Folsom AR. Evaluation of dietary, medical and lifestyle risk factors for incident kidney cancer in postmenopausal women. *Int J Cancer*. 2004;108:115-21.

Nield L, Moore HJ, Hooper L, Cruickshank JK, Vyas A, Whittaker V, et al. Dietary advice for treatment of type 2 diabetes mellitus in adults. SOURCE Cochrane Database of Systematic Reviews (2007) :3 Article Number: CD004097. Date of Publication: 2007 2007.

Nielsen JV, Jonsson E, Nilsson AK. Lasting improvement of hyperglycaemia and bodyweight: low-carbohydrate diet in type 2 diabetes--a brief report. *Ups J Med Sci* 2005;110:69-73.

Niemeier HM, Raynor HA, Lloyd-Richardson EE, Rogers ML, Wing RR. Fast food consumption and breakfast skipping: predictors of weight gain from adolescence to adulthood in a nationally representative sample. *J Adolesc Health* 2006;39:842-9.

Nilsson LM, Winkvist A, Brustad M, Jansson JH, Johansson I, Lenner P, et al. A traditional Sami diet score as a determinant of mortality in a general northern Swedish population. *Int J Circumpolar Health* 2012;71:1-12.

Nimptsch K, Berg-Beckhoff G, Linseisen J. Effect of dietary fatty acid intake on prospective weight change in the Heidelberg cohort of the European Prospective Investigation into Cancer and Nutrition. *Public Health Nutr* 2010;13:1636-46.

Nishimoto S. Dietary treatment of the cardiovascular patient with the hyperlipidemia and the long term dietary control of the hyperlipidemia. *Jpn Circ J* 1975;39:345-9.

Nissinen K, Mikkilä V, Mannisto S, Lahti-Koski M, Rasanen L, Viikari J, et al. Sweets and sugar-sweetened soft drink intake in childhood in relation to adult BMI and overweight. The Cardiovascular Risk in Young Finns Study. *Public Health Nutr* 2009;12:2018-26.

Njike V, Faridi Z, Dutta S, Gonzalez-Simon AL, Katz DL. Daily egg consumption in hyperlipidemic adults - Effects on endothelial function and cardiovascular risk. SOURCE Nutrition Journal (2010) 9:1 Article Number: 28. Date of Publication: 2010 2010.

Nkondjock A, Robidoux A, Paredes Y, Narod SA, Ghadirian P. Diet, lifestyle and BRCA-related breast cancer risk among French-Canadians. *Breast Cancer Res Treat*. 2006;98:285-94.

Nomura AM, Hankin JH, Lee J, Stemmermann GN. Cohort study of tofu intake and prostate cancer: no apparent association. *Cancer Epidemiol Biomarkers Prev* 2004;13:2277-9.

Nomura AMY, Hankin JH, Henderson BE, Wilkens LR, Murphy SP, Pike MC, et al. Dietary fiber and colorectal cancer risk: The multiethnic cohort study. *Cancer Causes and Control* 2007;18:753-764.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Norat T, Bingham S, Ferrari P, Slimani N, Jenab M, Mazuir M, et al. Meat, fish, and colorectal cancer risk: The European prospective investigation into cancer and nutrition. *Journal of the National Cancer Institute* 2005;97:906-916.

Norrish AE, Jackson RT, Sharpe SJ, Skeaff CM. Prostate cancer and dietary carotenoids. *Am J Epidemiol* 2000;151:119-23.

Nothlings U, Murphy SP, Wilkens LR, Henderson BE, Kolonel LN. Dietary glycemic load, added sugars, and carbohydrates as risk factors for pancreatic cancer: the Multiethnic Cohort Study. *Am J Clin Nutr*. 2007;86:1495-501.

Nowicka P, Höglund P, Pietrobelli A, Lissau I, Flodmark CE. Family Weight School treatment: 1-year results in obese adolescents. *International journal of pediatric obesity : IJPO : an official journal of the International Association for the Study of Obesity* 2008:141-7.

Nunez-Cordoba JM, Alonso A, Beunza JJ, Palma S, Gomez-Gracia E, Martinez-Gonzalez MA. Role of vegetables and fruits in Mediterranean diets to prevent hypertension. *Eur J Clin Nutr* 2009;63:605-12.

Nydahl M, Gustafsson IB, Eriksson H, Vessby B, Bjernulf A. Effects of a simplified programme for dietary treatment of hyperlipidaemia in primary health care. *Scand J Prim Health Care*. 1994;12:128-36.

Oba S, Nagata C, Nakamura K, Fujii K, Kawachi T, Takatsuka N, et al. Diet based on the Japanese Food Guide Spinning Top and subsequent mortality among men and women in a general Japanese population. *J Am Diet Assoc* 2009;109:1540-7.

Obarzanek E, Hunsberger SA, Van Horn L, Hartmuller VV, Barton BA, Stevens VJ, et al. Safety of a fat-reduced diet: the Dietary Intervention Study in Children (DISC). *Pediatrics*. 1997;100:51-9.

Obrador A. Fibre and colorectal cancer: a controversial question. *Br J Nutr* 2006;96 Suppl 1:S46-8.

Ochner CN, Lowe MR. Self-reported changes in dietary calcium and energy intake predict weight regain following a weight loss diet in obese women. *J Nutr* 2007;137:2324-8.

Ocke MC, Bueno-de-Mesquita HB, Feskens EJ, van Staveren WA, Kromhout D. Repeated measurements of vegetables, fruits, beta-carotene, and vitamins C and E in relation to lung cancer. The Zutphen Study. *Am J Epidemiol* 1997;145:358-65.

Ocke MC, Kromhout D, Menotti A, Aravanis C, Blackburn H, Buzina R, et al. Average intake of anti-oxidant (pro)vitamins and subsequent cancer mortality in the 16 cohorts of the Seven Countries Study. *Int J Cancer* 1995;61:480-4.

Oddy WH, Robinson M, Ambrosini GL, O'Sullivan TA, de Klerk NH, Beilin LJ, et al. The association between dietary patterns and mental health in early adolescence. *Prev Med*. 2009;49:39-44.

Oh K, Hu FB, Cho E, Rexrode KM, Stampfer MJ, Manson JE, et al. Carbohydrate intake, glycemic index, glycemic load, and dietary fiber in relation to risk of stroke in women. *Am J Epidemiol*. 2005;161:161-9.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Oh K, Willett WC, Fuchs CS, Giovannucci EL. Glycemic index, glycemic load, and carbohydrate intake in relation to risk of distal colorectal adenoma in women. *Cancer Epidemiol Biomarkers Prev.* 2004;13:1192-8.

Ohlin A, Rossner S. Trends in eating patterns, physical activity and socio-demographic factors in relation to postpartum body weight development. *Br J Nutr.* 1994;71:457-70.

Oomen CM, Feskens EJ, Rasanen L, Fidanza F, Nissinen AM, Menotti A, et al. Fish consumption and coronary heart disease mortality in Finland, Italy, and The Netherlands. *Am J Epidemiol* 2000;151:999-1006.

Osganian SK, Stampfer MJ, Rimm E, Spiegelman D, Hu FB, Manson JE, et al. Vitamin C and risk of coronary heart disease in women. *J Am Coll Cardiol* 2003;42:246-52.

Osler M, Andreasen AH, Hoidrup S. No inverse association between fish consumption and risk of death from all-causes, and incidence of coronary heart disease in middle-aged, Danish adults. *J Clin Epidemiol* 2003;56:274-9.

Osler M, Heitmann BL, Gerdes LU, Jorgensen LM, Schroll M. Dietary patterns and mortality in Danish men and women: a prospective observational study. *Br J Nutr* 2001;85:219-25.

Osler M, Schroll M. Diet and mortality in a cohort of elderly people in a north European community. *Int J Epidemiol* 1997;26:155-9.

Otsuka R, Imai T, Kato Y, Ando F, Shimokata H. Relationship between number of metabolic syndrome components and dietary factors in middle-aged and elderly Japanese subjects. *Hypertens Res.* 2010;33:548-54.

Padwal R. The Atkins diet led to more weight loss than the Zone diet in overweight and obese premenopausal women at 12 months. *Evidence-Based Medicine* 2007;12:138.

Paeratakul S, Popkin BM, Keyou G, Adair LS, Stevens J. Changes in diet and physical activity affect the body mass index of Chinese adults. *Int J Obes Relat Metab Disord.* 1998;22:424-31.

Pala V, Sieri S, Berrino F, Vineis P, Sacerdote C, Palli D, et al. Yogurt consumption and risk of colorectal cancer in the Italian European prospective investigation into cancer and nutrition cohort. *Int J Cancer* 2011;129:2712-9.

Palacios C, Joshipura KJ, Willett WC. Nutrition and health: Guidelines for dental practitioners. *Oral Diseases* 2009;15:369-381.

Pan A, Sun Q, Bernstein AM, Schulze MB, Manson JE, Stampfer MJ, et al. Red meat consumption and mortality: results from 2 prospective cohort studies. *Arch Intern Med* 2012;172:555-63.

Panagiotakos DB, Pitsavos C, Chrysohoou C, Skoumas I, Stefanadis C. Five-year incidence of cardiovascular disease and its predictors in Greece: the ATTICA study. *Vasc Med* 2008;13:113-21.

Panagiotakos DB, Pitsavos C, Chrysohoou C, Skoumas J, Stefanadis C. Status and management of blood lipids in Greek adults and their relation to socio-demographic, lifestyle and dietary factors: the ATTICA Study. *Blood lipids distribution in Greece. Atherosclerosis* 2004;173:353-61.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Panagiotakos DB, Zeimbekis A, Boutziouka V, Economou M, Kourlaba G, Toutouzas P, et al. Long-term fish intake is associated with better lipid profile, arterial blood pressure, and blood glucose levels in elderly people from Mediterranean islands (MEDIS epidemiological study). *Medical Science Monitor* 2007;13:CR307-CR312.

Pandey M, Shukla VK. Diet and gallbladder cancer: a case-control study. *Eur J Cancer Prev* 2002;11:365-8.

Papas MA, Giovannucci E, Platz EA. Fiber from fruit and colorectal neoplasia. *Cancer Epidemiol Biomarkers Prev* 2004;13:1267-70.

Park SK, Tucker KL, O'Neill MS, Sparrow D, Vokonas PS, Hu H, et al. Fruit, vegetable, and fish consumption and heart rate variability: the Veterans Administration Normative Aging Study. *Am J Clin Nutr* 2009;89:778-86.

Park SY, Murphy SP, Wilkens LR, Yamamoto JF, Sharma S, Hankin JH, et al. Dietary patterns using the Food Guide Pyramid groups are associated with sociodemographic and lifestyle factors: the multiethnic cohort study. *J Nutr*. 2005;135:843-9.

Park Y, Brinton LA, Subar AF, Hollenbeck A, Schatzkin A. Dietary fiber intake and risk of breast cancer in postmenopausal women: the National Institutes of Health-AARP Diet and Health Study. *Am J Clin Nutr* 2009;90:664-71.

Pascale RW, Wing RR, Butler BA, Mullen M, Bononi P. Effects of a behavioral weight loss program stressing calorie restriction versus calorie plus fat restriction in obese individuals with NIDDM or a family history of diabetes. *Diabetes Care* 1995;18:1241-8.

Patel AV, McCullough ML, Pavluck AL, Jacobs EJ, Thun MJ, Calle EE. Glycemic load, glycemic index, and carbohydrate intake in relation to pancreatic cancer risk in a large US cohort. *Cancer Causes and Control* 2007;18:287-294.

Patel PS, Forouhi NG, Kuijsten A, Schulze MB, van Woudenberg GJ, Ardanaz E, et al. The prospective association between total and type of fish intake and type 2 diabetes in 8 European countries: EPIC-InterAct Study. *Am J Clin Nutr* 2012;95:1445-53.

Patel PS, Sharp SJ, Luben RN, Khaw KT, Bingham SA, Wareham NJ, et al. Association between type of dietary fish and seafood intake and the risk of incident type 2 diabetes: the European prospective investigation of cancer (EPIC)-Norfolk cohort study. *Diabetes Care* 2009;32:1857-63.

Paterna S, Gaspare P, Fasullo S, Sarullo FM, Di Pasquale P. Normal-sodium diet compared with low-sodium diet in compensated congestive heart failure: Is sodium an old enemy or a new friend? *Clinical Science* 2008;114:221-230.

Payne ME, Anderson JJ, Steffens DC. Calcium and vitamin D intakes may be positively associated with brain lesions in depressed and nondepressed elders. *Nutr Res*. 2008;28:285-92.

Pereira MA, Kartashov AI, Ebbeling CB, Van Horn L, Slattery ML, Jacobs DR, Jr., et al. Fast-food habits, weight gain, and insulin resistance (the CARDIA study): 15-year prospective analysis. *Lancet* 2005;365:36-42.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Perez-Martinez P, Moreno-Conde M, Cruz-Teno C, Ruano J, Fuentes F, Delgado-Lista J, et al. Dietary fat differentially influences regulatory endothelial function during the postprandial state in patients with metabolic syndrome: From the LIPGENE study. *Atherosclerosis* 2010;209:533-538.

Peters ES, Lockett BG, Applebaum KM, Marsit CJ, McClean MD, Kelsey KT. Dairy products, leanness, and head and neck squamous cell carcinoma. *Head Neck* 2008;30:1193-205.

Pham TM, Fujino Y, Ide R, Kubo T, Shirane K, Tokui N, et al. Prospective study of vegetable consumption and liver cancer in Japan. *Int J Cancer* 2006;119:2408-11.

Pham TM, Fujino Y, Kikuchi S, Tamakoshi A, Matsuda S, Yoshimura T. Dietary patterns and risk of stomach cancer mortality: the Japan collaborative cohort study. *Ann Epidemiol* 2010;20:356-63.

Pham TM, Fujino Y, Kubo T, Ide R, Tokui N, Mizoue T, et al. Fish intake and the risk of fatal prostate cancer: findings from a cohort study in Japan. *Public Health Nutr* 2009;12:609-13.

Philippou E, Neary NM, Chaudhri O, Brynes AE, Dornhorst A, Leeds AR, et al. The effect of dietary glycemic index on weight maintenance in overweight subjects: a pilot study. *Obesity (Silver Spring)* 2009;17:396-401.

Phillips SM, Bandini LG, Cyr H, Colclough-Douglas S, Naumova E, Must A. Dairy food consumption and body weight and fatness studied longitudinally over the adolescent period. *Int J Obes Relat Metab Disord* 2003;27:1106-13.

Pijls LT, de Vries H, van Eijk JT, Donker AJ. Protein restriction, glomerular filtration rate and albuminuria in patients with type 2 diabetes mellitus: a randomized trial. *Eur J Clin Nutr* 2002;56:1200-7.

Pijls LTJ, De Vries H, Donker AJM, Van Eijk JTM. The effect of protein restriction on albuminuria in patients with type 2 diabetes mellitus: A randomized trial. *Nephrology Dialysis Transplantation* 1999;14:1445-1453.

Pilkington TR, Rosenoer VM, Gainsborough H, Carey M. Diet and weight-reduction in the obese. *Lancet* 1960:856-8.

Pitsavos C, Panagiotakos DB, Chrysohoou C, Papaioannou I, Papadimitriou L, Tousoulis D, et al. The adoption of mediterranean diet attenuates the development of acute coronary syndromes in people with the metabolic syndrome. *SOURCE Nutrition Journal* (2003) 2 (1-7) Article Number: 1. Date of Publication: 19 Mar 2003 2003.

Pitsavos C, Panagiotakos DB, Chrysohoou C, Skoumas J, Papaioannou I, Stefanadis C, et al. The effect of Mediterranean diet on the risk of the development of acute coronary syndromes in hypercholesterolemic people: a case-control study (CARDIO2000). *Coron Artery Dis* 2002;13:295-300.

Platz EA, Leitzmann MF, Michaud DS, Willett WC, Giovannucci E. Interrelation of Energy Intake, Body Size, and Physical Activity with Prostate Cancer in a Large Prospective Cohort Study. *Cancer Research* 2003;63:8542-8548.

Poddar KH, Hosig KW, Nickols-Richardson SM, Anderson ES, Herbert WG, Duncan SE. Low-fat dairy intake and body weight and composition changes in college students. *J Am Diet Assoc.* 2009;109:1433-8.

Posner BM. On the importance of dietary fat. Conclusions from the Framingham Study. *Acta Cardiol* 1993;48:452-5.

Posner BM, Cupples LA, Franz MM, Gagnon DR. Diet and heart disease risk factors in adult American men and women: the Framingham Offspring-Spouse nutrition studies. *Int J Epidemiol* 1993;22:1014-25.

Pott W, Albayrak O, Hebebrand J, Pauli-Pott U. Treating childhood obesity: family background variables and the child's success in a weight-control intervention. *Int J Eat Disord* 2009;42:284-9.

Potter JD, Steinmetz K. Vegetables, fruit and phytoestrogens as preventive agents. *IARC Sci Publ* 1996:61-90.

Provencher V, Begin C, Tremblay A, Mongeau L, Corneau L, Dodin S, et al. Health-At-Every-Size and eating behaviors: 1-year follow-up results of a size acceptance intervention. *J Am Diet Assoc* 2009;109:1854-61.

Psaltopoulou T, Kyrozi A, Stathopoulos P, Trichopoulos D, Vassilopoulos D, Trichopoulou A. Diet, physical activity and cognitive impairment among elders: the EPIC-Greece cohort (European Prospective Investigation into Cancer and Nutrition). *Public Health Nutr.* 2008;11:1054-62.

Purslow LR, Sandhu MS, Forouhi N, Young EH, Luben RN, Welch AA, et al. Energy intake at breakfast and weight change: prospective study of 6,764 middle-aged men and women. *Am J Epidemiol* 2008;167:188-92.

Qi L, Cornelis MC, Zhang C, van Dam RM, Hu FB. Genetic predisposition, Western dietary pattern, and the risk of type 2 diabetes in men. *Am J Clin Nutr* 2009;89:1453-8.

Qi L, van Dam RM, Rexrode K, Hu FB. Heme iron from diet as a risk factor for coronary heart disease in women with type 2 diabetes. *Diabetes Care* 2007;30:101-6.

Qin LQ, Xu JY, Wang PY, Hoshi K. Soyfood intake in the prevention of breast cancer risk in women: a meta-analysis of observational epidemiological studies. *J Nutr Sci Vitaminol (Tokyo)* 2006;52:428-36.

Randi G, Ferraroni M, Talamini R, Garavello W, Deandrea S, Decarli A, et al. Glycemic index, glycemic load and thyroid cancer risk. *Ann Oncol.* 2008;19:380-3.

Rapuri PB, Gallagher JC, Haynatzka V. Protein intake: effects on bone mineral density and the rate of bone loss in elderly women. *Am J Clin Nutr.* 2003;77:1517-25.

Rasanen M, Lehtinen JC, Niinikoski H, Keskinen S, Ruottinen S, Salminen M, et al. Dietary patterns and nutrient intakes of 7-year-old children taking part in an atherosclerosis prevention project in Finland. *J Am Diet Assoc* 2002;102:518-24.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Raynor HA, Jeffery RW, Tate DF, Wing RR. Relationship between changes in food group variety, dietary intake, and weight during obesity treatment. *Int J Obes Relat Metab Disord* 2004;28:813-20.

Raynor HA, Looney SM, Steeves EA, Spence M, Gorin AA. The effects of an energy density prescription on diet quality and weight loss: a pilot randomized controlled trial. *J Acad Nutr Diet* 2012;112:1397-402.

Raynor HA, Van Walleghen EL, Niemeier H, Butryn ML, Wing RR. Do food provisions packaged in single-servings reduce energy intake at breakfast during a brief behavioral weight-loss intervention? *J Am Diet Assoc* 2009;109:1922-5.

Razquin C, Alfredo Martinez J, Martinez-Gonzalez MA, Corella D, Santos JM, Marti A. The Mediterranean diet protects against waist circumference enlargement in 12Ala carriers for the PPARgamma gene: 2 years' follow-up of 774 subjects at high cardiovascular risk. *Br J Nutr*. 2009;102:672-9.

Razquin C, Martinez JA, Martinez-Gonzalez MA, Mitjavila MT, Estruch R, Marti A. A 3 years follow-up of a Mediterranean diet rich in virgin olive oil is associated with high plasma antioxidant capacity and reduced body weight gain. *Eur J Clin Nutr* 2009;63:1387-93.

Redman LM, Rood J, Anton SD, Champagne C, Smith SR, Ravussin E. Calorie restriction and bone health in young, overweight individuals. *Arch Intern Med* 2008;168:1859-66.

Reece EA, Hagay Z, Gay LJ, O'Connor T, DeGennaro N, Homko CJ, et al. A randomized clinical trial of a fiber-enriched diabetic diet vs. the standard American Diabetes Association-recommended diet in the management of diabetes mellitus in pregnancy. *Journal of Maternal-Fetal Investigation* 1995;5:8-12.

Reid V, Mulcahy R. Nutrient intakes and dietary compliance in cardiac patients: 6-year follow-up. *Hum Nutr Appl Nutr*. 1987;41:311-8.

Reinehr T, Kleber M, Toschke AM. Lifestyle intervention in obese children is associated with a decrease of the metabolic syndrome prevalence. *Atherosclerosis* 2009;174-80.

Reinehr T, Roth CL, Alexy U, Kersting M, Kiess W, Andler W. Ghrelin levels before and after reduction of overweight due to a low-fat high-carbohydrate diet in obese children and adolescents. *Int J Obes (Lond)* 2005;29:362-8.

Reinehr T, Schmidt C, Toschke AM, Andler W. Lifestyle intervention in obese children with nonalcoholic fatty liver disease: 2-Year follow-up study. *Archives of Disease in Childhood* 2009;94:437-442.

Reinehr T, Temmesfeld M, Kersting M, de Sousa G, Toschke AM. Four-year follow-up of children and adolescents participating in an obesity intervention program. *Int J Obes (Lond)* 2007;31:1074-7.

Remmell PS, Casey MP, McGandy RB, Stare FJ. A dietary program to lower serum cholesterol. *J Am Diet Assoc* 1969;54:13-9.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Renaud S, de Lorgeril M, Delaye J, Guidollet J, Jacquard F, Mamelle N, et al. Cretan Mediterranean diet for prevention of coronary heart disease. *Am J Clin Nutr* 1995;61:1360S-1367S.

Renner E. Dairy calcium, bone metabolism, and prevention of osteoporosis. *J Dairy Sci*. 1994;77:3498-505.

Reunanen A, Takkunen H, Knekt P, Seppanen R, Aromaa A. Body iron stores, dietary iron intake and coronary heart disease mortality. *J Intern Med* 1995;238:223-30.

Reynolds KD, Raczynski JM, Binkley D, Franklin FA, Duvall RC, Devane-Hart K, et al. Design of "High 5": a school-based study to promote fruit and vegetable consumption for reduction of cancer risk. *J Cancer Educ* 1998;13:169-77.

Rickman F, Mitchell N, Dingman J, Dalen JE. Changes in serum cholesterol during the Stillman diet. *JAMA*. 1974;228:54-8.

Riebe D, Blissmer B, Greene G, Caldwell M, Ruggiero L, Stillwell KM, et al. Long-term maintenance of exercise and healthy eating behaviors in overweight adults. *Prev Med* 2005;40:769-78.

Riedt CS, Schlussek Y, von Thun N, Ambia-Sobhan H, Stahl T, Field MP, et al. Premenopausal overweight women do not lose bone during moderate weight loss with adequate or higher calcium intake. *Am J Clin Nutr* 2007;85:972-80.

Rimm EB, Ascherio A, Giovannucci E, Spiegelman D, Stampfer MJ, Willett WC. Vegetable, fruit, and cereal fiber intake and risk of coronary heart disease among men. *JAMA* 1996;275:447-51.

Rimm EB, Katan MB, Ascherio A, Stampfer MJ, Willett WC. Relation between intake of flavonoids and risk for coronary heart disease in male health professionals. *Ann Intern Med* 1996;125:384-9.

Rissanen TH, Voutilainen S, Virtanen JK, Venho B, Vanharanta M, Mursu J, et al. Low intake of fruits, berries and vegetables is associated with excess mortality in men: the Kuopio Ischaemic Heart Disease Risk Factor (KIHD) Study. *J Nutr* 2003;133:199-204.

Robinson AH, Norman GJ, Sallis JF, Calfas KJ, Rock CL, Patrick K. Validating stage of change measures for physical activity and dietary behaviors for overweight women. *Int J Obes (Lond)* 2008;32:1137-44.

Robinson-O'Brien R, Perry CL, Wall MM, Story M, Neumark-Sztainer D. Adolescent and young adult vegetarianism: better dietary intake and weight outcomes but increased risk of disordered eating behaviors. *J Am Diet Assoc* 2009;109:648-55.

Roderick P. A randomized trial to evaluate the effectiveness of dietary advice by practice nurses in lowering diet-related coronary heart disease risk. *British Journal of General Practice* 1997;47:7-12.

Rogers IS, Emmett PM. Fat content of the diet among preschool children in southwest Britain: II. relationship with growth, blood lipids, and iron status. *Pediatrics*. 2001;108:E49.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Rohan TE, Howe GR, Friedenreich CM, Jain M, Miller AB. Dietary fiber, vitamins A, C, and E, and risk of breast cancer: a cohort study. *Cancer Causes Control*. 1993;4:29-37.

Rohan TE, Jain M, Howe GR, Miller AB. A cohort study of dietary carotenoids and lung cancer risk in women (Canada). *Cancer Causes Control* 2002;13:231-7.

Rolland-Cachera MF, Deheeger M, Akrouf M, Bellisle F. Influence of macronutrients on adiposity development: a follow up study of nutrition and growth from 10 months to 8 years of age. *Int J Obes Relat Metab Disord* 1995;19:573-8.

Rolls BJ, Roe LS, Beach AM, Kris-Etherton PM. Provision of foods differing in energy density affects long-term weight loss. *Obes Res* 2005;13:1052-60.

Romaguera D, Norat T, Mouw T, May AM, Bamia C, Slimani N, et al. Adherence to the Mediterranean diet is associated with lower abdominal adiposity in European men and women. *J Nutr* 2009;139:1728-37.

Romaguera D, Norat T, Vergnaud AC, Mouw T, May AM, Agudo A, et al. Mediterranean dietary patterns and prospective weight change in participants of the EPIC-PANACEA project. *Am J Clin Nutr* 2010;92:912-21.

Rosell M, Appleby P, Spencer E, Key T. Weight gain over 5 years in 21,966 meat-eating, fish-eating, vegetarian, and vegan men and women in EPIC-Oxford. *Int J Obes (Lond)* 2006;30:1389-96.

Rosell M, Hakansson NN, Wolk A. Association between dairy food consumption and weight change over 9 y in 19,352 perimenopausal women. *Am J Clin Nutr* 2006;84:1481-8.

Rosenheck R. Fast food consumption and increased caloric intake: a systematic review of a trajectory towards weight gain and obesity risk. *Obes Rev* 2008;9:535-47.

Roswall N, Olsen A, Christensen J, Dragsted LO, Overvad K, Tjønneland A. Micronutrient intake and breast cancer characteristics among postmenopausal women. *Eur J Cancer Prev*. 2010;19:360-5.

Roswall N, Olsen A, Christensen J, Dragsted LO, Overvad K, Tjønneland A. Micronutrient intake and risk of colon and rectal cancer in a Danish cohort. *Cancer Epidemiol*. 2010;34:40-6.

Rylander R, Axelsson G. Lung cancer risks in relation to vegetable and fruit consumption and smoking. *Int J Cancer* 2006;118:739-43.

Sabate J. Does nut consumption protect against ischaemic heart disease? *Eur J Clin Nutr* 1993;47 Suppl 1:S71-5.

Sabate J. Nut consumption and body weight. *Am J Clin Nutr* 2003;78:647S-650S.

Sabate J, Cordero-Macintyre Z, Siapco G, Torabian S, Haddad E. Does regular walnut consumption lead to weight gain? *Br J Nutr* 2005;94:859-64.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Sacks FM, Bray GA, Carey VJ, Smith SR, Ryan DH, Anton SD, et al. The new England journal of medicine: Comparison of weight-loss diets with different compositions of fat, protein, and carbohydrates. *New England Journal of Medicine* 2009;360:859-873.

Saelens BE, Couch SC, Wosje KS, Stark LJ, Daniels SR. Relations among milk and non-milk beverage consumption, calcium, and relative weight in high-weight status children. *Journal of Clinical Psychology in Medical Settings* 2006;13:117-125.

Sahay TB, Ashbury FD, Roberts M, Rootman I. Effective Components for Nutrition Interventions: A Review and Application of the Literature. *Health Promotion Practice* 2006;7:418-427.

Sahni S, Cupples LA, McLean RR, Tucker KL, Broe KE, Kiel DP, et al. Protective effect of high protein and calcium intake on the risk of hip fracture in the Framingham offspring cohort. *J Bone Miner Res* 2010;25:2494-500.

Sahyoun NR, Anderson AL, Tylavsky FA, Lee JS, Sellmeyer DE, Harris TB. Dietary glycemic index and glycemic load and the risk of type 2 diabetes in older adults. *Am J Clin Nutr* 2008;87:126-31.

Sahyoun NR, Jacques PF, Zhang XL, Juan W, McKeown NM. Whole-grain intake is inversely associated with the metabolic syndrome and mortality in older adults. *Am J Clin Nutr* 2006;83:124-31.

Salas-Salvado J, Bullo M, Babio N, Martinez-Gonzalez MA, Ibarrola-Jurado N, Basora J, et al. Reduction in the incidence of type 2 diabetes with the mediterranean diet: Results of the PREDIMED-Reus nutrition intervention randomized trial. *Diabetes Care* 2011;34:14-19.

Salas-Salvado J, Fernandez-Ballart J, Ros E, Martinez-Gonzalez MA, Fito M, Estruch R, et al. Effect of a Mediterranean diet supplemented with nuts on metabolic syndrome status: one-year results of the PREDIMED randomized trial. *Arch Intern Med* 2008;168:2449-58.

Salmeron J, Ascherio A, Rimm EB, Colditz GA, Spiegelman D, Jenkins DJ, et al. Dietary fiber, glycemic load, and risk of NIDDM in men. *Diabetes Care* 1997;20:545-50.

Salmeron J, Hu FB, Manson JE, Stampfer MJ, Colditz GA, Rimm EB, et al. Dietary fat intake and risk of type 2 diabetes in women. *Am J Clin Nutr*. 2001;73:1019-26.

Salmeron J, Manson JE, Stampfer MJ, Colditz GA, Wing AL, Willett WC. Dietary fiber, glycemic load, and risk of non-insulin-dependent diabetes mellitus in women. *JAMA* 1997;277:472-7.

Samaha FF, Foster GD, Makris AP. Low-carbohydrate diets, obesity, and metabolic risk factors for cardiovascular disease. *Current Atherosclerosis Reports* 2007;9:441-447.

Sanchez-Villegas A, Bes-Rastrollo M, Martinez-Gonzalez MA, Serra-Majem L. Adherence to a Mediterranean dietary pattern and weight gain in a follow-up study: the SUN cohort. *Int J Obes (Lond)*. 2006;30:350-8.

Sanchez-Villegas A, Delgado-Rodriguez M, Alonso A, Schlatter J, Lahortiga F, Serra Majem L, et al. Association of the Mediterranean dietary pattern with the incidence of depression: the Seguimiento Universidad de Navarra/University of Navarra follow-up (SUN) cohort. *Arch Gen Psychiatry*. 2009;66:1090-8.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Sansbury LB, Wanke K, Albert PS, Kahle L, Schatzkin A, Lanza E. The effect of strict adherence to a high-fiber, high-fruit and -vegetable, and low-fat eating pattern on adenoma recurrence. *Am J Epidemiol* 2009;170:576-84.

Sargeant LA, Wareham NJ, Bingham S, Day NE, Luben RN, Oakes S, et al. Vitamin C and hyperglycemia in the European Prospective Investigation into Cancer--Norfolk (EPIC-Norfolk) study: a population-based study. *Diabetes Care* 2000;23:726-32.

Saris WHM, Astrup A, Prentice AM, Zunft HJF, Formiguera X, Verboeket-Van De Venne WPHG, et al. Randomized controlled trial of changes in dietary carbohydrate/fat ratio and simple vs complex carbohydrates on body weight and blood lipids: The CARMEN study. *International Journal of Obesity* 2000;24:1310-1318.

Sarter B, Campbell TC, Fuhrman J. Effect of a high nutrient density diet on long-term weight loss: a retrospective chart review. *Altern Ther Health Med* 2008;14:48-53.

Sarwer DB, Wadden TA, Moore RH, Baker AW, Gibbons LM, Raper SE, et al. Preoperative eating behavior, postoperative dietary adherence, and weight loss after gastric bypass surgery. *Surg Obes Relat Dis* 2008;4:640-6.

Sasaki S, Ishikawa T, Yanagibori R, Amano K. Change and 1-year maintenance of nutrient and food group intakes at a 12-week worksite dietary intervention trial for men at high risk of coronary heart disease. *J Nutr Sci Vitaminol (Tokyo)* 2000;46:15-22.

Sato Y, Tsubono Y, Nakaya N, Ogawa K, Kurashima K, Kuriyama S, et al. Fruit and vegetable consumption and risk of colorectal cancer in Japan: The Miyagi Cohort Study. *Public Health Nutr* 2005;8:309-14.

Sauvaget C, Nagano J, Allen N, Grant EJ, Beral V. Intake of animal products and stroke mortality in the Hiroshima/Nagasaki Life Span Study. *Int J Epidemiol* 2003;32:536-43.

Sauvaget C, Nagano J, Allen N, Kodama K. Vegetable and fruit intake and stroke mortality in the Hiroshima/Nagasaki Life Span Study. *Stroke* 2003;34:2355-60.

Sauvaget C, Nagano J, Hayashi M, Spencer E, Shimizu Y, Allen N. Vegetables and fruit intake and cancer mortality in the Hiroshima/Nagasaki Life Span Study. *Br J Cancer* 2003;88:689-94.

Sauvaget C, Nagano J, Hayashi M, Yamada M. Animal protein, animal fat, and cholesterol intakes and risk of cerebral infarction mortality in the adult health study. *Stroke* 2004;35:1531-7.

Savage JS, Birch LL. Patterns of weight control strategies predict differences in women's 4-year weight gain. *Obesity (Silver Spring)* 2010;18:513-20.

Savage JS, Marini M, Birch LL. Dietary energy density predicts women's weight change over 6 y. *Am J Clin Nutr* 2008;88:677-84.

Savoie M, Nowicka P, Shaw M, Yu S, Dziura J, Chavent G, et al. Long-term results of an obesity program in an ethnically diverse pediatric population. *Pediatrics* 2011;127:402-10.

Scarmeas N, Luchsinger JA, Mayeux R, Schupf N, Stern Y. Mediterranean diet and mortality in a US population. *Arch Intern Med* 2008;168:1823-4.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Scarmeas N, Luchsinger JA, Mayeux R, Stern Y. Mediterranean diet and Alzheimer disease mortality. *Neurology*. 2007;69:1084-93.

Scarmeas N, Luchsinger JA, Schupf N, Brickman AM, Cosentino S, Tang MX, et al. Physical activity, diet, and risk of Alzheimer disease. *JAMA*. 2009;302:627-37.

Scarmeas N, Stern Y, Mayeux R, Manly JJ, Schupf N, Luchsinger JA. Mediterranean diet and mild cognitive impairment. *Arch Neurol*. 2009;66:216-25.

Scavone G, Manto A, Pitocco D, Gagliardi L, Caputo S, Mancini L, et al. Effect of carbohydrate counting and medical nutritional therapy on glycaemic control in Type 1 diabetic subjects: A pilot study. *Diabetic Medicine* 2010;27:477-479.

Schatzkin A. Going against the grain? Current status of the dietary fiber-colorectal cancer hypothesis. *Biofactors* 2000;12:305-11.

Schatzkin A, Lanza E. Polyps and vegetables (and fat, fibre): the polyp prevention trial. *IARC Sci Publ* 2002;156:463-6.

Schatzkin A, Park Y, Leitzmann MF, Hollenbeck AR, Cross AJ. Prospective study of dietary fiber, whole grain foods, and small intestinal cancer. *Gastroenterology* 2008;135:1163-7.

Schatzkin A, Subar AF, Thompson FE, Harlan LC, Tangrea J, Hollenbeck AR, et al. Design and serendipity in establishing a large cohort with wide dietary intake distributions : the National Institutes of Health-American Association of Retired Persons Diet and Health Study. *Am J Epidemiol* 2001;154:1119-25.

Schiepers OJ, de Groot RH, Jolles J, van Boxtel MP. Fish consumption, not fatty acid status, is related to quality of life in a healthy population. *Prostaglandins Leukot Essent Fatty Acids* 2010;83:31-5.

Scholtens S, Brunekreef B, Smit HA, Gast GC, Hoekstra MO, de Jongste JC, et al. Do differences in childhood diet explain the reduced overweight risk in breastfed children? *Obesity (Silver Spring)* 2008;16:2498-503.

Schroder KE. Effects of fruit consumption on body mass index and weight loss in a sample of overweight and obese dieters enrolled in a weight-loss intervention trial. *Nutrition* 2009;26:727-34.

Schroder KE. Effects of fruit consumption on body mass index and weight loss in a sample of overweight and obese dieters enrolled in a weight-loss intervention trial. *Nutrition* 2010;26:727-34.

Schulz M, Hoffmann K, Weikert C, Nothlings U, Schulze MB, Boeing H. Identification of a dietary pattern characterized by high-fat food choices associated with increased risk of breast cancer: The European Prospective Investigation into Cancer and Nutrition (EPIC)-Potsdam Study. *British Journal of Nutrition* 2008;100:942-946.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Schulz M, Kroke A, Liese AD, Hoffmann K, Bergmann MM, Boeing H. Food groups as predictors for short-term weight changes in men and women of the EPIC-Potsdam cohort. *J Nutr* 2002;132:1335-40.

Schulz M, Lahmann PH, Boeing H, Hoffmann K, Allen N, Key TJ, et al. Fruit and vegetable consumption and risk of epithelial ovarian cancer: the European Prospective Investigation into Cancer and Nutrition. *Cancer Epidemiol Biomarkers Prev* 2005;14:2531-5.

Schulz M, Nothlings U, Hoffmann K, Bergmann MM, Boeing H. Identification of a food pattern characterized by high-fiber and low-fat food choices associated with low prospective weight change in the EPIC-Potsdam cohort. *J Nutr* 2005;135:1183-9.

Schulze MB, Fung TT, Manson JE, Willett WC, Hu FB. Dietary patterns and changes in body weight in women. *Obesity (Silver Spring)* 2006;14:1444-53.

Schulze MB, Hoffmann K, Manson JE, Willett WC, Meigs JB, Weikert C, et al. Dietary pattern, inflammation, and incidence of type 2 diabetes in women. *Am J Clin Nutr* 2005;82:675-84; quiz 714-5.

Schulze MB, Schulz M, Heidemann C, Schienkiewitz A, Hoffmann K, Boeing H. Fiber and magnesium intake and incidence of type 2 diabetes: a prospective study and meta-analysis. *Arch Intern Med* 2007;167:956-65.

Scrafford CG, Tran NL, Barraj LM, Mink PJ. Egg consumption and CHD and stroke mortality: a prospective study of US adults. *Public Health Nutr* 2011;14:261-70.

Sellers TA, Alberts SR, Vierkant RA, Grabrick DM, Cerhan JR, Vachon CM, et al. High-folate diets and breast cancer survival in a prospective cohort study. *Nutr Cancer* 2002;44:139-44.

Sellmeyer DE, Stone KL, Sebastian A, Cummings SR. A high ratio of dietary animal to vegetable protein increases the rate of bone loss and the risk of fracture in postmenopausal women. Study of Osteoporotic Fractures Research Group. *Am J Clin Nutr* 2001;73:118-22.

Sengupta S, Tjandra JJ, Gibson PR. Dietary fiber and colorectal neoplasia. *Dis Colon Rectum* 2001;44:1016-33.

Serra-Majem L, La Vecchia C, Ribas-Barba L, Prieto-Ramos F, Lucchini F, Ramon JM, et al. Changes in diet and mortality from selected cancers in southern Mediterranean countries, 1960-1989. *Eur J Clin Nutr* 1993;47 Suppl 1:S25-34.

Seshadri P, Samaha FF, Stern L, Ahima RS, Daily D, Iqbal N. Adipocytokine changes caused by low-carbohydrate compared to conventional diets in obesity. *Metabolic Syndrome and Related Disorders* 2005;3:66-74.

Sesso HD, Gaziano JM, Jenkins DJ, Buring JE. Strawberry intake, lipids, C-reactive protein, and the risk of cardiovascular disease in women. *J Am Coll Nutr* 2007;26:303-10.

Sesso HD, Liu S, Gaziano JM, Buring JE. Dietary lycopene, tomato-based food products and cardiovascular disease in women. *J Nutr* 2003;133:2336-41.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Seymour JD, Calle EE, Flagg EW, Coates RJ, Ford ES, Thun MJ. Diet Quality Index as a predictor of short-term mortality in the American Cancer Society Cancer Prevention Study II Nutrition Cohort. *Am J Epidemiol* 2003;157:980-8.

Shah M, Adams-Huet B, Garg A. Effect of high-carbohydrate or high-cis-monounsaturated fat diets on blood pressure: A meta-analysis of intervention trials. *American Journal of Clinical Nutrition* 2007;85:1251-1256.

Shahar DR, Schwarzfuchs D, Fraser D, Vardi H, Thiery J, Fiedler GM, et al. Dairy calcium intake, serum vitamin D, and successful weight loss. *Am J Clin Nutr* 2010;92:1017-22.

Shaw DI, Tierney AC, McCarthy S, Upritchard J, Vermunt S, Gulseth HL, et al. LIPGENE food-exchange model for alteration of dietary fat quantity and quality in free-living participants from eight European countries. *British Journal of Nutrition* 2009;101:750-759.

Shekelle RB, Shryock AM, Paul O, Lepper M, Stamler J, Liu S, et al. Diet, serum cholesterol, and death from coronary heart disease. The Western Electric study. *N Engl J Med* 1981;304:65-70.

Shekelle RB, Stamler J. Dietary cholesterol and ischaemic heart disease. *Lancet* 1989;1:1177-9.

Sherafat-Kazemzadeh R, Egtesadi S, Mirmiran P, Gohari M, Farahani SJ, Esfahani FH, et al. Dietary patterns by reduced rank regression predicting changes in obesity indices in a cohort study: Tehran Lipid and Glucose Study. *Asia Pac J Clin Nutr* 2010;19:22-32.

Shi Z, Taylor AW, Hu G, Gill T, Wittert GA. Rice intake, weight change and risk of the metabolic syndrome development among Chinese adults: the Jiangsu Nutrition Study (JIN). *Asia Pac J Clin Nutr* 2012;21:35-43.

Shi Z, Yuan B, Hu G, Dai Y, Zuo H, Holmboe-Ottesen G. Dietary pattern and weight change in a 5-year follow-up among Chinese adults: results from the Jiangsu Nutrition Study. *Br J Nutr* 2011;105:1047-54.

Shimazu T, Inoue M, Sasazuki S, Iwasaki M, Sawada N, Yamaji T, et al. Isoflavone intake and risk of lung cancer: a prospective cohort study in Japan. *Am J Clin Nutr* 2010;91:722-8.

Shin A, Li H, Shu XO, Yang G, Gao YT, Zheng W. Dietary intake of calcium, fiber and other micronutrients in relation to colorectal cancer risk: Results from the Shanghai Women's Health Study. *Int J Cancer*. 2006;119:2938-42.

Shu XO, Zheng Y, Cai H, Gu K, Chen Z, Zheng W, et al. Soy food intake and breast cancer survival. *JAMA* 2009;302:2437-43.

Sidorenkov O, Nilssen O, Grjibovski AM. Determinants of cardiovascular and all-cause mortality in northwest Russia: a 10-year follow-up study. *Ann Epidemiol* 2012;22:57-65.

Siega-Riz AM, El Ghormli L, Mobley C, Gillis B, Stadler D, Hartstein J, et al. The effects of the HEALTHY study intervention on middle school student dietary intakes. *SOURCE International Journal of Behavioral Nutrition and Physical Activity* (2011) 8 Article Number: 7. Date of Publication: 4 Feb 2011 2011.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Siegel EM, Salemi JL, Villa LL, Ferenczy A, Franco EL, Giuliano AR. Dietary consumption of antioxidant nutrients and risk of incident cervical intraepithelial neoplasia. *Gynecol Oncol* 2010;118:289-94.

Siegel RM, Rich W, Joseph EC, Linhardt J, Knight J, Khoury J, et al. A 6-month, office-based, low-carbohydrate diet intervention in obese teens. *Clinical Pediatrics* 2009;48:745-749.

Sieri S, Krogh V, Pala V, Muti P, Micheli A, Evangelista A, et al. Dietary patterns and risk of breast cancer in the ORDET cohort. *Cancer Epidemiol Biomarkers Prev* 2004;13:567-72.

Sieri S, Pala V, Brighenti F, Pellegrini N, Muti P, Micheli A, et al. Dietary glycemic index, glycemic load, and the risk of breast cancer in an Italian prospective cohort study. *Am J Clin Nutr*. 2007;86:1160-6.

Silvera SA, Jain M, Howe GR, Miller AB, Rohan TE. Dietary carbohydrates and breast cancer risk: a prospective study of the roles of overall glycemic index and glycemic load. *Int J Cancer*. 2005;114:653-8.

Silvera SA, Jain M, Howe GR, Miller AB, Rohan TE. Energy balance and breast cancer risk: a prospective cohort study. *Breast Cancer Res Treat*. 2006;97:97-106.

Silvera SA, Jain M, Howe GR, Miller AB, Rohan TE. Dietary fiber intake and ovarian cancer risk: a prospective cohort study. *Cancer Causes Control* 2007;18:335-41.

Simell O, Niinikoski H, Ronnema T, Lapinleimu H, Routi T, Lagstrom H, et al. Special Turku Coronary Risk Factor Intervention Project for Babies (STRIP). *Am J Clin Nutr* 2000;72:1316S-1331S.

Simila ME, Valsta LM, Kontto JP, Albanes D, Virtamo J. Low-, medium- and high-glycaemic index carbohydrates and risk of type 2 diabetes in men. *Br J Nutr*. 2011;105:1258-64.

Simons CC, Hughes LA, Arts IC, Goldbohm RA, van den Brandt PA, Weijenberg MP. Dietary flavonol, flavone and catechin intake and risk of colorectal cancer in the Netherlands Cohort Study. *Int J Cancer*. 2009;125:2945-52.

Simonsen T, Nordoy A. Ischaemic heart disease, serum lipids and platelets in Norwegian populations with traditionally low or high fish consumption. *J Intern Med Suppl* 1989;731:83-9.

Sinclair AJ, Burdon MA, Nightingale PG, Ball AK, Good P, Matthews TD, et al. Low energy diet and intracranial pressure in women with idiopathic intracranial hypertension: Prospective cohort study. *BMJ* 2010;341:138.

Singh AS, Chinapaw MJM, Brug J, van Mechelen W. Process evaluation of a school-based weight gain prevention program: The Dutch Obesity Intervention in Teenagers (DOiT). *Health Education Research* 2009;24:772-777.

Singh PN, Fraser GE. Dietary risk factors for colon cancer in a low-risk population. *Am J Epidemiol*. 1998;148:761-74.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Singh RB, Niaz MA, Bishnoi I, Sharma JP, Gupta S, Rastogi SS, et al. Diet, antioxidant vitamins, oxidative stress and risk of coronary artery disease: the Peerzada Prospective Study. *Acta Cardiol.* 1994;49:453-67.

Singh RB, Niaz MA, Ghosh S. Effect on central obesity and associated disturbances of low-energy, fruit- and vegetable-enriched prudent diet in north Indians. *Postgrad Med J* 1994;70:895-900.

Singh RB, Rastogi SS, Verma R, Laxmi B, Singh R, Ghosh S, et al. Randomised controlled trial of cardioprotective diet in patients with recent acute myocardial infarction: results of one year follow up. *BMJ* 1992;304:1015-9.

Singh RB, Rastogi V, Rastogi SS, Niaz MA, Beegom R. Effect of diet and moderate exercise on central obesity and associated disturbances, myocardial infarction and mortality in patients with and without coronary artery disease. *Journal of the American College of Nutrition* 1996;15:592-601.

Singh RB, Singh NK, Rastogi SS, Mani UV, Niaz MA. Effects of diet and lifestyle changes on atherosclerotic risk factors after 24 weeks on the Indian Diet Heart Study. *Am J Cardiol* 1993;71:1283-8.

Sinha R, Cross AJ, Graubard BI, Leitzmann MF, Schatzkin A. Meat intake and mortality: a prospective study of over half a million people. *Arch Intern Med* 2009;169:562-71.

Skinner JD, Bounds W, Carruth BR, Ziegler P. Longitudinal calcium intake is negatively related to children's body fat indexes. *J Am Diet Assoc* 2003;103:1626-31.

Skuladottir H, Tjoenneland A, Overvad K, Stripp C, Christensen J, Raaschou-Nielsen O, et al. Does insufficient adjustment for smoking explain the preventive effects of fruit and vegetables on lung cancer? *Lung Cancer* 2004;45:1-10.

Slemenda CW, Christian JC, Reed T, Reister TK, Williams CJ, Johnston CC, Jr. Long-term bone loss in men: effects of genetic and environmental factors. *Ann Intern Med.* 1992;117:286-91.

Slimko ML, Mensah GA. The role of diets, food, and nutrients in the prevention and control of hypertension and prehypertension. *Cardiol Clin* 2010;28:665-74.

Sluijs I, Beulens JW, van der AD, Spijkerman AM, Grobbee DE, van der Schouw YT. Dietary intake of total, animal, and vegetable protein and risk of type 2 diabetes in the European Prospective Investigation into Cancer and Nutrition (EPIC)-NL study. *Diabetes Care* 2010;33:43-8.

Sluijs I, van der Schouw YT, van der AD, Spijkerman AM, Hu FB, Grobbee DE, et al. Carbohydrate quantity and quality and risk of type 2 diabetes in the European Prospective Investigation into Cancer and Nutrition-Netherlands (EPIC-NL) study. *Am J Clin Nutr* 2010;92:905-11.

Smedslund G, Byfuglien MG, Olsen SU, Birger Hagen K. Effectiveness and safety of dietary interventions for rheumatoid arthritis: a systematic review of randomized controlled trials (Brief record). *Journal of the American Dietetic Association* 2010:727-735.

Smit E, Garcia-Palmieri MR, Figueroa NR, McGee DL, Messina M, Freudenheim JL, et al. Protein and legume intake and prostate cancer mortality in Puerto Rican men. *Nutr Cancer* 2007;58:146-52.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Smith-Warner SA, Elmer PJ, Tharp TM, Fosdick L, Randall B, Gross M, et al. Increasing vegetable and fruit intake: Randomized intervention and monitoring in an at-risk population. *Cancer Epidemiology Biomarkers and Prevention* 2000;9:307-317.

Snowdon DA. Diet and ovarian cancer. *JAMA* 1985;254:356-7.

Snyder MP, Anliker J, Cunningham-Sabo L, Dixon LB, Altaba J, Chamberlain A, et al. The pathways study: A model for lowering the fat in school meals. *American Journal of Clinical Nutrition* 1999;69:810S-815S.

Sofi F, Cesari F, Abbate R, Gensini GF, Casini A. Adherence to Mediterranean diet and health status: meta-analysis. *BMJ* 2008;337:a1344.

Soinio M, Laakso M, Lehto S, Hakala P, Ronnema T. Dietary fat predicts coronary heart disease events in subjects with type 2 diabetes. *Diabetes Care* 2003;26:619-24.

Solfrizzi V, D'Introno A, Colacicco AM, Capurso C, Palasciano R, Capurso S, et al. Unsaturated fatty acids intake and all-causes mortality: a 8.5-year follow-up of the Italian Longitudinal Study on Aging. *Exp Gerontol* 2005;40:335-43.

Song EK. Adherence to the low-sodium diet plays a role in the interaction between depressive symptoms and prognosis in patients with heart failure. *J Cardiovasc Nurs.* 2009;24:299-305; quiz 306-7.

Song Y, Manson JE, Buring JE, Liu S. A prospective study of red meat consumption and type 2 diabetes in middle-aged and elderly women: the women's health study. *Diabetes Care* 2004;27:2108-15.

Song Y, Manson JE, Buring JE, Liu S. Dietary Magnesium Intake in Relation to Plasma Insulin Levels and Risk of Type 2 Diabetes in Women. *Diabetes Care* 2004;27:59-65.

Song Y, Manson JE, Buring JE, Sesso HD, Liu S. Associations of dietary flavonoids with risk of type 2 diabetes, and markers of insulin resistance and systemic inflammation in women: a prospective study and cross-sectional analysis. *J Am Coll Nutr* 2005;24:376-84.

Song Y, Sesso HD, Manson JE, Cook NR, Buring JE, Liu S. Dietary magnesium intake and risk of incident hypertension among middle-aged and older US women in a 10-year follow-up study. *Am J Cardiol* 2006;98:1616-21.

Sonnenberg L, Pencina M, Kimokoti R, Quatromoni P, Nam BH, D'Agostino R, et al. Dietary patterns and the metabolic syndrome in obese and non-obese Framingham women. *Obes Res* 2005;13:153-62.

Soriguer F, Almaraz MC, Garcia-Almeida JM, Cardona I, Linares F, Morcillo S, et al. Intake and home use of olive oil or mixed oils in relation to healthy lifestyles in a Mediterranean population. Findings from the prospective Pizarra study. *Br J Nutr.* 2010;103:114-22.

Sothorn MS, von Almen TK, Schumacher H, Zelman M, Farris RP, Carlisle L, et al. An effective multidisciplinary approach to weight reduction in youth. *Ann N Y Acad Sci* 1993;699:292-4.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Spector D, Anthony M, Alexander D, Arab L. Soy consumption and colorectal cancer. *Nutr Cancer* 2003;47:1-12.

Spencer EA, Appleby PN, Davey GK, Key TJ. Diet and body mass index in 38000 EPIC-Oxford meat-eaters, fish-eaters, vegetarians and vegans. *Int J Obes Relat Metab Disord.* 2003;27:728-34.

Sriamporn S, Parkin DM, Pisani P, Vatanasapt V, Suwanrungruang K, Kamsa-ard P, et al. A prospective study of diet, lifestyle, and genetic factors and the risk of cancer in Khon Kaen Province, northeast Thailand: description of the cohort. *Asian Pac J Cancer Prev.* 2005;6:295-303.

Stahelin HB, Gey F, Brubacher G. Preventive potential of antioxidative vitamins and carotenoids on cancer. *Int J Vitam Nutr Res Suppl* 1989;30:232-41.

Stahelin HB, Rosel F, Buess E, Brubacher G. Dietary risk factors for cancer in the Basel Study. *Bibl Nutr Dieta.* 1986:144-53.

Stamler J, Dolecek TA, Grandits GA. Chapter 13. Relation of food and nutrient intakes to body mass in the special intervention and usual care groups in the Multiple Risk Factor Intervention Trial. *American Journal of Clinical Nutrition* 1997;65:366S-373S.

Stamler R, Stamler J, Grimm R. Nutritional therapy for high blood pressure. Final report of a four-year randomized controlled trial - The Hypertension Control Program. *Journal of the American Medical Association* 1987;257:1484-1491.

Stamp LK, James MJ, Cleland LG. Diet and rheumatoid arthritis: a review of the literature. *Semin Arthritis Rheum.* 2005;35:77-94.

Stampfer MJ, Hu FB, Manson JE, Rimm EB, Willett WC. Primary prevention of coronary heart disease in women through diet and lifestyle. *N Engl J Med* 2000;343:16-22.

Steevens J, Schouten LJ, Goldbohm RA, van den Brandt PA. Vegetables and fruits consumption and risk of esophageal and gastric cancer subtypes in the Netherlands Cohort Study. *Int J Cancer* 2011;129:2681-93.

Steffen LM, Jacobs DR, Jr., Murtaugh MA, Moran A, Steinberger J, Hong CP, et al. Whole grain intake is associated with lower body mass and greater insulin sensitivity among adolescents. *Am J Epidemiol.* 2003;158:243-50.

Steinmetz KA, Potter JD. Vegetables, fruit, and cancer prevention: a review. *J Am Diet Assoc* 1996;96:1027-39.

Stellman SD, Garfinkel L. Artificial sweetener use and one-year weight change among women. *Prev Med.* 1986;15:195-202.

Stenius-Aarniala B, Poussa T, Kvarnstrom J, Gronlund EL, Ylikahri M, Mustajoki P. Immediate and long term effects of weight reduction in obese people with asthma: randomised controlled study. *BMJ* 2000;320:827-32.

Steyn NP, Lambert EV, Tabana H. Conference on "Multidisciplinary approaches to nutritional problems". Symposium on "Diabetes and health". Nutrition interventions for the prevention of type 2 diabetes. *Proc Nutr Soc.* 2009;68:55-70.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Stolzenberg-Solomon RZ, Pietinen P, Barrett MJ, Taylor PR, Virtamo J, Albanes D. Dietary and other methyl-group availability factors and pancreatic cancer risk in a cohort of male smokers. *Am J Epidemiol* 2001;153:680-7.

Stracke H, Renner E, Knie G, Leidig G, Minne H, Federlin K. Osteoporosis and bone metabolic parameters in dependence upon calcium intake through milk and milk products. *Eur J Clin Nutr*. 1993;47:617-22.

Stram DO, Hankin JH, Wilkens LR, Park S, Henderson BE, Nomura AM, et al. Prostate cancer incidence and intake of fruits, vegetables and related micronutrients: the multiethnic cohort study* (United States). *Cancer Causes Control*. 2006;17:193-207.

Strandberg TE, Strandberg AY, Pitkala K, Salomaa VV, Tilvis RS, Miettinen TA. Chocolate, well-being and health among elderly men. *Eur J Clin Nutr* 2008;62:247-53.

Stranges S, Sieri S, Vinceti M, Grioni S, Guallar E, Laclaustra M, et al. A prospective study of dietary selenium intake and risk of type 2 diabetes. *BMC Public Health* 2010;10:564.

Streppel MT, Ocke MC, Boshuizen HC, Kok FJ, Kromhout D. Dietary fiber intake in relation to coronary heart disease and all-cause mortality over 40 y: the Zutphen Study. *Am J Clin Nutr* 2008;88:1119-25.

Streppel MT, Ocke MC, Boshuizen HC, Kok FJ, Kromhout D. Long-term fish consumption and n-3 fatty acid intake in relation to (sudden) coronary heart disease death: the Zutphen study. *Eur Heart J* 2008;29:2024-30.

Summerbell Carolyn D, Cameron C, Glasziou Paul P. Advice on low-fat diets for obesity. *Cochrane Database of Systematic Reviews* 2008.

Sun Q, Spiegelman D, van Dam RM, Holmes MD, Malik VS, Willett WC, et al. White rice, brown rice, and risk of type 2 diabetes in US men and women. *Arch Intern Med* 2010;170:961-9.

Suzuki Y, Whiting SJ, Davison KS, Chilibeck PD. Total calcium intake is associated with cortical bone mineral density in a cohort of postmenopausal women not taking estrogen. *Journal of Nutrition, Health and Aging* 2003;7:296-299.

Sweeney ME, Hill JO, Heller PA, Baney R, DiGirolamo M. Severe vs moderate energy restriction with and without exercise in the treatment of obesity: efficiency of weight loss. *Am J Clin Nutr* 1993;57:127-34.

Svendsen M, Blomhoff R, Holme I, Tonstad S. The effect of an increased intake of vegetables and fruit on weight loss, blood pressure and antioxidant defense in subjects with sleep related breathing disorders. *Eur J Clin Nutr* 2007;61:1301-11.

Takachi R, Inoue M, Ishihara J, Kurahashi N, Iwasaki M, Sasazuki S, et al. Fruit and vegetable intake and risk of total cancer and cardiovascular disease: Japan Public Health Center-Based Prospective Study. *Am J Epidemiol* 2008;167:59-70.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Takashashi Y, Sasaki S, Takahashi M, Okubo S, Hayashi M, Tsugane S. A population-based dietary intervention trial in a high-risk area for stomach cancer and stroke: changes in intakes and related biomarkers. *Prev Med* 2003;37:432-41.

Talvia S, Lagstrom H, Rasanen M, Salminen M, Rasanen L, Salo P, et al. A randomized intervention since infancy to reduce intake of saturated fat: calorie (energy) and nutrient intakes up to the age of 10 years in the Special Turku Coronary Risk Factor Intervention Project. *Arch Pediatr Adolesc Med* 2004;158:41-7.

Tang L, Zirpoli GR, Guru K, Moysich KB, Zhang Y, Ambrosone CB, et al. Intake of cruciferous vegetables modifies bladder cancer survival. *Cancer Epidemiol Biomarkers Prev* 2010;19:1806-11.

Tanumihardjo SA, Valentine AR, Zhang Z, Whigham LD, Lai HJ, Atkinson RL. Strategies to increase vegetable or reduce energy and fat intake induce weight loss in adults. *Exp Biol Med (Maywood)* 2009;234:542-52.

Tapsell LC, Batterham MJ, Teuss G, Tan SY, Dalton S, Quick CJ, et al. Long-term effects of increased dietary polyunsaturated fat from walnuts on metabolic parameters in type II diabetes. *Eur J Clin Nutr* 2009;63:1008-15.

Tapsell LC, Gillen LJ, Patch CS, Batterham M, Owen A, Bare M, et al. Including walnuts in a low-fat/modified-fat diet improves HDL cholesterol-to-total cholesterol ratios in patients with type 2 diabetes. *Diabetes Care* 2004;27:2777-83.

Tasevska N, Sinha R, Kipnis V, Subar AF, Leitzmann MF, Hollenbeck AR, et al. A prospective study of meat, cooking methods, meat mutagens, heme iron, and lung cancer risks. *Am J Clin Nutr*. 2009;89:1884-94.

Tavani A, La Vecchia C. Fruit and vegetable consumption and cancer risk in a Mediterranean population. *Am J Clin Nutr* 1995;61:1374S-1377S.

Taylor FC, Irons LJ, Finn P, Summerbell CD. Controlled clinical trial of two weight reducing diets in a NHS hospital dietetic outpatient clinic - a pilot study. *J Hum Nutr Diet* 2003;16:85-7.

te Velde SJ, Twisk JW, Brug J. Tracking of fruit and vegetable consumption from adolescence into adulthood and its longitudinal association with overweight. *Br J Nutr* 2007;98:431-8.

Temple NJ, Gladwin KK. Fruit, vegetables, and the prevention of cancer: research challenges. *Nutrition* 2003;19:467-70.

ter Bogt NCW, Bemelmans WJE, Beltman FW, Broer J, Smit AJ, van der Meer K. Preventing weight gain: One-year results of a randomized lifestyle intervention. *American Journal of Preventive Medicine* 2009;37:270-277.

Terry P, Baron JA, Bergkvist L, Holmberg L, Wolk A. Dietary calcium and vitamin D intake and risk of colorectal cancer: a prospective cohort study in women. *Nutr Cancer* 2002;43:39-46.

Terry P, Giovannucci E, Michels KB, Bergkvist L, Hansen H, Holmberg L, et al. Fruit, vegetables, dietary fiber, and risk of colorectal cancer. *J Natl Cancer Inst* 2001;93:525-33.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Terry P, Jain M, Miller AB, Howe GR, Rohan TE. Dietary carotenoids and risk of breast cancer. *American Journal of Clinical Nutrition* 2002;76:883-888.

Terry P, Lichtenstein P, Feychting M, Ahlbom A, Wolk A. Fatty fish consumption and risk of prostate cancer. *Lancet* 2001;357:1764-6.

Terry P, Nyren O, Yuen J. Protective effect of fruits and vegetables on stomach cancer in a cohort of Swedish twins. *Int J Cancer* 1998;76:35-7.

Terry P, Wolk A, Vainio H, Weiderpass E. Fatty fish consumption lowers the risk of endometrial cancer: a nationwide case-control study in Sweden. *Cancer Epidemiol Biomarkers Prev* 2002;11:143-5.

Thiebaut AC, Chajes V, Gerber M, Boutron-Ruault MC, Joulin V, Lenoir G, et al. Dietary intakes of omega-6 and omega-3 polyunsaturated fatty acids and the risk of breast cancer. *Int J Cancer* 2009;124:924-31.

Thomas DE, Elliott EJ, Baur L. Low glycaemic index or low glycaemic load diets for overweight and obesity. *SOURCE Cochrane Database of Systematic Reviews (2007) :3 Article Number: CD005105.* Date of Publication: 2007 2007.

Thorogood M. Vegetarianism, coronary disease risk factors and coronary heart disease. *Curr Opin Lipidol* 1994;5:17-21.

Thorogood M, Mann J, Appleby P, McPherson K. Risk of death from cancer and ischaemic heart disease in meat and non-meat eaters. *BMJ* 1994;308:1667-70.

Thorpe DL, Knutsen SF, Beeson WL, Rajaram S, Fraser GE. Effects of meat consumption and vegetarian diet on risk of wrist fracture over 25 years in a cohort of peri- and postmenopausal women. *Public Health Nutr* 2008;11:564-72.

Thorsteinsson R, Johannesson A, Jonsson H, Thorhallsson T, Sigurdsson JA. Effects of dietary intervention on serum lipids in factory workers. *Scand J Prim Health Care* 1994;12:93-9.

Timonen M, Horrobin D, Jokelainen J, Laitinen J, Herva A, Rasanen P. Fish consumption and depression: the Northern Finland 1966 birth cohort study. *J Affect Disord* 2004;82:447-52.

Tinker LF, Rosal MC, Young AF, Perri MG, Patterson RE, Van Horn L, et al. Predictors of dietary change and maintenance in the Women's Health Initiative Dietary Modification Trial. *J Am Diet Assoc.* 2007;107:1155-66.

Tjonneland A, Olsen A, Boll K, Stripp C, Christensen J, Engholm G, et al. Study design, exposure variables, and socioeconomic determinants of participation in diet, cancer and health: a population-based prospective cohort study of 57,053 men and women in Denmark. *Scandinavian Journal of Public Health* 2007;35:432-441.

Todd S, Woodward M, Tunstall-Pedoe H, Bolton-Smith C. Dietary antioxidant vitamins and fiber in the etiology of cardiovascular disease and all-causes mortality: results from the Scottish Heart Health Study. *Am J Epidemiol* 1999;150:1073-80.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Tonstad S, Butler T, Yan R, Fraser GE. Type of vegetarian diet, body weight, and prevalence of type 2 diabetes. *Diabetes Care* 2009;32:791-6.

Toobert DJ, Glasgow RE, Strycker LA, Barrera M, Jr., Radcliffe JL, Wander RC, et al. Biologic and quality-of-life outcomes from the Mediterranean Lifestyle Program: a randomized clinical trial. *Diabetes Care* 2003;26:2288-93.

Torres MRSG, Francischetti EA, Genelhu V, Sanjuliani AF. Effect of a high-calcium energy-reduced diet on abdominal obesity and cardiometabolic risk factors in obese Brazilian subjects. *International Journal of Clinical Practice* 2010;64:1076-1083.

Tortosa A, Bes-Rastrollo M, Sanchez-Villegas A, Basterra-Gortari FJ, Nunez-Cordoba JM, Martinez-Gonzalez MA. Mediterranean diet inversely associated with the incidence of metabolic syndrome: The SUN prospective cohort. *Diabetes Care* 2007;30:2957-2959.

Trapp CB, Barnard ND. Usefulness of vegetarian and vegan diets for treating type 2 diabetes. *Current Diabetes Reports* 2010;10:152-158.

Tremblay A, Drapeau V, Doucet E, Almeras N, Despres JP, Bouchard C. Fat balance and ageing: results from the Quebec Family Study. *Br J Nutr.* 1998;79:413-8.

Trichopoulos D, Lagiou P. Mediterranean diet and overall mortality differences in the European Union. *Public Health Nutr* 2004;7:949-51.

Trichopoulou A, Bamia C, Lagiou P, Trichopoulos D. Conformity to traditional Mediterranean diet and breast cancer risk in the Greek EPIC (European Prospective Investigation into Cancer and Nutrition) cohort. *Am J Clin Nutr* 2010;92:620-5.

Trichopoulou A, Bamia C, Trichopoulos D. Mediterranean diet and survival among patients with coronary heart disease in Greece. *Arch Intern Med* 2005;165:929-35.

Trichopoulou A, Bamia C, Trichopoulos D. Anatomy of health effects of Mediterranean diet: Greek EPIC prospective cohort study. *BMJ* 2009;338:b2337.

Trichopoulou A, Costacou T, Bamia C, Trichopoulos D. Adherence to a Mediterranean diet and survival in a Greek population. *N Engl J Med* 2003;348:2599-608.

Trichopoulou A, Gnardellis C, Lagiou A, Benetou V, Naska A, Trichopoulos D. Physical activity and energy intake selectively predict the waist-to-hip ratio in men but not in women. *Am J Clin Nutr.* 2001;74:574-8.

Trichopoulou A, Kouris-Blazos A, Wahlqvist ML, Gnardellis C, Lagiou P, Polychronopoulos E, et al. Diet and overall survival in elderly people. *BMJ.* 1995;311:1457-60.

Trichopoulou A, Psaltopoulou T, Orfanos P, Trichopoulos D. Diet and physical activity in relation to overall mortality amongst adult diabetics in a general population cohort. *J Intern Med* 2006;259:583-91.

Trock BJ, Hilakivi-Clarke L, Clarke R. Meta-analysis of soy intake and breast cancer risk. *J Natl Cancer Inst* 2006;98:459-71.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Tucker KL, Chen H, Hannan MT, Cupples LA, Wilson PW, Felson D, et al. Bone mineral density and dietary patterns in older adults: the Framingham Osteoporosis Study. *Am J Clin Nutr*. 2002;76:245-52.

Tucker KL, Hallfrisch J, Qiao N, Muller D, Andres R, Fleg JL. The combination of high fruit and vegetable and low saturated fat intakes is more protective against mortality in aging men than is either alone: the Baltimore Longitudinal Study of Aging. *J Nutr* 2005;135:556-61.

Tucker LA, Thomas KS. Increasing total fiber intake reduces risk of weight and fat gains in women. *J Nutr*. 2009;139:576-81.

Tuomilehto H, Gylling H, Peltonen M, Martikainen T, Sahlman J, Kokkarinen J, et al. Sustained improvement in mild obstructive sleep apnea after a diet- and physical activity-based lifestyle intervention: postinterventional follow-up. *Am J Clin Nutr* 2010;92:688-96.

Turner SM, Evans DW, Ghosh P. Feasibility of long-term plasmacholesterol reduction by diet. *Lancet* 1972;1:172-4.

Turner-McGrievy GM, Barnard ND, Scialli AR. A two-year randomized weight loss trial comparing a vegan diet to a more moderate low-fat diet. *Obesity (Silver Spring)* 2007;15:2276-81.

Turunen AW, Verkasalo PK, Kiviranta H, Pukkala E, Jula A, Mannisto S, et al. Mortality in a cohort with high fish consumption. *Int J Epidemiol* 2008;37:1008-17.

Tzima N, Pitsavos C, Panagiotakos DB, Skoumas J, Zampelas A, Chrysohoou C, et al. Mediterranean diet and insulin sensitivity, lipid profile and blood pressure levels, in overweight and obese people; the Attica study. *Lipids Health Dis* 2007;6:22.

Tzonou A, Signorello LB, Lagiou P, Wu J, Trichopoulos D, Trichopoulou A. Diet and cancer of the prostate: a case-control study in Greece. *Int J Cancer*. 1999;80:704-8.

Umesawa M, Iso H, Date C, Yamamoto A, Toyoshima H, Watanabe Y, et al. Dietary intake of calcium in relation to mortality from cardiovascular disease: the JACC Study. *Stroke* 2006;37:20-6.

Umesawa M, Iso H, Date C, Yamamoto A, Toyoshima H, Watanabe Y, et al. Relations between dietary sodium and potassium intakes and mortality from cardiovascular disease: the Japan Collaborative Cohort Study for Evaluation of Cancer Risks. *Am J Clin Nutr* 2008;88:195-202.

Unick JL, Beavers D, Jakicic JM, Kitabchi AE, Knowler WC, Wadden TA, et al. Effectiveness of lifestyle interventions for individuals with severe obesity and type 2 diabetes: results from the Look AHEAD trial. *Diabetes Care* 2011;34:2152-7.

Urquiaga I, Echeverria G, Polic G, Castillo O, Liberona Y, Rozowski J, et al. Mediterranean food and diets, global resource for the control of metabolic syndrome and chronic diseases. *World Rev Nutr Diet* 2008;98:150-73.

Wagemakers JJ, Prynne CJ, Stephen AM, Wadsworth ME. Consumption of red or processed meat does not predict risk factors for coronary heart disease; results from a cohort of British adults in 1989 and 1999. *Eur J Clin Nutr*. 2009;63:303-11.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Vainio H, Weiderpass E. Fruit and vegetables in cancer prevention. *Nutr Cancer* 2006;54:111-42.

Vajifdar BU, Goyal VS, Lokhandwala YY, Mhamunkar SR, Mahadik SP, Gawad AK, et al. Is dietary fiber beneficial in chronic ischemic heart disease? *J Assoc Physicians India*. 2000;48:871-6.

Wakai K, Matsuo K, Nagata C, Mizoue T, Tanaka K, Tsuji I, et al. Lung cancer risk and consumption of vegetables and fruit: an evaluation based on a systematic review of epidemiological evidence from Japan. *Jpn J Clin Oncol* 2011;41:693-708.

Wal JS, McBurney MI, Cho S, Dhurandhar NV. Ready-to-eat cereal products as meal replacements for weight loss. *Int J Food Sci Nutr* 2007;58:331-40.

Waleekhachonloet OA, Limwattananon C, Limwattananon S, Gross CR. Group behavior therapy versus individual behavior therapy for healthy dieting and weight control management in overweight and obese women living in rural community. *Obesity Research and Clinical Practice* 2007;1:223-232.

Walker KZ, O'Dea K, Gomez M, Girgis S, Colagiuri R. Diet and exercise in the prevention of diabetes. *J Hum Nutr Diet*. 2010;23:344-52.

van Bakel MM, Kaaks R, Feskens EJ, Rohrmann S, Welch AA, Pala V, et al. Dietary glycaemic index and glycaemic load in the European Prospective Investigation into Cancer and Nutrition. *Eur J Clin Nutr*. 2009;63 Suppl 4:S188-205.

van Dam RM, Huang Z, Giovannucci E, Rimm EB, Hunter DJ, Colditz GA, et al. Diet and basal cell carcinoma of the skin in a prospective cohort of men. *Am J Clin Nutr* 2000;71:135-41.

van Dam RM, Rimm EB, Willett WC, Stampfer MJ, Hu FB. Dietary patterns and risk for type 2 diabetes mellitus in U.S. men. *Ann Intern Med*. 2002;136:201-9.

van Dam RM, Willett WC, Rimm EB, Stampfer MJ, Hu FB. Dietary fat and meat intake in relation to risk of type 2 diabetes in men. *Diabetes Care* 2002;25:417-24.

van de Laar FA, Akkermans RP, van Binsbergen JJ. Limited evidence for effects of diet for type 2 diabetes from systematic reviews. *European Journal of Clinical Nutrition* 2007;61:929-937.

van de Vijver LP, van den Bosch LM, van den Brandt PA, Goldbohm RA. Whole-grain consumption, dietary fibre intake and body mass index in the Netherlands cohort study. *Eur J Clin Nutr* 2009;63:31-8.

Van Den Brandt PA, Smits KM, Goldbohm RA, Weijnenberg MP. Magnesium intake and colorectal cancer risk in the Netherlands Cohort Study. *British Journal of Cancer* 2007;96:510-513.

van der Heijden AAWA, Hu FB, Rimm EB, van Dam RM. A prospective study of breakfast consumption and weight gain among U.S. men. *Obesity* 2007;15:2463-2469.

van der Pols JC, Gunnell D, Williams GM, Holly JM, Bain C, Martin RM. Childhood dairy and calcium intake and cardiovascular mortality in adulthood: 65-year follow-up of the Boyd Orr cohort. *Heart* 2009;95:1600-6.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

van Gils CH, Peeters PH, Bueno-de-Mesquita HB, Boshuizen HC, Lahmann PH, Clavel-Chapelon F, et al. Consumption of vegetables and fruits and risk of breast cancer. *JAMA* 2005;293:183-93.

Van Loan M. The role of dairy foods and dietary calcium in weight management. *Journal of the American College of Nutrition* 2009;28:120S-129S.

Van Strien T, Van de Laar FA. Intake of energy is best predicted by overeating tendency and consumption of fat is best predicted by dietary restraint: a 4-year follow-up of patients with newly diagnosed Type 2 diabetes. *Appetite*. 2008;50:544-7.

van Strien T, van de Laar FA, van Leeuwe JF, Lucassen PL, van den Hoogen HJ, Rutten GE, et al. The dieting dilemma in patients with newly diagnosed type 2 diabetes: does dietary restraint predict weight gain 4 years after diagnosis? *Health Psychol*. 2007;26:105-12.

van 't Veer P. Diet and breast cancer: trial and error? *Ann Med* 1994;26:453-60.

van Woudenberg GJ, Kuijsten A, Tigcheler B, Sijbrands EJ, van Rooij FJ, Hofman A, et al. Meat consumption and its association with C-reactive protein and incident type 2 diabetes: the Rotterdam Study. *Diabetes Care* 2012;35:1499-505.

van Woudenberg GJ, van Ballegooijen AJ, Kuijsten A, Sijbrands EJ, van Rooij FJ, Geleijnse JM, et al. Eating fish and risk of type 2 diabetes: A population-based, prospective follow-up study. *Diabetes Care* 2009;32:2021-6.

Wan YP, Xu RY, Wu YJ, Chen ZQ, Cai W. Diet intervention on obese children with hypertension in China. *World Journal of Pediatrics* 2009;5:269-274.

Vang A, Singh PN, Lee JW, Haddad EH, Brinegar CH. Meats, processed meats, obesity, weight gain and occurrence of diabetes among adults: findings from Adventist Health Studies. *Ann Nutr Metab* 2008;52:96-104.

Wang C, Harris WS, Chung M, Lichtenstein AH, Balk EM, Kupelnick B, et al. n-3 Fatty acids from fish or fish-oil supplements, but not alpha-linolenic acid, benefit cardiovascular disease outcomes in primary- and secondary-prevention studies: a systematic review. *Am J Clin Nutr* 2006;84:5-17.

Wang L, Manson JE, Buring JE, Lee IM, Sesso HD. Dietary intake of dairy products, calcium, and vitamin D and the risk of hypertension in middle-aged and older women. *Hypertension* 2008;51:1073-9.

Wang Y, Simpson JA, Wluka AE, English DR, Giles GG, Graves S, et al. Meat consumption and risk of primary hip and knee joint replacement due to osteoarthritis: a prospective cohort study. *BMC Musculoskelet Disord* 2011;12:17.

Wang YF, Yancy WS, Jr., Yu D, Champagne C, Appel LJ, Lin PH. The relationship between dietary protein intake and blood pressure: results from the PREMIER study. *J Hum Hypertens*. 2008;22:745-54.

Warensjo E, Jansson JH, Cederholm T, Boman K, Eliasson M, Hallmans G, et al. Biomarkers of milk fat and the risk of myocardial infarction in men and women: A prospective, matched case-control study. *American Journal of Clinical Nutrition* 2010;92:194-202.

Warensjo E, Sundstrom J, Vessby B, Cederholm T, Riserus U. Markers of dietary fat quality and fatty acid desaturation as predictors of total and cardiovascular mortality: a population-based prospective study. *Am J Clin Nutr* 2008;88:203-9.

Varraso R, Fung TT, Barr RG, Hu FB, Willett W, Camargo CA, Jr. Prospective study of dietary patterns and chronic obstructive pulmonary disease among US women. *Am J Clin Nutr*. 2007;86:488-95.

Varraso R, Jiang R, Barr RG, Willett WC, Camargo CA, Jr. Prospective study of cured meats consumption and risk of chronic obstructive pulmonary disease in men. *Am J Epidemiol*. 2007;166:1438-45.

Varraso R, Kauffmann F, Leynaert B, Le Moual N, Boutron-Ruault MC, Clavel-Chapelon F, et al. Dietary patterns and asthma in the E3N study. *Eur Respir J*. 2009;33:33-41.

Wassertheil-Smoller S, Langford HG, Blaufox MD, Oberman A, Hawkins M, Levine B, et al. Effective dietary intervention in hypertensives: sodium restriction and weight reduction. *J Am Diet Assoc* 1985;85:423-30.

Vatanparast H, Baxter-Jones A, Faulkner RA, Bailey DA, Whiting SJ. Positive effects of vegetable and fruit consumption and calcium intake on bone mineral accrual in boys during growth from childhood to adolescence: the University of Saskatchewan Pediatric Bone Mineral Accrual Study. *Am J Clin Nutr*. 2005;82:700-6.

Vazquez JA, Adibi SA. Protein sparing during treatment of obesity: ketogenic versus nonketogenic very low calorie diet. *Metabolism* 1992;41:406-14.

Webb PM, Byrne C, Schnitt SJ, Connolly JL, Jacobs TW, Baer HJ, et al. A prospective study of diet and benign breast disease. *Cancer Epidemiol Biomarkers Prev*. 2004;13:1106-13.

Veierod MB, Laake P, Thelle DS. Dietary fat intake and risk of prostate cancer: a prospective study of 25,708 Norwegian men. *Int J Cancer*. 1997;73:634-8.

Weijenberg MP, Mullie PF, Brants HA, Heinen MM, Goldbohm RA, van den Brandt PA. Dietary glycemic load, glycemic index and colorectal cancer risk: results from the Netherlands Cohort Study. *Int J Cancer*. 2008;122:620-9.

Weikert S, Boeing H, Pischon T, Olsen A, Tjønneland A, Overvad K, et al. Fruits and vegetables and renal cell carcinoma: findings from the European prospective investigation into cancer and nutrition (EPIC). *Int J Cancer* 2006;118:3133-9.

Veldhuis L, Koppes LL, Driessen MT, Samoocha D, Twisk JW. Effects of dietary fibre intake during adolescence on the components of the metabolic syndrome at the age of 36 years: the Amsterdam Growth and Health Longitudinal Study. *J Hum Nutr Diet*. 2010;23:601-8.

Velie EM, Schairer C, Flood A, He JP, Khattree R, Schatzkin A. Empirically derived dietary patterns and risk of postmenopausal breast cancer in a large prospective cohort study. *Am J Clin Nutr*. 2005;82:1308-19.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Wengreen HJ, Munger RG, West NA, Cutler DR, Corcoran CD, Zhang J, et al. Dietary protein intake and risk of osteoporotic hip fracture in elderly residents of Utah. *J Bone Miner Res* 2004;19:537-45.

Venkateswaran V, Haddad AQ, Klotz LH, Nam R, Fleshner NE. An understanding of low-carbohydrate, high-fat diets and cancer. *Seminars in Preventive and Alternative Medicine* 2006;2:136-140.

Venn BJ, Mann JI. Cereal grains, legumes and diabetes. *Eur J Clin Nutr* 2004;58:1443-61.

Wennberg M, Bergdahl IA, Hallmans G, Norberg M, Lundh T, Skerfving S, et al. Fish consumption and myocardial infarction: a second prospective biomarker study from northern Sweden. *Am J Clin Nutr* 2011;93:27-36.

Vergnaud AC, Estaquio C, Czernichow S, Peneau S, Hercberg S, Galan P, et al. Energy density and 6-year anthropometric changes in a middle-aged adult cohort. *Br J Nutr*. 2009;102:302-9.

Vergnaud AC, Peneau S, Chat-Yung S, Kesse E, Czernichow S, Galan P, et al. Dairy consumption and 6-y changes in body weight and waist circumference in middle-aged French adults. *Am J Clin Nutr* 2008;88:1248-55.

Westman EC, Samaha FF. Trial finds low carb diets reduce weight more than low fat diets in the short term, but have more minor adverse effects. *Evidence-based Cardiovascular Medicine* 2004;8:374-377.

Whaley SE, McGregor S, Jiang L, Gomez J, Harrison G, Jenks E. A WIC-based intervention to prevent early childhood overweight. *Journal of nutrition education and behavior* 2010:S47-51.

White LR, Petrovitch H, Ross GW, Masaki K, Hardman J, Nelson J, et al. Brain aging and midlife tofu consumption. *J Am Coll Nutr*. 2000;19:242-55.

Whiteman D, Muir J, Jones L, Murphy M, Key T. Dietary questions as determinants of mortality: the OXCHECK experience. *Public Health Nutr*. 1999;2:477-87.

Wideroff L, Potischman N, Glass AG, Greer CE, Manos MM, Scott DR, et al. A nested case-control study of dietary factors and the risk of incident cytological abnormalities of the cervix. *Nutr Cancer* 1998;30:130-6.

Vignolo M, Rossi F, Bardazza G, Pistorio A, Parodi A, Spigno S, et al. Five-year follow-up of a cognitive-behavioural lifestyle multidisciplinary programme for childhood obesity outpatient treatment. *Eur J Clin Nutr* 2008;62:1047-57.

Wilfley DE, Stein RI, Saelens BE, Mockus DS, Matt GE, Hayden-Wade HA, et al. Efficacy of maintenance treatment approaches for childhood overweight: A randomized controlled trial. *JAMA: Journal of the American Medical Association* 2007;298:1661-1673.

Wilfley DE, Tibbs TL, Van Buren D, Reach KP, Walker MS, Epstein LH. Lifestyle interventions in the treatment of childhood overweight: A meta-analytic review of randomized controlled trials. *Health Psychology* 2007;26:521-532.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Villareal R. A randomized, open-label, intervention study to evaluate the effect of diet composition on weight gain in obese, type 2 diabetes patients receiving intensive insulin therapy. *Journal* 2008;364.

Willcox BJ, Yano K, Chen R, Willcox DC, Rodriguez BL, Masaki KH, et al. How much should we eat? The association between energy intake and mortality in a 36-year follow-up study of Japanese-American men. *J Gerontol A Biol Sci Med Sci* 2004;59:789-95.

Villegas R, Gao YT, Dai Q, Yang G, Cai H, Li H, et al. Dietary calcium and magnesium intakes and the risk of type 2 diabetes: the Shanghai Women's Health Study. *Am J Clin Nutr* 2009;89:1059-67.

Villegas R, Gao YT, Yang G, Li HL, Elasy TA, Zheng W, et al. Legume and soy food intake and the incidence of type 2 diabetes in the Shanghai Women's Health Study. *Am J Clin Nutr* 2008;87:162-7.

Villegas R, Liu S, Gao YT, Yang G, Li H, Zheng W, et al. Prospective study of dietary carbohydrates, glycemic index, glycemic load, and incidence of type 2 diabetes mellitus in middle-aged Chinese women. *Arch Intern Med* 2007;167:2310-6.

Villegas R, Salim A, Flynn A, Perry IJ. Prudent diet and the risk of insulin resistance. *Nutr Metab Cardiovasc Dis* 2004;14:334-43.

Willett WC. Diet and breast cancer. *J Intern Med*. 2001;249:395-411.

Willett WC. Dietary fat plays a major role in obesity: No. *Obesity Reviews* 2002;3:59-68.

Willett WC, Stampfer MJ, Manson JE, Colditz GA, Speizer FE, Rosner BA, et al. Intake of trans fatty acids and risk of coronary heart disease among women. *Lancet*. 1993;341:581-5.

Williams CL, Bollella MC, Strobino BA, Spark A, Nicklas TA, Tolosi LB, et al. "Healthy-Start": Outcome of an intervention to promote a heart healthy diet in preschool children. *Journal of the American College of Nutrition* 2002;21:62-71.

Williams CL, Strobino BA. Childhood diet, overweight, and CVD risk factors: the Healthy Start project. *Prev Cardiol* 2008;11:11-20.

Williams FM, Skinner J, Spector TD, Cassidy A, Clark IM, Davidson RM, et al. Dietary garlic and hip osteoarthritis: evidence of a protective effect and putative mechanism of action. *BMC Musculoskelet Disord*. 2010;11:280.

Williams L, Germov J, Young A. Preventing weight gain: A population cohort study of the nature and effectiveness of mid-age women's weight control practices. *International Journal of Obesity* 2007;31:978-986.

Williams PT. Effects of diet, physical activity and performance, and body weight on incident gout in ostensibly healthy, vigorously active men. *Am J Clin Nutr* 2008;87:1480-7.

Williamson DA, Martin CK, Anton SD, York-Crowe E, Han H, Redman L, et al. Is caloric restriction associated with development of eating-disorder symptoms? Results from the CALERIE trial. *Health Psychol* 2008;27:S32-42.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Williamson DF. Dietary intake and physical activity as "predictors" of weight gain in observational, prospective studies of adults. *Nutr Rev.* 1996;54:S101-9.

Wilimink AB, Welch AA, Quick CR, Burns PJ, Hubbard CS, Bradbury AW, et al. Dietary folate and vitamin B6 are independent predictors of peripheral arterial occlusive disease. *J Vasc Surg* 2004;39:513-6.

Wing RR, Epstein LH, Shapira B. The effect of increasing initial weight loss with the Scarsdale Diet on subsequent weight loss in a behavioral treatment program. *J Consult Clin Psychol* 1982;50:446-7.

Vining EP, Pyzik P, McGrogan J, Hladky H, Anand A, Kriegler S, et al. Growth of children on the ketogenic diet. *Dev Med Child Neurol.* 2002;44:796-802.

Winters BL, Mitchell DC, Smiciklas-Wright H, Grosvenor MB, Liu W, Blackburn GL. Dietary patterns in women treated for breast cancer who successfully reduce fat intake: the Women's Intervention Nutrition Study (WINS). *J Am Diet Assoc* 2004;104:551-9.

Vioque J, Weinbrenner T, Castello A, Asensio L, Garcia de la Hera M. Intake of fruits and vegetables in relation to 10-year weight gain among Spanish adults. *Obesity (Silver Spring)* 2008;16:664-70.

Wise LA, Radin RG, Palmer JR, Kumanyika SK, Boggs DA, Rosenberg L. Intake of fruit, vegetables, and carotenoids in relation to risk of uterine leiomyomata. *Am J Clin Nutr* 2011;94:1620-31.

Visentin G. Towards evidence-based practice via practice-based evidence: the Italian experience. *Fam Pract* 2008;25 Suppl 1:i71-4.

Vobecky JS, Vobecky J, Normand L. Risk and benefit of low fat intake in childhood. *Ann Nutr Metab.* 1995;39:124-33.

Voko Z, Hollander M, Hofman A, Koudstaal PJ, Breteler MM. Dietary antioxidants and the risk of ischemic stroke: the Rotterdam Study. *Neurology* 2003;61:1273-5.

Wolever TM, Gibbs AL, Mehling C, Chiasson JL, Connelly PW, Josse RG, et al. The Canadian Trial of Carbohydrates in Diabetes (CCD), a 1-y controlled trial of low-glycemic-index dietary carbohydrate in type 2 diabetes: no effect on glycated hemoglobin but reduction in C-reactive protein. *American Journal of Clinical Nutrition* 2008;87:114-125.

Wolfe AR, Arroyo C, Tedders SH, Li Y, Dai Q, Zhang J. Dietary protein and protein-rich food in relation to severely depressed mood: A 10 year follow-up of a national cohort. *Prog Neuropsychopharmacol Biol Psychiatry.* 2011;35:232-8.

Wolfe AR, Ogbonna EM, Lim S, Li Y, Zhang J. Dietary linoleic and oleic fatty acids in relation to severe depressed mood: 10 years follow-up of a national cohort. *Prog Neuropsychopharmacol Biol Psychiatry.* 2009;33:972-7.

Woodside JV, McKinley MC, Young IS. Saturated and trans fatty acids and coronary heart disease. *Current Atherosclerosis Reports* 2008;10:460-466.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Voorrips LE, Goldbohm RA, Verhoeven DT, van Poppel GA, Sturmans F, Hermus RJ, et al. Vegetable and fruit consumption and lung cancer risk in the Netherlands Cohort Study on diet and cancer. *Cancer Causes Control* 2000;11:101-15.

Wright ME, Mayne ST, Stolzenberg-Solomon RZ, Li Z, Pietinen P, Taylor PR, et al. Development of a comprehensive dietary antioxidant index and application to lung cancer risk in a cohort of male smokers. *Am J Epidemiol* 2004;160:68-76.

Wright ME, Park Y, Subar AF, Freedman ND, Albanes D, Hollenbeck A, et al. Intakes of fruit, vegetables, and specific botanical groups in relation to lung cancer risk in the NIH-AARP Diet and Health Study. *Am J Epidemiol* 2008;168:1024-34.

Wu AH, Koh WP, Wang R, Lee HP, Yu MC. Soy intake and breast cancer risk in Singapore Chinese Health Study. *Br J Cancer* 2008;99:196-200.

Wu H, Pan A, Yu Z, Qi Q, Lu L, Zhang G, et al. Lifestyle counseling and supplementation with flaxseed or walnuts influence the management of metabolic syndrome. *J Nutr* 2010;140:1937-42.

Wu K, Hu FB, Fuchs C, Rimm EB, Willett WC, Giovannucci E. Dietary patterns and risk of colon cancer and adenoma in a cohort of men (United States). *Cancer Causes Control*. 2004;15:853-62.

Wu K, Hu FB, Willett WC, Giovannucci E. Dietary patterns and risk of prostate cancer in U.S. men. *Cancer Epidemiol Biomarkers Prev*. 2006;15:167-71.

Xun P, He K. Fish Consumption and Incidence of Diabetes: meta-analysis of data from 438,000 individuals in 12 independent prospective cohorts with an average 11-year follow-up. *Diabetes Care* 2012;35:930-8.

Yamagishi K, Iso H, Date C, Fukui M, Wakai K, Kikuchi S, et al. Fish, omega-3 polyunsaturated fatty acids, and mortality from cardiovascular diseases in a nationwide community-based cohort of Japanese men and women the JACC (Japan Collaborative Cohort Study for Evaluation of Cancer Risk) Study. *J Am Coll Cardiol* 2008;52:988-96.

Yamagishi K, Iso H, Yatsuya H, Tanabe N, Date C, Kikuchi S, et al. Dietary intake of saturated fatty acids and mortality from cardiovascular disease in Japanese: the Japan Collaborative Cohort Study for Evaluation of Cancer Risk (JACC) Study. *Am J Clin Nutr* 2010;92:759-65.

Yamamoto S, Sobue T, Kobayashi M, Sasaki S, Tsugane S. Soy, isoflavones, and breast cancer risk in Japan. *J Natl Cancer Inst* 2003;95:906-13.

Yang G, Shu XO, Jin F, Zhang X, Li HL, Li Q, et al. Longitudinal study of soy food intake and blood pressure among middle-aged and elderly Chinese women. *Am J Clin Nutr*. 2005;81:1012-7.

Yannakoulia M, Ntalla I, Papoutsakis C, Farmaki AE, Dedoussis GV. Consumption of vegetables, cooked meals, and eating dinner is negatively associated with overweight status in children. *Journal of Pediatrics* 2010;157:815-820.

Yaskin J, Toner RW, Goldfarb N. Obesity management interventions: A review of the evidence. *Population Health Management* 2009;12:305-316.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Yasmin, Mascie-Taylor CGN, Brown MJ, Hughes M. The effect of dietary intervention on changes in total cholesterol, blood pressure and weight in a Cambridge study. *International Journal of Clinical Practice* 1998;52:241-245.

Yau WY, Hankey GJ. Which dietary and lifestyle behaviours may be important in the aetiology (and prevention) of stroke? *J Clin Neurosci*. 2011;18:76-80.

Yeh CC, You SL, Chen CJ, Sung FC. Peanut consumption and reduced risk of colorectal cancer in women: a prospective study in Taiwan. *World J Gastroenterol* 2006;12:222-7.

Ylitalo V. Treatment of obese schoolchildren with special reference to the mode of therapy, cardiorespiratory performance and the carbohydrate and lipid metabolism. *Acta Paediatr Scand Suppl* 1981;290:1-108.

Yochum L, Kushi LH, Meyer K, Folsom AR. Dietary flavonoid intake and risk of cardiovascular disease in postmenopausal women. *Am J Epidemiol* 1999;149:943-9.

Yochum LA, Folsom AR, Kushi LH. Intake of antioxidant vitamins and risk of death from stroke in postmenopausal women. *Am J Clin Nutr* 2000;72:476-83.

Yoo S, Nicklas T, Baranowski T, Zakeri IF, Yang SJ, Srinivasan SR, et al. Comparison of dietary intakes associated with metabolic syndrome risk factors in young adults: the Bogalusa Heart Study. *Am J Clin Nutr*. 2004;80:841-8.

Yoon H, Benamouzig R, Little J, Francois-Collange M, Tome D. Systematic review of epidemiological studies on meat, dairy products and egg consumption and risk of colorectal adenomas. *Eur J Cancer Prev* 2000;9:151-64.

Yoshihara A, Watanabe R, Hanada N, Miyazaki H. A longitudinal study of the relationship between diet intake and dental caries and periodontal disease in elderly Japanese subjects. *Gerodontology*. 2009;26:130-6.

Yuan JM, Ross RK, Gao YT, Yu MC. Fish and shellfish consumption in relation to death from myocardial infarction among men in Shanghai, China. *Am J Epidemiol* 2001;154:809-16.

Yusof BN, Talib RA, Kamaruddin NA, Karim NA, Chinna K, Gilbertson H. A low-GI diet is associated with a short-term improvement of glycaemic control in Asian patients with type 2 diabetes. *Diabetes Obes Metab*. 2009;11:387-96.

Zambon A, Sartore G, Passera D, Francini-Pesenti F, Bassi A, Basso C, et al. Effects of hypocaloric dietary treatment enriched in oleic acid on LDL and HDL subclass distribution in mildly obese women. *J Intern Med* 1999;246:191-201.

Zazpe I, Bes-Rastrollo M, Ruiz-Canela M, Sanchez-Villegas A, Serrano-Martinez M, Angel Martinez-Gonzalez M. A brief assessment of eating habits and weight gain in a Mediterranean cohort. *Br J Nutr* 2011;105:765-75.

Zemel MB. The role of dairy foods in weight management. *Journal of the American College of Nutrition* 2005;24:537S-546S.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Zemel MB, Richards J, Milstead A, Campbell P. Effects of calcium and dairy on body composition and weight loss in African-American adults. *Obes Res* 2005;13:1218-25.

Zemel MB, Thompson W, Zemel P, Nocton AM, Milstead A, Morris K, et al. Dietary calcium and dairy products accelerate weight and fat loss during energy restriction in obese adults. *American Journal of Clinical Nutrition* 2002:342s-343s.

Zhang S, Hunter DJ, Forman MR, Rosner BA, Speizer FE, Colditz GA, et al. Dietary carotenoids and vitamins A, C, and E and risk of breast cancer. *J Natl Cancer Inst* 1999;91:547-56.

Zhang S, Hunter DJ, Rosner BA, Colditz GA, Fuchs CS, Speizer FE, et al. Dietary fat and protein in relation to risk of non-Hodgkin's lymphoma among women. *J Natl Cancer Inst*. 1999;91:1751-8.

Zhang X, Shu XO, Gao YT, Yang G, Li Q, Li H, et al. Soy food consumption is associated with lower risk of coronary heart disease in Chinese women. *J Nutr* 2003;133:2874-8.

Zhang X, Shu XO, Li H, Yang G, Li Q, Gao YT, et al. Prospective cohort study of soy food consumption and risk of bone fracture among postmenopausal women. *Arch Intern Med* 2005;165:1890-5.

Zhang X, Shu XO, Xiang YB, Yang G, Li H, Gao J, et al. Cruciferous vegetable consumption is associated with a reduced risk of total and cardiovascular disease mortality. *Am J Clin Nutr* 2011;94:240-6.

Zheng J, Huang T, Yu Y, Hu X, Yang B, Li D. Fish consumption and CHD mortality: an updated meta-analysis of seventeen cohort studies. *Public Health Nutr* 2012;15:725-37.

Zheng W, Anderson KE, Kushi LH, Sellers TA, Greenstein J, Hong CP, et al. A prospective cohort study of intake of calcium, vitamin D, and other micronutrients in relation to incidence of rectal cancer among postmenopausal women. *Cancer Epidemiol Biomarkers Prev* 1998;7:221-5.

Zheng W, Sellers TA, Doyle TJ, Kushi LH, Potter JD, Folsom AR. Retinol, antioxidant vitamins, and cancers of the upper digestive tract in a prospective cohort study of postmenopausal women. *Am J Epidemiol*. 1995;142:955-60.

Zhou Y, Tian C, Jia C. Association of fish and n-3 fatty acid intake with the risk of type 2 diabetes: a meta-analysis of prospective studies. *Br J Nutr* 2012;108:408-17.

Ziegler RG, Subar AF, Craft NE, Ursin G, Patterson BH, Graubard BI. Does beta-carotene explain why reduced cancer risk is associated with vegetable and fruit intake? *Cancer Res* 1992;52:2060s-2066s.

Zoltick ES, Sahni S, McLean RR, Quach L, Casey VA, Hannan MT. Dietary protein intake and subsequent falls in older men and women: The Framingham study. *Journal of Nutrition, Health and Aging* 2011;15:147-152.

Kapitel 3.3

Tea consumption and incidence of type 2 diabetes in Europe: the EPIC-InterAct case-cohort study. *PLoS One* 2012;7:e36910.

Acharya S, Maskarinec G, Williams AE, Oshiro C, Hebshi S, Murphy SP. Nutritional changes among premenopausal women undertaking a soya based dietary intervention study in Hawaii. *Journal of Human Nutrition and Dietetics* 2004;17:413-419.

Adamsson V, Reumark A, Fredriksson IB, Hammarstrom E, Vessby B, Johansson G, et al. Effects of a healthy Nordic diet on cardiovascular risk factors in hypercholesterolaemic subjects: A randomized controlled trial (NORDIET). *Journal of Internal Medicine* 2011;269:150-159.

Ajani UA, Hennekens CH, Spelsberg A, Manson JE. Alcohol consumption and risk of type 2 diabetes mellitus among US male physicians. *Arch Intern Med* 2000;160:1025-30.

Alkerwi A, Boutsen M, Vaillant M, Barre J, Lair ML, Albert A, et al. Alcohol consumption and the prevalence of metabolic syndrome: A meta-analysis of observational studies. *Atherosclerosis* 2009;204:624-635.

Astrup A, Chaput JP, Gilbert JA, Lorenzen JK. Dairy beverages and energy balance. *Physiology and Behavior* 2010;100:67-75.

Athyros VG, Liberopoulos EN, Mikhailidis DP, Papageorgiou AA, Ganotakis ES, Tziomalos K, et al. Association of drinking pattern and alcohol beverage type with the prevalence of metabolic syndrome, diabetes, coronary heart disease, stroke, and peripheral arterial disease in a Mediterranean cohort. *Angiology*. 2007;58:689-97.

Baik I, Shin C. Prospective study of alcohol consumption and metabolic syndrome. *American Journal of Clinical Nutrition* 2008;87:1455-1463.

Baliunas DO, Taylor BJ, Irving H, Roerecke M, Patra J, Mohapatra S, et al. Alcohol as a risk factor for type 2 diabetes: A systematic review and meta-analysis. *Diabetes Care* 2009;32:2123-32.

Balk L, Hoekstra T, Twisk J. Relationship between long-term coffee consumption and components of the metabolic syndrome: the Amsterdam Growth and Health Longitudinal Study. *Eur J Epidemiol* 2009;24:203-9.

Balkau B, Vierron E, Vernay M, Born C, Arondel D, Petrella A, et al. The impact of 3-year changes in lifestyle habits on metabolic syndrome parameters: the D.E.S.I.R study. *Eur J Cardiovasc Prev Rehabil*. 2006;13:334-40.

Banim PJ, Luben RN, Bulluck H, Sharp SJ, Wareham NJ, Khaw KT, et al. The aetiology of symptomatic gallstones quantification of the effects of obesity, alcohol and serum lipids on risk. Epidemiological and biomarker data from a UK prospective cohort study (EPIC-Norfolk). *Eur J Gastroenterol Hepatol* 2011;23:733-40.

Barstad B, Sorensen TI, Tjonneland A, Johansen D, Becker U, Andersen IB, et al. Intake of wine, beer and spirits and risk of gastric cancer. *Eur J Cancer Prev* 2005;14:239-43.

Basu A, Du M, Sanchez K, Leyva MJ, Betts NM, Blevins S, et al. Green tea minimally affects biomarkers of inflammation in obese subjects with metabolic syndrome. *Nutrition* 2011;27:206-213.

Batty GD, Lewars H, Emslie C, Gale CR, Hunt K. Internationally recognized guidelines for 'sensible' alcohol consumption: is exceeding them actually detrimental to health and social circumstances? Evidence from a population-based cohort study. *J Public Health (Oxf)* 2009;31:360-5.

Bayer O, von Kries R, Strauss A, Mitschek C, Toschke AM, Hose A, et al. Short- and mid-term effects of a setting based prevention program to reduce obesity risk factors in children: a cluster-randomized trial. *Clin Nutr* 2009;28:122-8.

Bellisle F, Altenburg de Assis MA, Fieux B, Preziosi P, Galan P, Guy-Grand B, et al. Use of 'light' foods and drinks in French adults: biological, anthropometric and nutritional correlates. *J Hum Nutr Diet*. 2001;14:191-206.

Bellisle F, Drewnowski A. Intense sweeteners, energy intake and the control of body weight. *Eur J Clin Nutr* 2007;61:691-700.

Benetou V, Orfanos P, Zylis D, Sieri S, Contiero P, Tumino R, et al. Diet and hip fractures among elderly Europeans in the EPIC cohort. *Eur J Clin Nutr*. 2011;65:132-9.

Beresford SA, Johnson KC, Ritenbaugh C, Lasser NL, Snetselaar LG, Black HR, et al. Low-fat dietary pattern and risk of colorectal cancer: the Women's Health Initiative Randomized Controlled Dietary Modification Trial. *JAMA* 2006;295:643-54.

Berkey CS, Rockett HR, Colditz GA. Weight gain in older adolescent females: the internet, sleep, coffee, and alcohol. *J Pediatr*. 2008;153:635-9, 639 e1.

Berkey CS, Rockett HR, Field AE, Gillman MW, Colditz GA. Sugar-added beverages and adolescent weight change. *Obes Res* 2004;12:778-88.

Berkey CS, Rockett HR, Willett WC, Colditz GA. Milk, dairy fat, dietary calcium, and weight gain: a longitudinal study of adolescents. *Arch Pediatr Adolesc Med* 2005;159:543-50.

Bes-Rastrollo M, van Dam RM, Martinez-Gonzalez MA, Li TY, Sampson LL, Hu FB. Prospective study of dietary energy density and weight gain in women. *Am J Clin Nutr* 2008;88:769-77.

Beulens JWJ, de Zoete EC, Kok FJ, Schaafsma G, Hendriks HFJ. Effect of moderate alcohol consumption on adipokines and insulin sensitivity in lean and overweight men: A diet intervention study. *European Journal of Clinical Nutrition* 2008;62:1098-1105.

Bihan H, Takbou K, Cohen R, Michault A, Boitou F, Reach G, et al. Impact of short-duration lifestyle intervention in collaboration with general practitioners in patients with the metabolic syndrome. *Diabetes Metab* 2009;35:185-91.

Blum JE, Davee AM, Beaudoin CM, Jenkins PL, Kaley LA, Wigand DA. Reduced availability of sugar-sweetened beverages and diet soda has a limited impact on beverage consumption patterns in Maine high school youth. *J Nutr Educ Behav* 2008;40:341-7.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Bo S, Durazzo M, Guidi S, Carello M, Sacerdote C, Silli B, et al. Dietary magnesium and fiber intakes and inflammatory and metabolic indicators in middle-aged subjects from a population-based cohort. *Am J Clin Nutr* 2006;84:1062-9.

Bolling BW, Chen CYO, Blumberg JB. Tea and health: Preventive and therapeutic usefulness in the elderly? *Current Opinion in Clinical Nutrition and Metabolic Care* 2009;12:42-48.

Bonita JS, Mandarano M, Shuta D, Vinson J. Coffee and cardiovascular disease: In vitro, cellular, animal, and human studies. *Pharmacological Research* 2007;55:187-198.

Bostick RM, Potter JD, Kushi LH, Sellers TA, Steinmetz KA, McKenzie DR, et al. Sugar, meat, and fat intake, and non-dietary risk factors for colon cancer incidence in Iowa women (United States). *Cancer Causes Control*. 1994;5:38-52.

Breitling LP, Arndt V, Drath C, Rothenbacher D, Brenner H. Smoking and gamma-glutamyltransferase: opposite interactions with alcohol consumption and body mass index. *PLoS One*. 2010;5.

Breslow RA, Chen CM, Graubard BI, Mukamal KJ. Prospective study of alcohol consumption quantity and frequency and cancer-specific mortality in the US population. *Am J Epidemiol* 2011;174:1044-53.

Brown AL, Lane J, Holyoak C, Nicol B, Mayes AE, Dadd T. Health effects of green tea catechins in overweight and obese men: a randomised controlled cross-over trial. *Br J Nutr* 2011;106:1880-9.

Bueno MB, Cesar CL, Martini LA, Fisberg RM. Dietary calcium intake and overweight: an epidemiologic view. *Nutrition* 2008;24:1110-5.

Burke V, Lee AH, Hunter E, Spargo R, Smith R, Beilin LJ, et al. Alcohol intake and incidence of coronary in Australian aborigines. *Alcohol and Alcoholism* 2007;42:49-54.

Burrows T, Janet WM, Collins CE. Long-term changes in food consumption trends in overweight children in the HIKCUPS intervention. *J Pediatr Gastroenterol Nutr* 2011;53:543-7.

Burrows T, Warren JM, Baur LA, Collins CE. Impact of a child obesity intervention on dietary intake and behaviors. *Int J Obes (Lond)* 2008;32:1481-8.

Byrne C, Webb PM, Jacobs TW, Peiro G, Schnitt SJ, Connolly JL, et al. Alcohol consumption and incidence of benign breast disease. *Cancer Epidemiol Biomarkers Prev*. 2002;11:1369-74.

Cade JE, Burley VJ, Greenwood DC. The UK Women's Cohort Study: comparison of vegetarians, fish-eaters and meat-eaters. *Public Health Nutr*. 2004;7:871-8.

Canas JA, Damaso L, Altomare A, Killen K, Balagopal B. Fruit and vegetable juice concentrate (FVJC) improves glucose disposition index (GDI) and serum retinol (SR) in obese children. a 6-month pilot, randomized, double-blind placebo-controlled study. In: Reviews E, editor. 93rd Annual Meeting and Expo of the Endocrine Society, ENDO 2011 Boston, MA, United States; 2011. p 3

Champagne CM, Broyles ST, Moran LD, Cash KC, Levy EJ, Lin PH, et al. Dietary intakes associated with successful weight loss and maintenance during the Weight Loss Maintenance trial. *J Am Diet Assoc* 2011;111:1826-35.

Chang SC, Ziegler RG, Dunn B, Stolzenberg-Solomon R, Lacey JV, Jr., Huang WY, et al. Association of energy intake and energy balance with postmenopausal breast cancer in the prostate, lung, colorectal, and ovarian cancer screening trial. *Cancer Epidemiol Biomarkers Prev*. 2006;15:334-41.

Chen L, Appel LJ, Loria C, Lin PH, Champagne CM, Elmer PJ, et al. Reduction in consumption of sugar-sweetened beverages is associated with weight loss: the PREMIER trial. *Am J Clin Nutr* 2009;89:1299-306.

Chen M, Pan A, Malik VS, Hu FB. Effects of dairy intake on body weight and fat: a meta-analysis of randomized controlled trials. *Am J Clin Nutr* 2012;96:735-47.

Christensen R, Lorenzen JK, Svith CR, Bartels EM, Melanson EL, Saris WH, et al. Effect of calcium from dairy and dietary supplements on faecal fat excretion: a meta-analysis of randomized controlled trials. *Obes Rev* 2009;10:475-86.

Chu SL, Fu H, Yang JX, Liu GX, Dou P, Zhang L, et al. A randomized double-blind placebo-controlled study of Pu'er tea extract on the regulation of metabolic syndrome. *Chin J Integr Med* 2011;17:492-8.

Clavien H, Theintz G, Rizzoli R, Bonjour JP. Does puberty alter dietary habits in adolescents living in a western society? *J Adolesc Health*. 1996;19:68-75.

Clement Y. Can green tea do that? A literature review of the clinical evidence. *Preventive Medicine* 2009;49:83-87.

Colditz GA, Giovannucci E, Rimm EB, Stampfer MJ, Rosner B, Speizer FE, et al. Alcohol intake in relation to diet and obesity in women and men. *Am J Clin Nutr* 1991;54:49-55.

Collins CE, Warren JM, Neve M, McCoy P, Stokes B. Systematic review of interventions in the management of overweight and obese children which include a dietary component. *International Journal of Evidence-Based Healthcare* 2007;5:2-53.

Crawley HF, While D. The diet and body weight of British teenage smokers at 16-17 years. *Eur J Clin Nutr*. 1995;49:904-14.

Crichton GE, Murphy KJ, Howe PR, Buckley JD, Bryan J. Dairy consumption and working memory performance in overweight and obese adults. *Appetite* 2012;59:34-40.

Crichton GE, PR CH, Buckley JD, Coates AM, Murphy KJ. Dairy consumption and cardiometabolic health: outcomes of a 12-month crossover trial. *Nutr Metab (Lond)* 2012;9:19.

Cullmann M, Hilding A, Ostenson CG. Alcohol consumption and risk of pre-diabetes and type 2 diabetes development in a Swedish population. *Diabet Med* 2012;29:441-52.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

De Bacquer D, Clays E, Delanghe J, De Backer G. Epidemiological evidence for an association between habitual tea consumption and markers of chronic inflammation. *Atherosclerosis* 2006;189:428-435.

De Koning L, Malik VS, Rimm EB, Willett WC, Hu FB. Sugar-sweetened and artificially sweetened beverage consumption and risk of type 2 diabetes in men. *American Journal of Clinical Nutrition* 2011;93:1321-1327.

de La Hunty A, Gibson S, Ashwell M. A review of the effectiveness of aspartamen in helping with weight control. *Nutrition Bulletin* 2006;31:115-128.

de Ruyter JC, Olthof MR, Seidell JC, Katan MB. A Trial of Sugar-free or Sugar-Sweetened Beverages and Body Weight in Children. *N Engl J Med* 2012.

D'Elia L, Barba G, Cappuccio FP, Strazzullo P. Potassium intake, stroke, and cardiovascular disease: A meta-analysis of prospective studies. *Journal of the American College of Cardiology* 2011;57:1210-1219.

Denova-Gutierrez E, Talavera JO, Huitron-Bravo G, Mendez-Hernandez P, Salmeron J. Sweetened beverage consumption and increased risk of metabolic syndrome in Mexican adults. *Public Health Nutr* 2010;13:835-42.

Dhingra R, Sullivan L, Jacques PF, Wang TJ, Fox CS, Meigs JB, et al. Soft drink consumption and risk of developing cardiometabolic risk factors and the metabolic syndrome in middle-aged adults in the community. *Circulation* 2007;116:480-8.

Di Pierro F, Menghi AB, Barreca A, Lucarelli M, Calandrelli A. GreenSelect[®] Phytosome as an adjunct to a low-calorie diet for treatment of obesity: a clinical trial. *Alternative Medicine Review* 2009;14 154-60.

Dorn JM, Hovey K, Muti P, Freudenheim JL, Russell M, Nochajski TH, et al. Alcohol drinking patterns differentially affect central adiposity as measured by abdominal height in women and men. *J Nutr* 2003;133:2655-62.

Doymaz S, Neuspiel DR. The influence of pediatric resident counseling on limiting sugar-sweetened drinks in children. *Clin Pediatr (Phila)* 2009;48:777-9.

Drewnowski A, Almiron-Roig E, Marmonier C, Lluch A. Dietary energy density and body weight: Is there a relationship? *Nutrition Reviews* 2004;62:403-413.

Driessen MT, Koppes LL, Veldhuis L, Samoocha D, Twisk JW. Coffee consumption is not related to the metabolic syndrome at the age of 36 years: the Amsterdam Growth and Health Longitudinal Study. *Eur J Clin Nutr*. 2009;63:536-42.

Drogan D, Hoffmann K, Schulz M, Bergmann MM, Boeing H, Weikert C. A food pattern predicting prospective weight change is associated with risk of fatal but not with nonfatal cardiovascular disease. *J Nutr* 2007;137:1961-7.

Dubois L, Farmer A, Girard M, Peterson K. Regular sugar-sweetened beverage consumption between meals increases risk of overweight among preschool-aged children. *J Am Diet Assoc* 2007;107:924-34; discussion 934-5.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Duffey KJ, Steffen LM, Van Horn L, Jacobs DR, Jr., Popkin BM. Dietary patterns matter: diet beverages and cardiometabolic risks in the longitudinal Coronary Artery Risk Development in Young Adults (CARDIA) Study. *Am J Clin Nutr* 2012;95:909-15.

Eagan MS, Lyle RM, Gunther CW, Peacock M, Teegarden D. Effect of 1-year dairy product intervention on fat mass in young women: 6-month follow-up. *Obesity (Silver Spring)*. 2006;14:2242-8.

Ebbeling CB, Feldman HA, Chomitz VR, Antonelli TA, Gortmaker SL, Osganian SK, et al. A Randomized Trial of Sugar-Sweetened Beverages and Adolescent Body Weight. *N Engl J Med* 2012.

Ebbeling CB, Feldman HA, Chomitz VR, Antonelli TA, Gortmaker SL, Osganian SK, et al. Dublett A randomized trial of sugar-sweetened beverages and adolescent body weight. *N Engl J Med* 2012;367:1407-16.

Ebbeling CB, Feldman HA, Osganian SK, Chomitz VR, Ellenbogen SJ, Ludwig DS. Effects of decreasing sugar-sweetened beverage consumption on body weight in adolescents: a randomized, controlled pilot study. *Pediatrics* 2006;117:673-80.

Elwood P, Hughes J, Fehily A. Milk, heart disease and obesity: An examination of the evidence. *British Journal of Cardiology* 2005;12:283-290.

Elwood PC, Pickering JE, Fehily AM. Milk and dairy consumption, diabetes and the metabolic syndrome: the Caerphilly prospective study. *J Epidemiol Community Health* 2007;61:695-8.

Elwood PC, Pickering JE, Givens DI, Gallacher JE. The consumption of milk and dairy foods and the incidence of vascular disease and diabetes: an overview of the evidence. *Lipids* 2010;45:925-39.

Engeset D, Alsaker E, Ciampi A, Lund E. Dietary patterns and lifestyle factors in the Norwegian EPIC cohort: the Norwegian Women and Cancer (NOWAC) study. *Eur J Clin Nutr*. 2005;59:675-84.

Eshak ES, Iso H, Mizoue T, Inoue M, Noda M, Tsugane S. Soft drink, 100% fruit juice, and vegetable juice intakes and risk of diabetes mellitus. *Clin Nutr* 2012.

Faghih S, Abadi AR, Hedayati M, Kimiagar SM. Comparison of the effects of cows' milk, fortified soy milk, and calcium supplement on weight and fat loss in premenopausal overweight and obese women. *Nutr Metab Cardiovasc Dis* 2011;21:499-503.

Faith MS, Dennison BA, Edmunds LS, Stratton HH. Fruit juice intake predicts increased adiposity gain in children from low-income families: weight status-by-environment interaction. *Pediatrics* 2006;118:2066-75.

Fan AZ, Russell M, Stranges S, Dorn J, Trevisan M. Association of lifetime alcohol drinking trajectories with cardiometabolic risk. *J Clin Endocrinol Metab* 2008;93:154-61.

Farchi G, Fidanza F, Mariotti S, Menotti A. Is diet an independent risk factor for mortality? 20 year mortality in the Italian rural cohorts of the Seven Countries Study. *Eur J Clin Nutr* 1994;48:19-29.

Feig DI. Sugar-sweetened beverages and hypertension. *Future Cardiology* 2010;6:773-776.

Field AE, Coakley EH, Must A, Spadano JL, Laird N, Dietz WH, et al. Impact of overweight on the risk of developing common chronic diseases during a 10-year period. *Arch Intern Med* 2001;161:1581-6.

Field AE, Gillman MW, Rosner B, Rockett HR, Colditz GA. Association between fruit and vegetable intake and change in body mass index among a large sample of children and adolescents in the United States. *Int J Obes Relat Metab Disord*. 2003;27:821-6.

Fiorito LM, Marini M, Francis LA, Smiciklas-Wright H, Birch LL. Beverage intake of girls at age 5 y predicts adiposity and weight status in childhood and adolescence. *Am J Clin Nutr* 2009;90:935-42.

Flanagan DE, Moore VM, Godsland IF, Cockington RA, Robinson JS, Phillips DI. Alcohol consumption and insulin resistance in young adults. *Eur J Clin Invest*. 2000;30:297-301.

Flechtner-Mors M, Biesalski HK, Jenkinson CP, Adler G, Ditschuneit HH. Effects of moderate consumption of white wine on weight loss in overweight and obese subjects. *Int J Obes Relat Metab Disord* 2004;28:1420-6.

Flood A, Mitchell N, Jaeb M, Finch EA, Laqua PS, Welsh EM, et al. Energy density and weight change in a long-term weight-loss trial. *The International Journal of Behavioral Nutrition and Physical Activity* 2009;6.

Forshee RA, Anderson PA, Storey ML. Sugar-sweetened beverages and body mass index in children and adolescents: a meta-analysis. *Am J Clin Nutr* 2008;87:1662-71.

Franceschi S, Dal Maso L, Augustin L, Negri E, Parpinel M, Boyle P, et al. Dietary glycemic load and colorectal cancer risk. *Ann Oncol* 2001;12:173-8.

Friberg E, Orsini N, Mantzoros CS, Wolk A. Coffee drinking and risk of endometrial cancer--a population-based cohort study. *Int J Cancer* 2009;125:2413-7.

Fuhrman BJ, Smit E, Crespo CJ, Garcia-Palmieri MR. Coffee intake and risk of incident diabetes in Puerto Rican men: results from the Puerto Rico Heart Health Program. *Public Health Nutr* 2009;12:842-8.

Fumeron F, Lamri A, Abi Khalil C, Jaziri R, Porchay-Balderelli I, Lantieri O, et al. Dairy consumption and the incidence of hyperglycemia and the metabolic syndrome: results from a french prospective study, Data from the Epidemiological Study on the Insulin Resistance Syndrome (DESIR). *Diabetes Care* 2011;34:813-7.

Fumeron F, Lamri A, Emery N, Bellili N, Jaziri R, Porchay-Balderelli I, et al. Dairy products and the metabolic syndrome in a prospective study, DESIR. *J Am Coll Nutr* 2011;30:454S-63S.

Fung TT, Chiuve SE, McCullough ML, Rexrode KM, Logroscino G, Hu FB. Adherence to a DASH-style diet and risk of coronary heart disease and stroke in women. *Arch Intern Med* 2008;168:713-20.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Fung TT, Malik V, Rexrode KM, Manson JE, Willett WC, Hu FB. Sweetened beverage consumption and risk of coronary heart disease in women. *Am J Clin Nutr*. 2009;89:1037-42.

Gapstur SM, Potter JD, Sellers TA, Kushi LH, Folsom AR. Alcohol consumption and postmenopausal endometrial cancer: results from the Iowa Women's Health Study. *Cancer Causes Control*. 1993;4:323-9.

Gardener H, Rundek T, Markert M, Wright CB, Elkind MSV, Sacco RL. Diet Soft Drink Consumption is Associated with an Increased Risk of Vascular Events in the Northern Manhattan Study. *Journal of General Internal Medicine* 2012;1-7.

Gartside PS, Glueck CJ. The important role of modifiable dietary and behavioral characteristics in the causation and prevention of coronary heart disease hospitalization and mortality: the prospective NHANES I follow-up study. *J Am Coll Nutr* 1995;14:71-9.

Gavrilyuk O, Braaten T, Skeie G, Lund E. Coffee consumption and endometrial cancer risk among postmenopausal women in the norwegian women and cancer (NOWAC) study. *International Journal of Gynecological Cancer* 2011;21:S509.

Geleijnse JM, Launer LJ, Van der Kuip DA, Hofman A, Witteman JC. Inverse association of tea and flavonoid intakes with incident myocardial infarction: the Rotterdam Study. *Am J Clin Nutr* 2002;75:880-6.

Ghayour-Mobarhan M, Sahebkar A, Vakili R, Safarian M, Nematy M, Lotfian E, et al. Investigation of the effect of high dairy diet on body mass index and body fat in overweight and obese children. *Indian Journal of Pediatrics* 2009;76:1145-1150.

Gibson S. Sugar-sweetened soft drinks and obesity: a systematic review of the evidence from observational studies and interventions. *Nutr Res Rev* 2008;21:134-47.

Giggey PP, Wendell CR, Zonderman AB, Waldstein SR. Greater coffee intake in men is associated with steeper age-related increases in blood pressure. *Am J Hypertens* 2011;24:310-5.

Gilbert JA, Joannise DR, Chaput JP, Miegueu P, Cianflone K, Almeras N, et al. Milk supplementation facilitates appetite control in obese women during weight loss: a randomised, single-blind, placebo-controlled trial. *Br J Nutr* 2011;105:133-43.

Goldbohm RA, Chorus AM, Galindo Garre F, Schouten LJ, van den Brandt PA. Dairy consumption and 10-y total and cardiovascular mortality: a prospective cohort study in the Netherlands. *Am J Clin Nutr*. 2011;93:615-27.

Greenberg JA, Axen KV, Schnoll R, Boozer CN. Coffee, tea and diabetes: the role of weight loss and caffeine. *Int J Obes (Lond)* 2005;29:1121-9.

Greenberg JA, Boozer CN, Geliebter A. Coffee, diabetes, and weight control. *Am J Clin Nutr* 2006;84:682-93.

Gronbaek M. Factors influencing the relation between alcohol and cardiovascular disease. *Curr Opin Lipidol* 2006;17:17-21.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Gronbaek M, Becker U, Johansen D, Gottschau A, Schnohr P, Hein HO, et al. Type of alcohol consumed and mortality from all causes, coronary heart disease, and cancer. *Ann Intern Med* 2000;133:411-9.

Gronbaek M, Deis A, Sorensen TI, Becker U, Borch-Johnsen K, Muller C, et al. Influence of sex, age, body mass index, and smoking on alcohol intake and mortality. *BMJ*. 1994;308:302-6.

Grove KA, Lambert JD. Laboratory, epidemiological, and human intervention studies show that tea (*Camellia sinensis*) may be useful in the prevention of obesity. *Journal of Nutrition* 2010;140:446-453.

Halkjaer J, Sorensen TI, Tjonneland A, Togo P, Holst C, Heitmann BL. Food and drinking patterns as predictors of 6-year BMI-adjusted changes in waist circumference. *Br J Nutr* 2004;92:735-48.

Hankinson SE, Willett WC, Manson JE, Hunter DJ, Colditz GA, Stampfer MJ, et al. Alcohol, height, and adiposity in relation to estrogen and prolactin levels in postmenopausal women. *J Natl Cancer Inst* 1995;87:1297-302.

Hansel B, Kontush A, Bruckert E. Is a cardioprotective action of alcohol a myth? *Curr Opin Cardiol* 2012;27:550-5.

Happonen P, Laara E, Hiltunen L, Luukinen H. Coffee consumption and mortality in a 14-year follow-up of an elderly northern Finnish population. *Br J Nutr* 2008;99:1354-61.

Harvey-Berino J, Gold BC, Lauber R, Starinski A. The impact of calcium and dairy product consumption on weight loss. *Obes Res* 2005;13:1720-6.

Hashibe M, Sankaranarayanan R, Thomas G, Kuruvilla B, Mathew B, Somanathan T, et al. Body mass index, tobacco chewing, alcohol drinking and the risk of oral submucous fibrosis in Kerala, India. *Cancer Causes Control*. 2002;13:55-64.

Hayashino Y, Fukuhara S, Okamura T, Tanaka T, Ueshima H. High oolong tea consumption predicts future risk of diabetes among Japanese male workers: a prospective cohort study. *Diabet Med* 2011;28:805-10.

Heaney RP, Davies KM, Barger-Lux MJ. Calcium and Weight: Clinical Studies. *Journal of the American College of Nutrition* 2002;21:152S-155S.

Hinton PS, Rector RS, Donnelly JE, Smith BK, Bailey B. Total body bone mineral content and density during weight loss and maintenance on a low- or recommended-dairy weight-maintenance diet in obese men and women. *Eur J Clin Nutr* 2010;64:392-9.

Hirst K, Baranowski T, DeBar L, Foster GD, Kaufman F, Kennel P, et al. HEALTHY study rationale, design and methods: moderating risk of type 2 diabetes in multi-ethnic middle school students. *Int J Obes (Lond)* 2009;33 Suppl 4:S4-20.

Hjellvik V, Tverdal A, Strom H. Boiled coffee intake and subsequent risk for type 2 diabetes. *Epidemiology* 2011;22:418-21.

Hodgson JM, Puddey IB, Woodman RJ, Mulder TP, Fuchs D, Scott K, et al. Effects of black tea on blood pressure: a randomized controlled trial. *Arch Intern Med* 2012;172:186-8.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Hsu CH, Liao YL, Lin SC, Tsai TH, Huang CJ, Chou P. Does supplementation with green tea extract improve insulin resistance in obese type 2 diabetics? A randomized, double-blind, and placebo-controlled clinical trial. *Altern Med Rev* 2011;16:157-63.

Hu FB, Malik VS. Sugar-sweetened beverages and risk of obesity and type 2 diabetes: epidemiologic evidence. *Physiol Behav* 2010;100:47-54.

Huh SY, Rifas-Shiman SL, Rich-Edwards JW, Taveras EM, Gillman MW. Prospective association between milk intake and adiposity in preschool-aged children. *J Am Diet Assoc* 2010;110:563-70.

Huxley R, Lee CM, Barzi F, Timmermeister L, Czernichow S, Perkovic V, et al. Coffee, decaffeinated coffee, and tea consumption in relation to incident type 2 diabetes mellitus: a systematic review with meta-analysis. *Arch Intern Med* 2009;169:2053-63.

Jago R, Nicklas T, Yang SJ, Baranowski T, Zakeri I, Berenson GS. Physical activity and health enhancing dietary behaviors in young adults: Bogalusa Heart Study. *Prev Med* 2005;41:194-202.

James J, Thomas P, Cavan D, Kerr D. Preventing childhood obesity by reducing consumption of carbonated drinks: cluster randomised controlled trial. *BMJ* 2004;328:1237.

Jee SH, He J, Appel LJ, Whelton PK, Suh I, Klag MJ. Coffee consumption and serum lipids: a meta-analysis of randomized controlled clinical trials. *Am J Epidemiol* 2001;153:353-62.

Jing Y, Han G, Hu Y, Bi Y, Li L, Zhu D. Tea consumption and risk of type 2 diabetes: a meta-analysis of cohort studies. *J Gen Intern Med* 2009;24:557-62.

Johansen KS, Bjorge B, Hjellset VT, Holmboe-Ottesen G, Raberg M, Wandel M. Changes in food habits and motivation for healthy eating among Pakistani women living in Norway: results from the InnvaDiab-DEPLAN study. *Public Health Nutr* 2010;13:858-67.

Johnson L, Mander AP, Jones LR, Emmett PM, Jebb SA. Is sugar-sweetened beverage consumption associated with increased fatness in children? *Nutrition* 2007;23:557-63.

Joosten MM, Grobbee DE, van der AD, Verschuren WM, Hendriks HF, Beulens JW. Combined effect of alcohol consumption and lifestyle behaviors on risk of type 2 diabetes. *Am J Clin Nutr* 2010;91:1777-83.

Kakuta Y, Nakaya N, Nagase S, Fujita M, Koizumi T, Okamura C, et al. Case-control study of green tea consumption and the risk of endometrial endometrioid adenocarcinoma. *Cancer Causes Control* 2009;20:617-24.

Kanders BS, Lavin PT, Kowalchuk MB, Greenberg I, Blackburn GL. An evaluation of the effect of aspartame on weight loss. *Appetite* 1988;11 Suppl 1:73-84.

Kant AK, Schatzkin A, Graubard BI, Schairer C. A prospective study of diet quality and mortality in women. *JAMA*. 2000;283:2109-15.

Kastorini CM, Millionis HJ, Esposito K, Giugliano D, Goudevenos JA, Panagiotakos DB. The effect of mediterranean diet on metabolic syndrome and its components: A meta-analysis of 50 studies and 534,906 individuals. *Journal of the American College of Cardiology* 2011;57:1299-1313.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Kennedy BM, Champagne CM, Ryan DH, Newton R, Jr., Conish BK, Harsha DW, et al. The "Rolling Store:" an economical and environmental approach to the prevention of weight gain in African American women. *Ethn Dis* 2009;19:7-12.

Kerr WC, Ye Y. Relationship of life-course drinking patterns to diabetes, heart problems, and hypertension among those 40 and older in the 2005 U.S. National Alcohol Survey. *J Stud Alcohol Drugs* 2010;71:515-25.

Kipping RR, Jago R, Lawlor DA. Diet outcomes of a pilot school-based randomised controlled obesity prevention study with 9-10 year olds in England. *Prev Med* 2010;51:56-62.

Klesges RC, Klesges LM, Haddock CK, Eck LH. A longitudinal analysis of the impact of dietary intake and physical activity on weight change in adults. *Am J Clin Nutr* 1992;55:818-22.

Koh-Banerjee P, Chu NF, Spiegelman D, Rosner B, Colditz G, Willett W, et al. Prospective study of the association of changes in dietary intake, physical activity, alcohol consumption, and smoking with 9-y gain in waist circumference among 16 587 US men. *Am J Clin Nutr*. 2003;78:719-27.

Koppes LL, Boon N, Nooyens AC, van Mechelen W, Saris WH. Macronutrient distribution over a period of 23 years in relation to energy intake and body fatness. *Br J Nutr*. 2009;101:108-15.

Koppes LL, Dekker JM, Hendriks HF, Bouter LM, Heine RJ. Moderate alcohol consumption lowers the risk of type 2 diabetes: a meta-analysis of prospective observational studies. *Diabetes Care* 2005;28:719-25.

Kovacs EM, Lejeune MP, Nijs I, Westerterp-Plantenga MS. Effects of green tea on weight maintenance after body-weight loss. *Br J Nutr* 2004;91:431-7.

Kral TV, Stunkard AJ, Berkowitz RI, Stallings VA, Moore RH, Faith MS. Beverage consumption patterns of children born at different risk of obesity. *Obesity (Silver Spring)* 2008;16:1802-8.

Kratz M, Baars T, Guyenet S. The relationship between high-fat dairy consumption and obesity, cardiovascular, and metabolic disease. *Eur J Nutr* 2012.

Kritchevsky SB. Dietary lipids and the low blood cholesterol-cancer association. *American Journal of Epidemiology* 1992;135:509-520.

Kromhout D, Saris WH, Horst CH. Energy intake, energy expenditure, and smoking in relation to body fatness: the Zutphen Study. *Am J Clin Nutr*. 1988;47:668-74.

Kuriyama S. The relation between green tea consumption and cardiovascular disease as evidenced by epidemiological studies. *J Nutr* 2008;138:1548S-1553S.

Lahti-Koski M, Pietinen P, Heliovaara M, Vartiainen E. Associations of body mass index and obesity with physical activity, food choices, alcohol intake, and smoking in the 1982-1997 FINRISK Studies. *Am J Clin Nutr* 2002;75:809-17.

Langsetmo L, Hanley DA, Prior JC, Barr SI, Anastassiades T, Towheed T, et al. Dietary patterns and incident low-trauma fractures in postmenopausal women and men aged ≥ 50 y: a population-based cohort study. *Am J Clin Nutr* 2011;93:192-9.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

LaRowe TL, Adams AK, Jobe JB, Cronin KA, Vannatter SM, Prince RJ. Dietary intakes and physical activity among preschool-aged children living in rural American Indian communities before a family-based healthy lifestyle intervention. *J Am Diet Assoc* 2010;110:1049-57.

Larsson SC, Wolk A. Coffee consumption and risk of liver cancer: a meta-analysis. *Gastroenterology* 2007;132:1740-5.

Laurson K, Eisenmann JC, Moore S. Lack of association between television viewing, soft drinks, physical activity and body mass index in children. *Acta Paediatr* 2008;97:795-800.

Lee DH, Ha MH, Christiani DC. Body weight, alcohol consumption and liver enzyme activity--a 4-year follow-up study. *Int J Epidemiol*. 2001;30:766-70.

Libuda L, Alexy U, Sichert-Hellert W, Stehle P, Karaolis-Danckert N, Buyken AE, et al. Pattern of beverage consumption and long-term association with body-weight status in German adolescents--results from the DONALD study. *Br J Nutr*. 2008;99:1370-9.

Lien do TK, Nhung BT, Khan NC, Hop le T, Nga NT, Hung NT, et al. Impact of milk consumption on performance and health of primary school children in rural Vietnam. *Asia Pac J Clin Nutr*. 2009;18:326-34.

Lin WY, Xavier Pi-Sunyer F, Chen CC, Davidson LE, Liu CS, Li TC, et al. Coffee consumption is inversely associated with type 2 diabetes in Chinese. *Eur J Clin Invest* 2011;41:659-66.

Liu S, Serdula MK, Williamson DF, Mokdad AH, Byers T. A prospective study of alcohol intake and change in body weight among US adults. *Am J Epidemiol* 1994;140:912-20.

Loerbroks A, Schouten LJ, Goldbohm RA, van den Brandt PA. Alcohol consumption, cigarette smoking, and endometrial cancer risk: results from the Netherlands Cohort Study. *Cancer Causes Control*. 2007;18:551-60.

Loos RJ, Rankinen T, Leon AS, Skinner JS, Wilmore JH, Rao DC, et al. Calcium intake is associated with adiposity in black and white men and white women of the HERITAGE Family Study. *J Nutr* 2004;134:1772-8.

Lopez-Garcia E, Rodriguez-Artalejo F, Li TY, Mukamal KJ, Hu FB, van Dam RM. Coffee consumption and mortality in women with cardiovascular disease. *Am J Clin Nutr* 2011;94:218-24.

Louie JC, Flood VM, Hector DJ, Rangan AM, Gill TP. Dairy consumption and overweight and obesity: a systematic review of prospective cohort studies. *Obes Rev* 2011;12:e582-92.

Lucas M, Mirzaei F, Pan A, Okereke OI, Willett WC, O'Reilly EJ, et al. Coffee, caffeine, and risk of depression among women. *Arch Intern Med* 2011;171:1571-8.

Ludwig DS, Peterson KE, Gortmaker SL. Relation between consumption of sugar-sweetened drinks and childhood obesity: a prospective, observational analysis. *Lancet* 2001;357:505-8.

Maclure KM, Hayes KC, Colditz GA, Stampfer MJ, Speizer FE, Willett WC. Weight, diet, and the risk of symptomatic gallstones in middle-aged women. *N Engl J Med* 1989;321:563-9.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Maersk M, Belza A, Stodkilde-Jorgensen H, Ringgaard S, Chabanova E, Thomsen H, et al. Sucrose-sweetened beverages increase fat storage in the liver, muscle, and visceral fat depot: a 6-mo randomized intervention study. *Am J Clin Nutr* 2012;95:283-9.

Mai XM, Becker AB, Sellers EA, Liem JJ, Kozyrskyj AL. Infrequent milk consumption plus being overweight may have great risk for asthma in girls. *Allergy* 2007;62:1295-301.

Maki T, Pham NM, Yoshida D, Yin G, Ohnaka K, Takayanagi R, et al. The relationship of coffee and green tea consumption with high-sensitivity C-reactive protein in Japanese men and women. *Clin Chem Lab Med*. 2010;48:849-54.

Malik VS, Schulze MB, Hu FB. Intake of sugar-sweetened beverages and weight gain: a systematic review. *Am J Clin Nutr* 2006;84:274-88.

Maskarinec G, Meng L, Kolonel LN. Alcohol intake, body weight, and mortality in a multiethnic prospective cohort. *Epidemiology* 1998;9:654-61.

Mattes RD, Shikany JM, Kaiser KA, Allison DB. Nutritively sweetened beverage consumption and body weight: A systematic review and meta-analysis of randomized experiments. *Obesity Reviews* 2011;12:346-365.

McConahy KL, Smiciklas-Wright H, Birch LL, Mitchell DC, Picciano MF. Food portions are positively related to energy intake and body weight in early childhood. *J Pediatr* 2002;140:340-7.

Moore TJ, Alsabeeh N, Apovian CM, Murphy MC, Coffman GA, Cullum-Dugan D, et al. Weight, blood pressure, and dietary benefits after 12 months of a web-based nutrition education program (DASH for health): Longitudinal observational study. *SOURCE Journal of Medical Internet Research* (2008) 10:4 Article Number: e52. Date of Publication: October-December 2008 2008.

Mozaffarian D, Hao T, Rimm EB, Willett WC, Hu FB. Changes in diet and lifestyle and long-term weight gain in women and men. *N Engl J Med* 2011;364:2392-404.

Muckelbauer R, Libuda L, Clausen K, Toschke AM, Reinehr T, Kersting M. Promotion and provision of drinking water in schools for overweight prevention: Randomized, controlled cluster trial. *Pediatrics* 2009;123:e661-e667.

Muckelbauer R, Libuda L, Clausen K, Toschke AM, Reinehr T, Kersting M. Immigrational background affects the effectiveness of a school-based overweight prevention program promoting water consumption. *Obesity (Silver Spring)* 2010;18:528-34.

Muley A, Muley P, Shah M. Coffee to reduce risk of type 2 diabetes?: a systematic review. *Curr Diabetes Rev* 2012;8:162-8.

Mundt CA, Baxter-Jones AD, Whiting SJ, Bailey DA, Faulkner RA, Mirwald RL. Relationships of activity and sugar drink intake on fat mass development in youths. *Med Sci Sports Exerc*. 2006;38:1245-54.

Murakami K, Okubo H, Sasaki S. Effect of dietary factors on incidence of type 2 diabetes: a systematic review of cohort studies. *J Nutr Sci Vitaminol (Tokyo)* 2005;51:292-310.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Murphy RA, Mourtzakis M, Chu QSC, Baracos VE, Reiman T, Mazurak VC. Nutritional intervention with fish oil provides a benefit over standard of care for weight and skeletal muscle mass in patients with nonsmall cell lung cancer receiving chemotherapy. *Cancer* 2011;117:1775-1782.

Newby PK, Peterson KE, Berkey CS, Leppert J, Willett WC, Colditz GA. Beverage consumption is not associated with changes in weight and body mass index among low-income preschool children in North Dakota. *J Am Diet Assoc* 2004;104:1086-94.

Nissinen K, Mikkilä V, Mannisto S, Lahti-Koski M, Rasanen L, Viikari J, et al. Sweets and sugar-sweetened soft drink intake in childhood in relation to adult BMI and overweight. The Cardiovascular Risk in Young Finns Study. *Public Health Nutr* 2009;12:2018-26.

Novotny R, Daida YG, Acharya S, Grove JS, Vogt TM. Dairy intake is associated with lower body fat and soda intake with greater weight in adolescent girls. *J Nutr*. 2004;134:1905-9.

Oba S, Nagata C, Nakamura K, Fujii K, Kawachi T, Takatsuka N, et al. Diet based on the Japanese Food Guide Spinning Top and subsequent mortality among men and women in a general Japanese population. *J Am Diet Assoc* 2009;109:1540-7.

Oba S, Nagata C, Nakamura K, Fujii K, Kawachi T, Takatsuka N, et al. Consumption of coffee, green tea, oolong tea, black tea, chocolate snacks and the caffeine content in relation to risk of diabetes in Japanese men and women. *Br J Nutr* 2010;103:453-9.

Onat A, Hergenc G, Kucukdurmaz Z, Ugur M, Kaya Z, Can G, et al. Moderate and heavy alcohol consumption among Turks: long-term impact on mortality and cardiometabolic risk. *Turk Kardiyol Dern Ars* 2009;37:83-90.

Ortiz-Moncada R, Garcia M, Gonzalez-Zapata LI, Fernandez E, Alvarez-Dardet C. Incidence of overweight and obesity in a Mediterranean population-based cohort: the Cornella Health Interview Survey Follow-up Study (CHIS.FU). *Prev Med* 2010;50:45-9.

Osler M, Heitmann BL, Gerdes LU, Jorgensen LM, Schroll M. Dietary patterns and mortality in Danish men and women: a prospective observational study. *Br J Nutr* 2001;85:219-25.

Paganini-Hill A, Kawas CH, Corrada MM. Non-alcoholic beverage and caffeine consumption and mortality: the Leisure World Cohort Study. *Prev Med* 2007;44:305-10.

Pai JK, Mukamal KJ, Rimm EB. Long-term alcohol consumption in relation to all-cause and cardiovascular mortality among survivors of myocardial infarction: the Health Professionals Follow-up Study. *Eur Heart J* 2012;33:1598-605.

Pala V, Sieri S, Berrino F, Vineis P, Sacerdote C, Palli D, et al. Yogurt consumption and risk of colorectal cancer in the Italian European prospective investigation into cancer and nutrition cohort. *Int J Cancer* 2011;129:2712-9.

Pan A, Malik VS, Schulze MB, Manson JE, Willett WC, Hu FB. Plain-water intake and risk of type 2 diabetes in young and middle-aged women. *Am J Clin Nutr* 2012;95:1454-60.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Park SY, Murphy SP, Wilkens LR, Yamamoto JF, Sharma S, Hankin JH, et al. Dietary patterns using the Food Guide Pyramid groups are associated with sociodemographic and lifestyle factors: the multiethnic cohort study. *J Nutr*. 2005;135:843-9.

Patra J, Taylor B, Irving H, Roerecke M, Baliunas D, Mohapatra S, et al. Alcohol consumption and the risk of morbidity and mortality for different stroke types--a systematic review and meta-analysis. *BMC Public Health* 2010;10:258.

Paynter NP, Yeh HC, Voutilainen S, Schmidt MI, Heiss G, Folsom AR, et al. Coffee and sweetened beverage consumption and the risk of type 2 diabetes mellitus: the atherosclerosis risk in communities study. *Am J Epidemiol* 2006;164:1075-84.

Pereira MA, Jacobs DR, Jr. Sugar-sweetened beverages, weight gain and nutritional epidemiological study design. *Br J Nutr* 2008;99:1169-70.

Pereira MA, Jacobs DR, Jr., Van Horn L, Slattery ML, Kartashov AI, Ludwig DS. Dairy consumption, obesity, and the insulin resistance syndrome in young adults: the CARDIA Study. *JAMA* 2002;287:2081-9.

Pereira MA, Parker ED, Folsom AR. Coffee consumption and risk of type 2 diabetes mellitus: an 11-year prospective study of 28 812 postmenopausal women. *Arch Intern Med* 2006;166:1311-6.

Potter JD, Steinmetz K. Vegetables, fruit and phytoestrogens as preventive agents. *IARC Sci Publ* 1996:61-90.

Puddey IB, Parker M, Beilin LJ, Vandongen R, Masarei JR. Effects of alcohol and caloric restrictions on blood pressure and serum lipids in overweight men. *Hypertension* 1992;20:533-41.

Qi Q, Chu AY, Kang JH, Jensen MK, Curhan GC, Pasquale LR, et al. Sugar-Sweetened Beverages and Genetic Risk of Obesity. *N Engl J Med* 2012.

Raben A, Vasilaras TH, Moller AC, Astrup A. Sucrose compared with artificial sweeteners: different effects on ad libitum food intake and body weight after 10 wk of supplementation in overweight subjects. *Am J Clin Nutr* 2002;76:721-9.

Rajeshwari R, Yang SJ, Nicklas TA, Berenson GS. Secular trends in children's sweetened-beverage consumption (1973 to 1994): the Bogalusa Heart Study. *J Am Diet Assoc*. 2005;105:208-14.

Rajpathak SN, Rimm EB, Rosner B, Willett WC, Hu FB. Calcium and dairy intakes in relation to long-term weight gain in US men. *Am J Clin Nutr* 2006;83:559-66.

Rockell JE, Williams SM, Taylor RW, Grant AM, Jones IE, Goulding A. Two-year changes in bone and body composition in young children with a history of prolonged milk avoidance. *Osteoporos Int*. 2005;16:1016-23.

Rodearmel SJ, Wyatt HR, Stroebele N, Smith SM, Ogden LG, Hill JO. Small changes in dietary sugar and physical activity as an approach to preventing excessive weight gain: the America on the Move family study. *Pediatrics* 2007;120:e869-79.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Roerecke M, Greenfield TK, Kerr WC, Bondy S, Cohen J, Rehm J. Heavy drinking occasions in relation to ischaemic heart disease mortality-- an 11-22 year follow-up of the 1984 and 1995 US National Alcohol Surveys. *Int J Epidemiol* 2011;40:1401-10.

Roerecke M, Rehm J. The cardioprotective association of average alcohol consumption and ischaemic heart disease: a systematic review and meta-analysis. *Addiction* 2012;107:1246-60.

Roh WG, Shin HC, Choi JH, Lee YJ, Kim K. Alcohol consumption and higher incidence of impaired fasting glucose or type 2 diabetes in obese Korean men. *Alcohol* 2009;43:643-8.

Rolland-Cachera MF, Deheeger M, Akrouf M, Bellisle F. Influence of macronutrients on adiposity development: a follow up study of nutrition and growth from 10 months to 8 years of age. *Int J Obes Relat Metab Disord* 1995;19:573-8.

Romelsjö A, Allebeck P, Andreasson S, Leifman A. Alcohol, mortality and cardiovascular events in a 35 year follow-up of a nationwide representative cohort of 50,000 Swedish conscripts up to age 55. *Alcohol* 2012;47:322-7.

Ronksley PE, Brien SE, Turner BJ, Mukamal KJ, Ghali WA. Association of alcohol consumption with selected cardiovascular disease outcomes: a systematic review and meta-analysis. *BMJ* 2011;342:d671.

Rosell M, Hakansson NN, Wolk A. Association between dairy food consumption and weight change over 9 y in 19,352 perimenopausal women. *Am J Clin Nutr* 2006;84:1481-8.

Rosenbloom JI, Mukamal KJ, Frost LE, Mittleman MA. Alcohol consumption patterns, beverage type, and long-term mortality among women survivors of acute myocardial infarction. *Am J Cardiol* 2012;109:147-52.

Ruusunen A, Lehto SM, Tolmunen T, Mursu J, Kaplan GA, Voutilainen S. Coffee, tea and caffeine intake and the risk of severe depression in middle-aged Finnish men: the Kuopio Ischaemic Heart Disease Risk Factor Study. *Public Health Nutr*. 2010;13:1215-20.

Saelens BE, Couch SC, Wosje KS, Stark LJ, Daniels SR. Relations among milk and non-milk beverage consumption, calcium, and relative weight in high-weight status children. *Journal of Clinical Psychology in Medical Settings* 2006;13:117-125.

Sahyoun NR, Jacques PF, Zhang XL, Juan W, McKeown NM. Whole-grain intake is inversely associated with the metabolic syndrome and mortality in older adults. *Am J Clin Nutr* 2006;83:124-31.

Salinardi TC, Rubin KH, Black RM, St-Onge MP. Coffee manooligosaccharides, consumed as part of a free-living, weight-maintaining diet, increase the proportional reduction in body volume in overweight men. *J Nutr* 2010;140:1943-8.

Sanchez-Villegas A, Bes-Rastrollo M, Martinez-Gonzalez MA, Serra-Majem L. Adherence to a Mediterranean dietary pattern and weight gain in a follow-up study: the SUN cohort. *Int J Obes (Lond)*. 2006;30:350-8.

Sayon-Orea C, Martinez-Gonzalez MA, Bes-Rastrollo M. Alcohol consumption and body weight: a systematic review. *Nutr Rev* 2011;69:419-31.

Schroder KE. Effects of fruit consumption on body mass index and weight loss in a sample of overweight and obese dieters enrolled in a weight-loss intervention trial. *Nutrition* 2010;26:727-34.

Schulz M, Nothlings U, Hoffmann K, Bergmann MM, Boeing H. Identification of a food pattern characterized by high-fiber and low-fat food choices associated with low prospective weight change in the EPIC-Potsdam cohort. *J Nutr* 2005;135:1183-9.

Schulze MB, Manson JE, Ludwig DS, Colditz GA, Stampfer MJ, Willett WC, et al. Sugar-sweetened beverages, weight gain, and incidence of type 2 diabetes in young and middle-aged women. *JAMA* 2004;292:927-34.

Schulze MB, Manson JE, Willett WC, Hu FB. Processed meat intake and incidence of Type 2 diabetes in younger and middle-aged women. *Diabetologia* 2003;46:1465-73.

Schutze M, Boeing H, Pischon T, Rehm J, Kehoe T, Gmel G, et al. Alcohol attributable burden of incidence of cancer in eight European countries based on results from prospective cohort study. *BMJ* 2011;342:d1584.

Schutze M, Schulz M, Steffen A, Bergmann MM, Kroke A, Lissner L, et al. Beer consumption and the 'beer belly': scientific basis or common belief? *Eur J Clin Nutr* 2009;63:1143-9.

Shahar DR, Abel R, Elhayany A, Vardi H, Fraser D. Does dairy calcium intake enhance weight loss among overweight diabetic patients? *Diabetes Care* 2007;30:485-9.

Shahar DR, Schwarzfuchs D, Fraser D, Vardi H, Thiery J, Fiedler GM, et al. Dairy calcium intake, serum vitamin D, and successful weight loss. *Am J Clin Nutr* 2010;92:1017-22.

Shaw DI, Tierney AC, McCarthy S, Upritchard J, Vermunt S, Gulseth HL, et al. LIPGENE food-exchange model for alteration of dietary fat quantity and quality in free-living participants from eight European countries. *British Journal of Nutrition* 2009;101:750-759.

Shen Z, Li Y, Yu C, Shen Y, Xu L, Xu C, et al. A cohort study of the effect of alcohol consumption and obesity on serum liver enzyme levels. *Eur J Gastroenterol Hepatol* 2010;22:820-5.

Shenoy SF, Poston WS, Reeves RS, Kazaks AG, Holt RR, Keen CL, et al. Weight loss in individuals with metabolic syndrome given DASH diet counseling when provided a low sodium vegetable juice: a randomized controlled trial. *Nutr J* 2010;9:8.

Shenoy SF, Poston WSC, Reeves RS, Kazaks AG, Holt RR, Keen CL, et al. Weight loss in individuals with metabolic syndrome given DASH diet counseling when provided a low sodium vegetable juice: A randomized controlled trial. *SOURCE Nutrition Journal* (2010) 9:1 Article Number: 8. Date of Publication: 2010 2010.

Shimizu N, Nagata C, Shimizu H, Kametani M, Takeyama N, Ohnuma T, et al. Height, weight, and alcohol consumption in relation to the risk of colorectal cancer in Japan: a prospective study. *Br J Cancer* 2003;88:1038-43.

Sichieri R, Paula Trotte A, de Souza RA, Veiga GV. School randomised trial on prevention of excessive weight gain by discouraging students from drinking sodas. *Public Health Nutr* 2009;12:197-202.

Sieri S, Krogh V, Saieva C, Grobbee DE, Bergmann M, Rohrmann S, et al. Alcohol consumption patterns, diet and body weight in 10 European countries. *Eur J Clin Nutr* 2009;63 Suppl 4:S81-100.

Silvera SA, Jain M, Howe GR, Miller AB, Rohan TE. Energy balance and breast cancer risk: a prospective cohort study. *Breast Cancer Res Treat.* 2006;97:97-106.

Sinha R, Cross AJ, Daniel CR, Graubard BI, Wu JW, Hollenbeck AR, et al. Caffeinated and decaffeinated coffee and tea intakes and risk of colorectal cancer in a large prospective study. *Am J Clin Nutr* 2012;96:374-81.

Skinner JD, Bounds W, Carruth BR, Ziegler P. Longitudinal calcium intake is negatively related to children's body fat indexes. *J Am Diet Assoc* 2003;103:1626-31.

Skinner JD, Carruth BR, Moran J, 3rd, Houck K, Coletta F. Fruit juice intake is not related to children's growth. *Pediatrics.* 1999;103:58-64.

Sluijs I, Forouhi NG, Beulens JW, van der Schouw YT, Agnoli C, Arriola L, et al. The amount and type of dairy product intake and incident type 2 diabetes: results from the EPIC-InterAct Study. *Am J Clin Nutr* 2012;96:382-90.

Smith B, Wingard DL, Smith TC, Kritz-Silverstein D, Barrett-Connor E. Does coffee consumption reduce the risk of type 2 diabetes in individuals with impaired glucose? *Diabetes Care* 2006;29:2385-90.

Snijder MB, van Dam RM, Stehouwer CD, Hiddink GJ, Heine RJ, Dekker JM. A prospective study of dairy consumption in relation to changes in metabolic risk factors: the Hoorn Study. *Obesity (Silver Spring)* 2008;16:706-9.

Soenen S, Hochstenbach-Waelen A, Westerterp-Plantenga MS. Efficacy of alpha-lactalbumin and milk protein on weight loss and body composition during energy restriction. *Obesity (Silver Spring)* 2011;19:370-9.

Steffensen FH, Sorensen HT, Brock A, Vilstrup H, Lauritzen T. Alcohol consumption and serum liver-derived enzymes in a Danish population aged 30-50 years. *Int J Epidemiol.* 1997;26:92-9.

Summerbell CD, Watts C, Higgins JP, Garrow JS. Randomised controlled trial of novel, simple, and well supervised weight reducing diets in outpatients. *BMJ* 1998;317:1487-9.

Sun CL, Yuan JM, Koh WP, Yu MC. Green tea, black tea and colorectal cancer risk: a meta-analysis of epidemiologic studies. *Carcinogenesis* 2006;27:1301-9.

Sun CL, Yuan JM, Koh WP, Yu MC. Green tea, black tea and breast cancer risk: a meta-analysis of epidemiological studies. *Carcinogenesis* 2006;27:1310-5.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Suzuki R, Iwasaki M, Inoue M, Sasazuki S, Sawada N, Yamaji T, et al. Alcohol consumption-associated breast cancer incidence and potential effect modifiers: The Japan Public Health Center-based prospective study. *International Journal of Cancer* 2010;127:685-695.

Tam CS, Garnett SP, Cowell CT, Campbell K, Cabrera G, Baur LA. Soft drink consumption and excess weight gain in Australian school students: results from the Nepean study. *Int J Obes (Lond)* 2006;30:1091-3.

Tamakoshi A, Lin Y, Kawado M, Yagyu K, Kikuchi S, Iso H. Effect of coffee consumption on all-cause and total cancer mortality: findings from the JACC study. *Eur J Epidemiol* 2011;26:285-93.

Tate DF, Turner-McGrievy G, Lyons E, Stevens J, Erickson K, Polzien K, et al. Replacing caloric beverages with water or diet beverages for weight loss in adults: main results of the Choose Healthy Options Consciously Everyday (CHOICE) randomized clinical trial. *Am J Clin Nutr* 2012;95:555-63.

Tate DF, Turner-McGrievy G, Stevens J, Erickson K, Polzien K, Diamond M, et al. Replacing caloric beverages with water or diet beverages for weight loss in adults: Results of a 6-month randomized controlled trial. 29th Annual Scientific Meeting of the Obesity Society, *Obesity* 2011;19:S68.

Taveras EM, Gortmaker SL, Hohman KH, Horan CM, Kleinman KP, Mitchell K, et al. Randomized controlled trial to improve primary care to prevent and manage childhood obesity: the High Five for Kids study. *Arch Pediatr Adolesc Med* 2011;165:714-22.

Tong X, Dong JY, Wu ZW, Li W, Qin LQ. Dairy consumption and risk of type 2 diabetes mellitus: a meta-analysis of cohort studies. *Eur J Clin Nutr* 2011;65:1027-31.

Torres MRSG, Francischetti EA, Genelhu V, Sanjuliani AF. Effect of a high-calcium energy-reduced diet on abdominal obesity and cardiometabolic risk factors in obese Brazilian subjects. *International Journal of Clinical Practice* 2010;64:1076-1083.

Toubro S, Astrup A. Randomised comparison of diets for maintaining obese subjects' weight after major weight loss: ad lib, low fat, high carbohydrate diet v fixed energy intake. *BMJ* 1997;314:29-34.

Trichopoulos D, Bamia C, Lagiou P, Fedirko V, Trepo E, Jenab M, et al. Hepatocellular carcinoma risk factors and disease burden in a European cohort: a nested case-control study. *J Natl Cancer Inst* 2011;103:1686-95.

Troy JD, Hartge P, Weissfeld JL, Oken MM, Colditz GA, Mechanic LE, et al. Associations between anthropometry, cigarette smoking, alcohol consumption, and non-Hodgkin lymphoma in the Prostate, Lung, Colorectal, and Ovarian Cancer Screening Trial. *Am J Epidemiol.* 2010;171:1270-81.

Vadstrup ES, Petersen L, Sorensen TI, Gronbaek M. Waist circumference in relation to history of amount and type of alcohol: results from the Copenhagen City Heart Study. *Int J Obes Relat Metab Disord.* 2003;27:238-46.

Wagner G, Kindrick S, Hertzler S, DiSilvestro RA. Effects of various forms of calcium on body weight and bone turnover markers in women participating in a weight loss program. *J Am Coll Nutr* 2007;26:456-61.

Waki K, Noda M, Sasaki S, Matsumura Y, Takahashi Y, Isogawa A, et al. Alcohol consumption and other risk factors for self-reported diabetes among middle-aged Japanese: a population-based prospective study in the JPHC study cohort I. *Diabet Med*. 2005;22:323-31.

van Dam RM. Coffee consumption and risk of type 2 diabetes, cardiovascular diseases, and cancer. *Appl Physiol Nutr Metab* 2008;33:1269-83.

van Woudenberg GJ, Vliegenthart R, van Rooij FJ, Hofman A, Oudkerk M, Witteman JC, et al. Coffee consumption and coronary calcification: the Rotterdam Coronary Calcification Study. *Arterioscler Thromb Vasc Biol*. 2008;28:1018-23.

Wannamethee G, Shaper AG. Blood lipids: the relationship with alcohol intake, smoking, and body weight. *J Epidemiol Community Health* 1992;46:197-202.

Vartanian LR, Schwartz MB, Brownell KD. Effects of soft drink consumption on nutrition and health: a systematic review and meta-analysis. *Am J Public Health* 2007;97:667-75.

Watanabe M, Barzi F, Neal B, Ueshima H, Miyoshi Y, Okayama A, et al. Alcohol consumption and the risk of diabetes by body mass index levels in a cohort of 5,636 Japanese. *Diabetes Res Clin Pract* 2002;57:191-7.

Vatten LJ, Solvoll K, Loken EB. Coffee consumption and the risk of breast cancer. A prospective study of 14,593 Norwegian women. *Br J Cancer*. 1990;62:267-70.

Veitch J, Singh A, van Stralen MM, van Mechelen W, Brug J, Chinapaw MJ. Reduction in sugar-sweetened beverages is not associated with more water or diet drinks. *Public Health Nutr* 2011;14:1388-93.

Welsh J, Dietz W. Sugar-sweetened beverage consumption is associated with weight gain and incidence of type 2 diabetes. *Clinical Diabetes* 2005;23:150-152.

Welsh JA, Cogswell ME, Rogers S, Rockett H, Mei Z, Grummer-Strawn LM. Overweight among low-income preschool children associated with the consumption of sweet drinks: Missouri, 1999-2002. *Pediatrics* 2005;115:e223-9.

Vergnaud AC, Peneau S, Chat-Yung S, Kesse E, Czernichow S, Galan P, et al. Dairy consumption and 6-y changes in body weight and waist circumference in middle-aged French adults. *Am J Clin Nutr* 2008;88:1248-55.

Wilfley DE, Tibbs TL, Van Buren D, Reach KP, Walker MS, Epstein LH. Lifestyle interventions in the treatment of childhood overweight: A meta-analytic review of randomized controlled trials. *Health Psychology* 2007;26:521-532.

Williams PT. The relationships of vigorous exercise, alcohol, and adiposity to low and high high-density lipoprotein-cholesterol levels. *Metabolism*. 2004;53:700-9.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Wilson KM, Kasperzyk JL, Rider JR, Kenfield S, van Dam RM, Stampfer MJ, et al. Coffee consumption and prostate cancer risk and progression in the Health Professionals Follow-up Study. *J Natl Cancer Inst* 2011;103:876-84.

Wirfalt E, Hedblad B, Gullberg B, Mattisson I, Andren C, Rosander U, et al. Food patterns and components of the metabolic syndrome in men and women: a cross-sectional study within the Malmo Diet and Cancer cohort. *Am J Epidemiol* 2001;154:1150-9.

Wolff E, Dansinger ML. Soft drinks and weight gain: How strong is the link? *SOURCE MedGenMed Medscape General Medicine* (2008) 10:8 Article Number: 189. Date of Publication: 2008 2008.

Yang CC, Chiou WB. Substitution of healthy for unhealthy beverages among college students. A health-concerns and behavioral-economics perspective. *Appetite* 2010;54:512-6.

Yang G, Zheng W, Xiang YB, Gao J, Li HL, Zhang X, et al. Green tea consumption and colorectal cancer risk: a report from the Shanghai Men's Health Study. *Carcinogenesis* 2011;32:1684-8.

Yannakoulia M, Ntalla I, Papoutsakis C, Farmaki AE, Dedoussis GV. Consumption of vegetables, cooked meals, and eating dinner is negatively associated with overweight status in children. *Journal of Pediatrics* 2010;157:815-820.

Yoneda T, Shoji K, Takase H, Hibi M, Hase T, Meguro S, et al. Effectiveness and safety of 1-year ad libitum consumption of a high-catechin beverage under nutritional guidance. *Metab Syndr Relat Disord* 2009;7:349-56.

Yoo S, Nicklas T, Baranowski T, Zakeri IF, Yang SJ, Srinivasan SR, et al. Comparison of dietary intakes associated with metabolic syndrome risk factors in young adults: the Bogalusa Heart Study. *Am J Clin Nutr*. 2004;80:841-8.

Yoshita K, Miura K, Morikawa Y, Ishizaki M, Kido T, Naruse Y, et al. Relationship of alcohol consumption to 7-year blood pressure change in Japanese men. *J Hypertens* 2005;23:1485-90.

Zemel B, Shi H, Greer B, Dirienzo D, Zemel PC. Regulation of adiposity by dietary calcium. *FASEB J* 2000;14:1132-8.

Zemel MB. The role of dairy foods in weight management. *Journal of the American College of Nutrition* 2005;24:537S-546S.

Zemel MB, Richards J, Mathis S, Milstead A, Gebhardt L, Silva E. Dairy augmentation of total and central fat loss in obese subjects. *Int J Obes (Lond)* 2005;29:391-7.

Zemel MB, Thompson W, Zemel P, Nocton AM, Milstead A, Morris K, et al. Dietary calcium and dairy products accelerate weight and fat loss during energy restriction in obese adults. *American Journal of Clinical Nutrition* 2002:342s-343s.

Zhang M, Lee AH, Binns CW, Xie X. Green tea consumption enhances survival of epithelial ovarian cancer. *Int J Cancer* 2004;112:465-9.

Zheng XX, Xu YL, Li SH, Liu XX, Hui R, Huang XH. Green tea intake lowers fasting serum total and LDL cholesterol in adults: a meta-analysis of 14 randomized controlled trials. *Am J Clin Nutr* 2011;94:601-10

Kapitel 3.4

Amorim AR, Linne YM, Lourenco PM. Diet or exercise, or both, for weight reduction in women after childbirth. *Cochrane Database Syst Rev* 2007;CD005627.

Atlantis E, Barnes EH, Singh MA. Efficacy of exercise for treating overweight in children and adolescents: a systematic review. *Int J Obes (Lond)* 2006;30:1027-40.

Bennett B, Sothorn MS. Diet, exercise, behavior: the promise and limits of lifestyle change. *Semin Pediatr Surg* 2009;18:152-8.

Benson AC, Torode ME, Fiatarone Singh MA. Effects of resistance training on metabolic fitness in children and adolescents: a systematic review. *Obes Rev* 2008;9:43-66.

Bes-Rastrollo M, van Dam RM, Martinez-Gonzalez MA, Li TY, Sampson LL, Hu FB. Prospective study of dietary energy density and weight gain in women. *Am J Clin Nutr* 2008;88:769-77.

Boone-Heinonen J, Evenson KR, Taber DR, Gordon-Larsen P. Walking for prevention of cardiovascular disease in men and women: a systematic review of observational studies. *Obes Rev* 2009;10:204-17.

Brown T, Avenell A, Edmunds LD, Moore H, Whittaker V, Avery L, et al. Systematic review of long-term lifestyle interventions to prevent weight gain and morbidity in adults. *Obes Rev* 2009;10:627-38.

Caperchione CM, Kolt GS, Mummery WK. Physical activity in culturally and linguistically diverse migrant groups to Western society: a review of barriers, enablers and experiences. *Sports Med* 2009;39:167-77.

Chang SC, Ziegler RG, Dunn B, Stolzenberg-Solomon R, Lacey JV, Jr., Huang WY, et al. Association of energy intake and energy balance with postmenopausal breast cancer in the prostate, lung, colorectal, and ovarian cancer screening trial. *Cancer Epidemiol Biomarkers Prev.* 2006;15:334-41.

Conn VS, Hafdahl AR, Mehr DR, LeMaster JW, Brown SA, Nielsen PJ. Metabolic effects of interventions to increase exercise in adults with type 2 diabetes. *Diabetologia* 2007;50:913-921.

Dumler F. Dietary sodium intake and arterial blood pressure. *J Ren Nutr* 2009;19:57-60.

Durand CP, Andalib M, Dunton GF, Wolch J, Pentz MA. A systematic review of built environment factors related to physical activity and obesity risk: implications for smart growth urban planning. *Obes Rev* 2011;12:e173-82.

Fagard RH. Exercise is good for your blood pressure: Effects of endurance training and resistance training. *Clinical and Experimental Pharmacology and Physiology* 2006;33:853-856.

Faulkner GEJ, Buliung RN, Flora PK, Fusco C. Active school transport, physical activity levels and body weight of children and youth: A systematic review. *Preventive Medicine* 2009;48:3-8.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Flood A, Mitchell N, Jaeb M, Finch EA, Laqua PS, Welsh EM, et al. Energy density and weight change in a long-term weight-loss trial. *The International Journal of Behavioral Nutrition and Physical Activity* 2009;6.

Fogelholm M. Walking for the management of obesity. *Disease Management and Health Outcomes* 2005;13:9-18.

Giada F, Biffi A, Agostoni P, Anedda A, Belardinelli R, Carlon R, et al. Exercise prescription for the prevention and treatment of cardiovascular diseases: part I. *J Cardiovasc Med (Hagerstown)* 2008;9:529-44.

Hallal PC, Victora CG, Azevedo MR, Wells JCK. Adolescent physical activity and health: A systematic review. *Sports Medicine* 2006;36:1019-1030.

Harris KC, Kuramoto LK, Schulzer M, Retallack JE. Effect of school-based physical activity interventions on body mass index in children: a meta-analysis. *CMAJ* 2009;180:719-26.

He FJ, MacGregor GA. Reducing population salt intake worldwide: from evidence to implementation. *Prog Cardiovasc Dis* 2010;52:363-82.

Heymsfield SB, Harp JB, Reitman ML, Beetsch JW, Schoeller DA, Erondy N, et al. Why do obese patients not lose more weight when treated with low-calorie diets? A mechanistic perspective. *Am J Clin Nutr* 2007;85:346-54.

Hollenberg NK. The influence of dietary sodium on blood pressure. *Journal of the American College of Nutrition* 2006;25:240S-246S.

Huybrechts I, De Bourdeaudhuij I, De Henauw S. Environmental factors: opportunities and barriers for physical activity, and healthy eating among children and adolescents. *Verh K Acad Geneesk Belg* 2010;72:277-93.

Janssen I, LeBlanc AG. Systematic review of the health benefits of physical activity and fitness in school-aged children and youth. *SOURCE International Journal of Behavioral Nutrition and Physical Activity* (2010) 7 Article Number: 40. Date of Publication: 11 May 2010.

Jolly CA. Is dietary restriction beneficial for human health, such as for immune function? *Curr Opin Lipidol* 2007;18:53-7.

Kadoglou NPE, Iliadis F, Liapis CD. Exercise and Carotid Atherosclerosis. *European Journal of Vascular and Endovascular Surgery* 2008;35:264-272.

Kelly SA, Melnyk BM. Systematic review of multicomponent interventions with overweight middle adolescents: implications for clinical practice and research. *Worldviews Evid Based Nurs* 2008;5:113-35.

Lachowycz K, Jones AP. Greenspace and obesity: a systematic review of the evidence. *Obes Rev* 2011;12:e183-9.

Lovasi GS, Hutson MA, Guerra M, Neckerman KM. Built environments and obesity in disadvantaged populations. *Epidemiologic Reviews* 2009;31:7-20.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Orozco LJ, Buchleitner AM, Gimenez-Perez G, Figuls MR, Richter B, Mauricio D. Exercise or exercise and diet for preventing type 2 diabetes mellitus. SOURCE Cochrane Database of Systematic Reviews (2008) :3 Article Number: CD003054. Date of Publication: 2008.

Pescatello LS. Exercise and hypertension: recent advances in exercise prescription. *Curr Hypertens Rep* 2005;7:281-6.

Pichon Riviere A, Augustovski F, Garcia Marti S, Aruj P, Buendia J, Galante J, et al. Physical activity for the treatment of morbid obesity (Structured abstract). Ciudad de Buenos Aires: Institute for Clinical Effectiveness and Health Policy (IECS) 2009.

Rhodes RE, Warburton DER, Murray H. Characteristics of physical activity guidelines and their effect on adherence: A review of randomized trials. *Sports Medicine* 2009;39:355-375.

Silvera SA, Jain M, Howe GR, Miller AB, Rohan TE. Energy balance and breast cancer risk: a prospective cohort study. *Breast Cancer Res Treat.* 2006;97:97-106.

Soderlund A, Fischer A, Johansson T. Physical activity, diet and behaviour modification in the treatment of overweight and obese adults: a systematic review. *Perspect Public Health* 2009;129:132-42.

Summerbell CD, Douthwaite W, Whittaker V, Ells LJ, Hillier F, Smith S, et al. The association between diet and physical activity and subsequent excess weight gain and obesity assessed at 5 years of age or older: a systematic review of the epidemiological evidence. *Int J Obes (Lond)* 2009;33 Suppl 3:S1-92.

Tambalis K, Panagiotakos DB, Kavouras SA, Sidossis LS. Responses of blood lipids to aerobic, resistance, and combined aerobic with resistance exercise training: A systematic review of current evidence. *Angiology* 2009;60:614-632.

Tsai AG, Wadden TA. The evolution of very-low-calorie diets: an update and meta-analysis. *Obesity (Silver Spring)* 2006;14:1283-93.

Umpierre D, Ribeiro PA, Kramer CK, Leitao CB, Zucatti AT, Azevedo MJ, et al. Physical activity advice only or structured exercise training and association with HbA1c levels in type 2 diabetes: a systematic review and meta-analysis. *JAMA* 2011;305:1790-9.

Weinheimer EM, Sands LP, Campbell WW. A systematic review of the separate and combined effects of energy restriction and exercise on fat-free mass in middle-aged and older adults: implications for sarcopenic obesity. *Nutr Rev* 2010;68:375-88.

Welch G, Wesolowski C, Piepul B, Kuhn J, Romanelli J, Garb J. Physical activity predicts weight loss following gastric bypass surgery: findings from a support group survey. *Obes Surg* 2008;18:517-24.

Yates T, Khunti K, Bull F, Gorely T, Davies MJ. The role of physical activity in the management of impaired glucose tolerance: A systematic review. *Diabetologia* 2007;50:1116-1126.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Kapitel 5

Kostnadseffektivitet

Burns C, Sacks G, Rayner M, Bilenkij G, Swinburn B. Correctly calculating the cost of food. *Nutr Rev* 2010;68:182-3; author reply 184-5.

Cecchini M, Sassi F, Lauer JA, Lee YY, Guajardo-Barron V, Chisholm D. Tackling of unhealthy diets, physical inactivity, and obesity: health effects and cost-effectiveness. *Lancet* 2010;376:1775-84.

Cobiac L, Vos T, Veerman L. Cost-effectiveness of Weight Watchers and the Lighten Up to a Healthy Lifestyle program. *Aust N Z J Public Health* 2010;34:240-7.

Continuing medical education program: A comparison of diet and exercise therapy versus laparoscopic Roux-en-Y gastric bypass surgery for morbid obesity: A decision analysis model; A cost-effectiveness analysis of intraoperative cholangiography in the prevention of bile duct injury during laparoscopic cholecystectomy. *Journal of the American College of Surgeons* 2003;196:501-503.

Cost-effectiveness of diet and exercise interventions to reduce overweight and obesity. Forster, M., Veerman, J.L., Barendregt, J.J. and Vos, T., *International Journal of Obesity*, 35 (2011) 1071-8. (<http://www.ncbi.nlm.nih.gov/pubmed/21224825>)

Cost-effectiveness of exercise and diet in overweight and obese adults with knee osteoarthritis. Sevick, M.A., Miller, G.D., Loeser, R.F., Williamson, J.D. and Messier, S.P. *Medicine and Science in Sports and Exercise*, 41 (2009) 1167-74. (<http://www.ncbi.nlm.nih.gov/pubmed/19461553>)

Costs of Mediterranean and western dietary patterns in a Spanish cohort and their relationship with prospective weight change. Lopez, C.N., Martinez-Gonzalez, M.A., Sanchez-Villegas, A., Alonso, A., Pimenta, A.M. and Bes-Rastrollo, M., *J Epidemiol Community Health*, 63 (2009) 920-7.

Drewnowski A, Eichelsdoerfer P. The Mediterranean diet: does it have to cost more? *Public Health Nutr* 2009;12:1621-8.

Finkelstein EA, Ruhm CJ, Kosa KM. Economic causes and consequences of obesity. *Annu Rev Public Health* 2005;26:239-57.

Flynn TJ, Walsh MF. Thirty-month evaluation of a popular very-low-calorie diet program. *Arch Fam Med* 1993;2:1042-8.

Fry J, Finley W. The prevalence and costs of obesity in the EU. *Proc Nutr Soc* 2005;64:359-62.

Jensen C, Flum DR. The costs of nonsurgical and surgical weight loss interventions: is an ounce of prevention really worth a pound of cure? (Structured abstract). *Surgery for Obesity and Related Diseases* 2005:353-357.

Johannesson M, Fagerberg B. A health-economic comparison of diet and drug treatment in obese men with mild hypertension. *J Hypertens* 1992;10:1063-70.

John J, Wenig CM, Wolfenstetter SB. Recent economic findings on childhood obesity: cost-of-illness and cost-effectiveness of interventions. *Curr Opin Clin Nutr Metab Care* 2010;13:305-13.

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Kouris-Blazos A, Wahlqvist ML. Health economics of weight management: evidence and cost. *Asia Pac J Clin Nutr* 2007;16 Suppl 1:329-38.

Lopez CN, Martinez-Gonzalez MA, Sanchez-Villegas A, Alonso A, Pimenta AM, Bes-Rastrollo M. Costs of Mediterranean and western dietary patterns in a Spanish cohort and their relationship with prospective weight change. *Journal of Epidemiology and Community Health* 2009;63:920-927.

McAuley KA, Taylor RW, Farmer VL, Hansen P, Williams SM, Booker CS, et al. Economic evaluation of a community-based obesity prevention program in children: the APPLE project (Provisional abstract). *Obesity* 2010:131-136.

McCarron DA, Heaney RP. Estimated healthcare savings associated with adequate dairy food intake. *Am J Hypertens* 2004;17:88-97.

Meenan RT, Vogt TM, Williams AE, Stevens VJ, Albright CL, Nigg C. Economic evaluation of a worksite obesity prevention and intervention trial among hotel workers in Hawaii. *Journal of Occupational and Environmental Medicine* 2010;52:S8-S13.

Olsen J, Willaing I, Ladelund S, Jorgensen T, Gundgaard J, Sorensen J. Cost-effectiveness of nutritional counseling for obese patients and patients at risk of ischemic heart disease. *Int J Technol Assess Health Care* 2005;21:194-202.

Pacy PJ, Webster JD, Pearson M, Garrow JS. A cross-sectional cost/benefit audit in a hospital obesity clinic. *Hum Nutr Appl Nutr* 1987;41:38-46.

Raffensperger JF. The least-cost low-carbohydrate diet is expensive. *Nutr Res* 2008;28:6-12.

Raynor HA, Kilanowski CK, Esterlis I, Epstein LH. A cost-analysis of adopting a healthful diet in a family-based obesity treatment program. *J Am Diet Assoc* 2002;102:645-56.

Roux L, Kuntz KM, Donaldson C, Goldie SJ. Economic evaluation of weight loss interventions in overweight and obese women. *Obesity* 2006;14:1093-1106.

Schroder H, Marrugat J, Covas MI. High monetary costs of dietary patterns associated with lower body mass index: a population-based study. *Int J Obes (Lond)* 2006;30:1574-9.

Sevick MA, Miller GD, Loeser RF, Williamson JD, Messier SP. Cost-effectiveness of exercise and diet in overweight and obese adults with knee osteoarthritis. *Med Sci Sports Exerc* 2009;41:1167-74.

Van Baal PHM, Van Den Berg M, Hoogenveen RT, Vijgen SMC, Engelfriet PM. Cost-effectiveness of a low-calorie diet and orlistat for obese persons: Modeling long-term health gains through prevention of obesity-related chronic diseases. *Value in Health* 2008;11:1033-1040.

Wake M, Baur LA, Gerner B, Gibbons K, Gold L, Gunn J, et al. Outcomes and costs of primary care surveillance and intervention for overweight or obese children: the LEAP 2 randomised controlled trial (Structured abstract). *BMJ* 2009.

Wake M, Gold L, McCallum Z, Gerner B, Waters E. Economic evaluation of a primary care trial to reduce weight gain in overweight/obese children: the LEAP trial. *Ambul Pediatr* 2008;8:336-41.

Samband BMI/Vikt

Författare, år	Från sökning	Orsak till exkludering
Andreyeva, 2011	Taxation	Tvärsnittsstudier
Asfaw, 2006	Econlit	Tvärsnittsstudier
Asfaw, 2006b	Econlit	Tvärsnittsstudier
Beydoun, 2008		Tvärsnittsstudier
Boizot, 2010	Econlit	Ej refereegranskad
Bonnet, 2011	Taxation	Ej relevant
Brooks, 2010	Taxation	Ej relevant
Cash, 2007	Econlit	Översiktsartikel
Chen, 2002	Econlit	Modellstudie
Chouinard, 2007	Econlit	Modellstudie
Cotti, 2012	Taxation	Tvärsnittsstudier
Couch, 2011		Debattartikel
Dharmesena, 2011		Tvärsnittsstudier
Epstein, 2012	Taxation	Systematisk översikt med tvärsnittsstudier och studier i laboratoriemiljö
Finkelstein, 2010	Taxation	Tvärsnittsstudier
Fletcher, 2010a	Econlit/Taxation	Tvärsnittsstudier
Fletcher, 2010b	Econlit	Tvärsnittsstudier
Fletcher, 2010c		Översiktsartikel
Fletcher, 2011	Econlit, Taxation	Tvärsnittsstudier
Gandal, 2010	Econlit	Tvärsnittsstudier
Giesen, 2011	Taxation	Laboratoriemiljö
Giesen, 2012	Taxation	Laboratoriemiljö
Gordon-Larsen, 2011	Taxation	Ej relevant
Goldman, 2008		Ej peer-reviewed
Gustavsen, 2011	Econlit	Tvärsnittsstudier
Horgren, 2002		Ej relevant
Lacanilao, 2011	Econlit	Ej relevant
Kennedy, 2010	Econlit	Modellstudie
Khan, 2012	Taxation	Ej relevant
Kimbrow, 2010	Taxation	Ej relevant
Kotakorpi, 2011	Econlit	Tvärsnittsstudier
Kuchler, 2005	Econlit	Tvärsnittsstudier
Lin, 2011	Taxation	Modellstudie
Lordan, 2011	Econlit	Ej relevant
Lu, 2010	Taxation	Tvärsnittsstudier
Maddock, 2010	Taxation	Översiktsartikel
Miao, 2011	Econlit	Modellstudie
Meyerhoefer, 2010	Econlit	Tvärsnittsstudier
Nnoaham, 2009	Taxation	Modellstudie
Nordström, 2011a	Econlit	Tvärsnittsstudier
Nordström, 2011b	Econlit	Modellstudie
Powell, 2007		Tvärsnittsstudier
Powell, 2009a	Econlit	Tvärsnittsstudier
Powell, 2009b	Econlit/Taxation	Ej relevant
Powell, 2009d	Taxation	Tvärsnittsstudier
Powell, 2009e	Taxation	Översiktsartikel
Powell, 2011	Econlit	Tvärsnittsstudier
Sacks, 2011	Taxation	Modellstudie
Schroeter, 2008	Econlit	Modellstudie
Smed, 2007	Econlit	Tvärsnittsstudier
Staudigel, 2010	Econlit/Taxation	Tvärsnittsstudier
Sturm, 2010	Taxation	Tvärsnittsstudier

SBU. Mat vid fetma. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2013. SBU-rapport nr 218.

Sturm, 2011	Taxation	Tvärsnittsstudier
Thow, 2010	Taxation	Översiktsartikel
Tiffin, 2011	Taxation	Modellstudie
Tobey		Översiktsartikel
Wang, 2012	Econlit	Modellstudie
Yaniv, 2009	Econlit	Modellstudie
Zheng, 2008	Econlit	Tvärsnittsstudier