



Bilaga 4 Studier som bedömts ha låg kvalitet samt studier som bedömts vara ej relevanta

Ej relevanta studier. Kohort- samt fall-kontrolldesign

Albert HB, Godskesen M, Korsholm L, Westergaard JG. Risk factors in developing pregnancy-related pelvic girdle pain. *Acta Obstet Gynecol Scand.* 2006;85(5):539-44.

Biering-Sorensen F. A prospective study of low back pain in a general population. I. Occurrence, recurrence and aetiology. *Scand J Rehabil Med.* 1983;15(2):71-9.

Biering-Sorensen F, Thomsen C. Medical, social and occupational history as risk indicators for low-back trouble in a general population. *Spine (Phila Pa 1976).* 1986;11(7):720-5.

Bildt C, Alfredsson L, Punnett L, Theobald H, Torgen M, Wikman A. Effects of drop out in a longitudinal study of musculoskeletal disorders. *Occup Environ Med.* 2001;58(3):194-9.

Bjorksten MG, Talback M. A follow-up study of psychosocial factors and musculoskeletal problems among unskilled female workers with monotonous work. *Eur J Public Health.* 2001;11(1):102-8.

Boos N, Rieder R, Schade V, Spratt KF, Semmer N, Aebi M. 1995 Volvo Award in clinical sciences. The diagnostic accuracy of magnetic resonance imaging, work perception, and psychosocial factors in identifying symptomatic disc herniations. *Spine (Phila Pa 1976).* 1995;20(24):2613-25.

Bridger RS, Brasher K, Bennett A. Sustaining person-environment fit with a changing workforce. *Ergonomics.* 2012.

Brynhildsen J, Hansson A, Persson A, Hammar M. Follow-up of patients with low back pain during pregnancy. *Obstet Gynecol.* 1998;91(2):182-6.

Canivet C, Ostergren PO, Choi B, Nilsson P, af Sillen U, Moghadassi M, et al. Sleeping problems as a risk factor for subsequent musculoskeletal pain and the role of job strain: results from a one-year follow-up of the Malmo Shoulder Neck Study Cohort. *International journal of behavioral medicine.* 2008;15(4):254-62.

Daltroy LH, Larson MG, Wright EA, Malspeis S, Fossel AH, Ryan J, et al. A case-control study of risk factors for industrial low back injury: implications for primary and secondary prevention programs. *Am J Ind Med.* 1991;20(4):505-15.

de Zwart BC, Broersen JP, van der Beek AJ, Frings-Dresen MH, Van Dijk FJ. Selection related to musculoskeletal complaints among employees. *Occup Environ Med*. 1997;54(11):800-6.

Elfering A, Mannion AF, Jacobshagen N, Tamcan O, Muller U. Beliefs about back pain predict the recovery rate over 52 consecutive weeks. *Scand J Work Environ Health*. 2009;35(6):437-45.

Engkvist IL. Back injuries among nurses - A comparison of the accident processes after a 10-year follow-up. *Safety Science*. 2008;46(2):291-301.

Engkvist IL, Hagberg M, Hjelm EW, Menckel E, Ekenvall L. The accident process preceding overexertion back injuries in nursing personnel. PROSA study group. *Scand J Work Environ Health*. 1998;24(5):367-75.

Engkvist IL, Hjelm EW, Hagberg M, Menckel E, Ekenvall L. Risk indicators for reported over-exertion back injuries among female nursing personnel. *Epidemiology (Cambridge, Mass)*. 2000;11(5):519-22.

Ferguson SA, Allread WG, Burr DL, Heaney C, Marras WS. Biomechanical, psychosocial and individual risk factors predicting low back functional impairment among furniture distribution employees. *Clinical Biomechanics*. 2012;27(2):117-23.

Ferguson SA, Marras WS, Burr DL. The influence of individual low back health status on workplace trunk kinematics and risk of low back disorder. *Ergonomics*. 2004;47(11):1226-37.

Gatchel RJ, Polatin PB, Mayer TG. The dominant role of psychosocial risk factors in the development of chronic low back pain disability. *Spine (Phila Pa 1976)*. 1995;20(24):2702-9.

Hartvigsen J, Bakketeig LS, Leboeuf-Yde C, Engberg M, Lauritzen T. [The association between physical workload and low back pain clouded by the "healthy worker" effect]. *Ugeskr Laeger*. 2002;164(21):2765-8.

Hellsing A, Bryngelsson I. Predictors of musculoskeletal pain in men: a twenty-year follow-up from examination at enlistment. *Spine (Phila Pa 1976)*. 2000;25(23):3080-6.

Holmberg S, Thelin A, Stiernstrom EL, Svardsudd K. The impact of physical work exposure on musculoskeletal symptoms among farmers and rural non-farmers. A population-based study. *Annals of Agricultural and Environmental Medicine*. 2003;10(2):179-84.

- Holmberg S, Thelin A, Stiernstrom EL, Svardsudd K. Low back pain comorbidity among male farmers and rural referents: a population-based study. *Ann Agric Environ Med.* 2005;12(2):261-8.
- Holtermann A, Blangsted AK, Hansen K, Christensen H, Sogaard K. What characterizes cleaners sustaining good musculoskeletal health after years with physically heavy work? *International Archives of Occupational and Environmental Health.* 2009;82(8):1015-22.
- Hägg O, Fritzell P, Nordwall A. Characteristics of patients with chronic low back pain selected for surgery: a comparison with the general population reported from the Swedish Lumbar Spine Study including commentary by Fanuele J. *Spine (Phila Pa 1976).* 2002;27(11):1223-31.
- Josephson M, Ahlberg G, Harenstam A, Svensson H, Theorell T, Wiktorin C, et al. Paid and unpaid work, and its relation to low back and neck/shoulder disorders among women. *Women & health.* 2003;37(2):17-30.
- Juhl M, Andersen PK, Olsen J, Andersen AM. Psychosocial and physical work environment, and risk of pelvic pain in pregnancy. A study within the Danish national birth cohort. *J Epidemiol Community Health.* 2005;59(7):580-5.
- Keeney BJ, Turner JA, Fulton-Kehoe D, Wickizer TM, Chan KC, Franklin GM. Early predictors of occupational back reinjury: results from a prospective study of workers in Washington State. *Spine (Phila Pa 1976).* 2013;38(2):178-87.
- Kishi R, Doi R, Fukuchi Y, Satoh H, Satoh T, Ono A, et al. Subjective symptoms and neurobehavioral performances of ex-mercury miners at an average of 18 years after the cessation of chronic exposure to mercury vapor. *Mercury Workers Study Group. Environmental research.* 1993;62(2):289-302.
- Kuh DJ, Coggan D, Mann S, Cooper C, Yusuf E. Height, occupation and back pain in a national prospective study. *Br J Rheumatol.* 1993;32(10):911-6.
- Kuijjer PP, van der Beek AJ, van Dieen JH, Visser B, Frings-Dresen MH. Effect of job rotation on need for recovery, musculoskeletal complaints, and sick leave due to musculoskeletal complaints: a prospective study among refuse collectors. *Am J Ind Med.* 2005;47(5):394-402.
- Lin MY, Ahern JE, Gershon RR, Grimes M. The use of total quality improvement techniques to determine risk factors for back injuries in hospital workers. *Clinical performance and quality health care.* 1998;6(1):23-7.

- Lotters F, Burdorf A, Kuiper J, Miedema H. Model for the work-relatedness of low-back pain. *Scand J Work Environ Health*. 2003;29(6):431-40.
- Manninen P, Heliovaara M, Riihimaki H, Makela P. Does psychological distress predict disability? *Int J Epidemiol*. 1997;26(5):1063-70.
- Marras WS, Lavender SA, Ferguson SA, Splittstoesser RE, Yang G. Quantitative dynamic measures of physical exposure predict low back functional impairment. *Spine (Phila Pa 1976)*. 2010;35(8):914-23.
- Masset DF, Piette AG, Malchaire JB. Relation between functional characteristics of the trunk and the occurrence of low back pain. Associated risk factors. *Spine (Phila Pa 1976)*. 1998;23(3):359-65.
- Melloh M, Salathe CR, Elfering A, Kaser A, Barz T, Aghayev E, et al. Occupational, personal and psychosocial resources for preventing persistent low back pain. *Int J Occup Saf Ergon*. 2013;19(1):29-40.
- Messing K, Stock SR, Tissot F. Should studies of risk factors for musculoskeletal disorders be stratified by gender? Lessons from the 1998 Quebec Health and Social Survey. *Scand J Work Environ Health*. 2009;35(2):96-112.
- Miwa S, Yokogawa A, Kobayashi T, Nishimura T, Igarashi K, Inatani H, et al. Risk factors of recurrent lumbar disc herniation: A single center study and review of the literature. *Journal of Spinal Disorders and Techniques*. 2012;14.
- Nyman T, Mulder M, Iliadou A, Svartengren M, Wiktorin C. Physical workload, low back pain and neck-shoulder pain: a Swedish twin study. *Occupational and Environmental Medicine [Internet]*. 2009; (6):[395-401 pp.]. Available from: <http://www.mrw.interscience.wiley.com/cochrane/clcentral/articles/243/CN-00713243/frame.html>.
- Papageorgiou AC, Croft PR, Thomas E, Silman AJ, Macfarlane GJ. Psychosocial risks for low back pain: are these related to work? *Ann Rheum Dis*. 1998;57(8):500-2.
- Piterman L, Dunt D. Occupational lower-back injuries in a primary medical care setting: a five-year follow-up study. *Med J Aust*. 1987;147(6):276-9.
- Riihimaki H, Wickstrom G, Hanninen K, Luopajarvi T. Predictors of sciatic pain among concrete reinforcement workers and house painters--a five-year follow-up. *Scand J Work Environ Health*. 1989;15(6):415-23.
- Riyazi N, Rosendaal FR, Slagboom E, Kroon HM, Breedveld FC, Kloppenburg M. Risk factors in familial osteoarthritis: the GARP sibling study. *Osteoarthritis Cartilage*. 2008;16(6):654-9.

Låg kvalitet. Studier av kohort- samt fall-kontrolldesign

Bergenudd H, Johnell O. Somatic versus nonsomatic shoulder and back pain experience in middle age in relation to body build, physical fitness, bone mineral content, gamma-glutamyltransferase, occupational workload, and psychosocial factors. *Spine (Phila Pa 1976)*. 1991;16(9):1051-5.

Bigos SJ, Battie MC, Spengler DM, Fisher LD, Fordyce WE, Hansson T, et al. A longitudinal, prospective study of industrial back injury reporting. *Clinical orthopaedics and related research*. 1992(279):21-34.

Bigos SJ, Battie MC, Spengler DM, Fisher LD, Fordyce WE, Hansson TH, et al. A prospective study of work perceptions and psychosocial factors affecting the report of back injury. *Spine (Phila Pa 1976)*. 1991;16(1):1-6.

Boos N, Semmer N, Elfering A, Schade V, Gal I, Zanetti M, et al. Natural history of individuals with asymptomatic disc abnormalities in magnetic resonance imaging: predictors of low back pain-related medical consultation and work incapacity. *Spine (Phila Pa 1976)*. 2000;25(12):1484-92.

Clays E, De Bacquer D, Leynen F, Kornitzer M, Kittel F, De Backer G. The impact of psychosocial factors on low back pain: longitudinal results from the Belstress study. *Spine (Phila Pa 1976)*. 2007;32(2):262-8.

Coupland CA, Grainge MJ, Cliffe SJ, Hosking DJ, Chilvers CE. Occupational activity and bone mineral density in postmenopausal women in England. *Osteoporos Int*. 2000;11(4):310-5.

de Zwart BC, Broersen JP, Frings-Dresen MH, van Dijk FJ. Repeated survey on changes in musculoskeletal complaints relative to age and work demands. *Occup Environ Med*. 1997;54(11):793-9.

Elfering A, Semmer N, Birkhofer D, Zanetti M, Hodler J, Boos N. Risk factors for lumbar disc degeneration: a 5-year prospective MRI study in asymptomatic individuals. *Spine (Phila Pa 1976)*. 2002;27(2):125-34.

Elfering A, Semmer NK, Schade V, Grund S, Boos N. Supportive colleague, unsupportive supervisor: the role of provider-specific constellations of social support at work in the development of low back pain. *Journal of occupational health psychology*. 2002;7(2):130-40.

Eriksen W, Natvig B, Bruusgaard D. Smoking, heavy physical work and low back pain: a four-year prospective study. *Occup Med (Lond)*. 1999;49(3):155-60.

Gnudi S, Sitta E, Gnudi F, Pignotti E. Relationship of a lifelong physical workload with physical function and low back pain in retired women. *Aging Clinical & Experimental Research*. 2009;21(1):55-61.

Holtermann A, Clausen T, Aust B, Mortensen OS, Andersen LL. Risk for low back pain from different frequencies, load mass and trunk postures of lifting and carrying among female healthcare workers. *Int Arch Occup Environ Health*. 2012.

Holtermann A, Clausen T, Aust B, Mortensen OS, Andersen LL. Does occupational lifting and carrying among female health care workers contribute to an escalation of pain-day frequency? *Eur J Pain*. 2013;17(2):290-6.

Jensen A, Kaerlev L, Tuchsén F, Hannerz H, Dahl S, Nielsen PS, et al. Locomotor diseases among male long-haul truck drivers and other professional drivers. *Int Arch Occup Environ Health*. 2008;81(7):821-7.

Jensen JC, Haahr JP, Frost P, Andersen JH. Do work-related factors affect care-seeking in general practice for back pain or upper extremity pain? *Int Arch Occup Environ Health*. 2012.

Josephson M, Lagerstrom M, Hagberg M, Wigaeus Hjelm E. Musculoskeletal symptoms and job strain among nursing personnel: a study over a three year period. *Occup Environ Med*. 1997;54(9):681-5.

Kaaria SM, Malkia EA, Luukkonen RA, Leino-Arjas PI. Pain and clinical findings in the low back: a study of industrial employees with 5-, 10-, and 28-year follow-ups. *Eur J Pain*. 2010;14(7):759-63.

Kelsey JL, Githens PB, White AA, 3rd, Holford TR, Walter SD, O'Connor T, et al. An epidemiologic study of lifting and twisting on the job and risk for acute prolapsed lumbar intervertebral disc. *Journal of orthopaedic research : official publication of the Orthopaedic Research Society*. 1984;2(1):61-6.

Knox JB, Orchowski JR, Scher DL, Owens BD, Burks R, Belmont PJ, Jr. Occupational driving as a risk factor for low back pain in active-duty military service members. *Spine J*. 2013.

Lau B, Knardahl S. Perceived job insecurity, job predictability, personality, and health. *J Occup Environ Med*. 2008;50(2):172-81.

Lipscomb HJ, Loomis D, McDonald MA, Kucera K, Marshall S, Li L. Musculoskeletal symptoms among commercial fishers in North Carolina. *Appl Ergon*. 2004;35(5):417-26.

Lonnberg F, Pedersen PA, Siersma V. Early predictors of the long-term outcome of low back pain--results of a 22-year prospective cohort study. *Family practice*. 2010;27(6):609-14.

Massaccesi M, Pagnotta A, Soccetti A, Masali M, Masiero C, Greco F. Investigation of work-related disorders in truck drivers using RULA method. *Appl Ergon*. 2003;34(4):303-7.

Melloh M, Elfering A, Stanton TR, Kaser A, Salathe CR, Barz T, et al. Who is likely to develop persistent low back pain? A longitudinal analysis of prognostic occupational factors. *Work*. 2013;46(3):297-311.

Miedema HS, Chorus AM, Wevers CW, van der Linden S. Chronicity of back problems during working life. *Spine (Phila Pa 1976)*. 1998;23(18):2021-8; discussion 8-9.

Milosavljevic S, Bagheri N, Vasiljev RM, McBride DI, Rehn B. Does daily exposure to whole-body vibration and mechanical shock relate to the prevalence of low back and neck pain in a rural workforce? *Ann Occup Hyg*. 2012;56(1):10-7.

Nuwayhid IA, Stewart W, Johnson JV. Work activities and the onset of first-time low back pain among New York City fire fighters. *Am J Epidemiol*. 1993;137(5):539-48.

Palmer KT, Griffin M, Ntani G, Shambrook J, McNee P, Sampson M, et al. Professional driving and prolapsed lumbar intervertebral disc diagnosed by magnetic resonance imaging: a case-control study. *Scand J Work Environ Health*. 2012;38(6):577-81.

Palmer KT, Harris CE, Griffin MJ, Bennett J, Reading I, Sampson M, et al. Case-control study of low-back pain referred for magnetic resonance imaging, with special focus on whole-body vibration. *Scand J Work Environ Health*. 2008;34(5):364-73.

Plouvier S, Leclerc A, Chastang JF, Bonenfant S, Goldberg M. Socioeconomic position and low-back pain--the role of biomechanical strains and psychosocial work factors in the GAZEL cohort. *Scand J Work Environ Health*. 2009;35(6):429-36.

Plouvier S, Renahy E, Chastang JF, Bonenfant S, Leclerc A. Biomechanical strains and low back disorders: quantifying the effects of the number of years of exposure on various types of pain. *Occup Environ Med*. 2008;65(4):268-74.

Power C, Frank J, Hertzman C, Schierhout G, Li L. Predictors of low back pain onset in a prospective British study. *Am J Public Health*. 2001;91(10):1671-8.

Reigo T, Tropp H, Timpka T. Absence of back disorders in adults and work-related predictive factors in a 5-year perspective. *Eur Spine J*. 2001;10(3):215-20; discussion 21.

- Riihimaki H, Viikari-Juntura E, Moneta G, Kuha J, Videman T, Tola S. Incidence of sciatic pain among men in machine operating, dynamic physical work, and sedentary work. A three-year follow-up. *Spine (Phila Pa 1976)*. 1994;19(2):138-42.
- Rohrer MH, Santos-Eggimann B, Paccaud F, Haller-Maslov E. Epidemiologic study of low back pain in 1398 Swiss conscripts between 1985 and 1992. *Eur Spine J*. 1994;3(1):2-7.
- Roy TC, Lopez HP, Piva SR. Loads Worn by Soldiers Predict Episodes of Low Back Pain during Deployment to Afghanistan. *Spine (Phila Pa 1976)*. 2013.
- Rugulies R, Krause N. Effort-reward imbalance and incidence of low back and neck injuries in San Francisco transit operators. *Occup Environ Med*. 2008;65(8):525-33.
- Stevenson JM, Weber CL, Smith JT, Dumas GA, Albert WJ. A longitudinal study of the development of low back pain in an industrial population. *Spine (Phila Pa 1976)*. 2001;26(12):1370-7.
- Yassi A, Khokhar J, Tate R, Cooper J, Snow C, Vallentyne S. The epidemiology of back injuries in nurses at a large Canadian tertiary care hospital: implications for prevention. *Occup Med (Lond)*. 1995;45(4):215-20.
- Zwerling C, Ryan J, Schootman M. A case-control study of risk factors for industrial low back injury. The utility of preplacement screening in defining high-risk groups. *Spine (Phila Pa 1976)*. 1993;18(9):1242-7.

Ej relevanta studier. Tvärsnittsdesign

Aydog ST, Turbedar E, Demirel AH, Tetik O, Akin A, Doral MN. Cervical and lumbar spinal changes diagnosed in four-view radiographs of 732 military pilots. *Aviat Space Environ Med.* 2004;75(2):154-7.

Bartels S, Niederman B, Waters TR. Job hazards for musculoskeletal disorders for youth working on farms. *J Agric Saf Health.* 2000;6(3):191-201.

Benach J, Gimeno D, Benavides FG, Martinez JM, Torne Mdel M. Types of employment and health in the European union: changes from 1995 to 2000. *Eur J Public Health.* 2004;14(3):314-21.

Benavides FG, Benach J, Diez-Roux AV, Roman C. How do types of employment relate to health indicators? Findings from the second European survey on working conditions. *J Epidemiol Community Health.* 2000;54(7):494-501.

Bern SH, Brauer C, Moller KL, Koblauch H, Thygesen LC, Simonsen EB, et al. Baggage handler seniority and musculoskeletal symptoms: is heavy lifting in awkward positions associated with the risk of pain? *BMJ Open.* 2013;3(11):e004055.

Brinckmann P, Frobin W, Biggeman M, Tillotson M, Burton K. Quantification of overload injuries to thoracolumbar vertebrae and discs in persons exposed to heavy physical exertions or vibration at the workplace. Part II. Occurrence and magnitude of overload injury in exposed cohorts. *Clinical Biomechanics.* 1998;13(SUPPL. 2):I-S36.

Brown JJ, Wells GA, Trottier AJ, Bonneau J, Ferris B. Back pain in a large Canadian police force. *Spine (Phila Pa 1976).* 1998;23(7):821-7.

Burton AK, Tillotson KM, Troup JD. Prediction of low-back trouble frequency in a working population. *Spine (Phila Pa 1976).* 1989;14(9):939-46.

Byrns G, Agnew J, Curbow B. Attributions, stress, and work-related low back pain. *Appl Occup Environ Hyg.* 2002;17(11):752-64.

Cherry N. Women and work stress: Evidence from the 1946 birth cohort. *Ergonomics.* 1984;27(5):519-26.

Christensen H, Pedersen MB, Sjogaard G. A national cross-sectional study in the Danish wood and furniture industry on working postures and manual materials handling. *Ergonomics.* 1995;38(4):793-805.

Daniels C, Huang GD, Feuerstein M, Lopez M. Self-report measure of low back-related biomechanical exposures: clinical validation. *J Occup Rehabil.* 2005;15(2):113-28.

D'Arcy LP, Sasai Y, Stearns SC. Do assistive devices, training, and workload affect injury incidence? Prevention efforts by nursing homes and back injuries among nursing assistants. *Journal of Advanced Nursing.* 2012;68(4):836-45.

Dennerlein JT, Hopcia K, Sembajwe G, Kenwood C, Stoddard AM, Tveito TH, et al. Ergonomic practices within patient care units are associated with musculoskeletal pain and limitations. *Am J Ind Med.* 2012;55(2):107-16.

Dosoglu M, Yldz U, Is M, Orhan Z. Low back pain among nurses: A review. *Neurosurgery Quarterly.* 2009;19(1):29-34.

Duquette J, Lortie M, Rossignol M. Perception of difficulties for the back related to assembly work: general findings and impact of back health. *Appl Ergon.* 1997;28(5-6):389-96.

Elfering A, Grebner S, Gerber H, Semmer NK. Workplace observation of work stressors, catecholamines and musculoskeletal pain among male employees. *Scand J Work Environ Health.* 2008;34(5):337-44.

Elovainio M, Sinervo T. Psychosocial stressors at work, psychological stress and musculoskeletal symptoms in the care for the elderly. *Work & Stress.* 1997;11(4):351-61.

Erdinc O. Upper extremity musculoskeletal discomfort among occupational notebook personal computer users: Work interference, associations with risk factors and the use of notebook computer stand and docking station. *Work: Journal of Prevention, Assessment & Rehabilitation.* 2011;39(4):455-63.

Fathallah FA, Marras WS, Parnianpour M. The role of complex, simultaneous trunk motions in the risk of occupation-related low back disorders. *Spine.* 1998;23(9):1035-42.

Feuerstein M, Harrington CB, Lopez M, Haufler A. How do job stress and ergonomic factors impact clinic visits in acute low back pain? A prospective study. *J Occup Environ Med.* 2006;48(6):607-14.

Feuerstein M, Sult S, Houle M. Environmental stressors and chronic low back pain: Life events, family and work environment. *Pain.* 1985;22(3):295-307.

Frymoyer JW, Pope MH, Costanza MC. Epidemiologic studies of low-back pain. *Spine*. 1980;5(5):419-23.

Geiger-Brown J, Trinkoff AM, Nielsen K, Lirtmunlikaporn S, Brady B, Vasquez EI. Nurses' perception of their work environment, health, and well-being: a qualitative perspective. *AAOHN J*. 2004;52(1):16-22.

Genevay S, Cedraschi C, Courvoisier DS, Perneger TV, Grandjean R, Griesser AC, et al. Work related characteristics of back and neck pain among employees of a Swiss University Hospital. *Joint Bone Spine*. 2011;78(4):392-7.

Gershon RR, Lin S, Li X. Work stress in aging police officers. *J Occup Environ Med*. 2002;44(2):160-7.

Griefahn B, Mehnert P, Brode P, Forsthoff A. Working in moderate cold: A possible risk to health. *Journal of Occupational Health*. 1997;39(1):36-44.

Hagg O, Fritzell P, Nordwall A. Characteristics of patients with chronic low back pain selected for surgery: A comparison with the general population reported from the Swedish Lumbar Spine Study. *Spine (Phila Pa 1976)*. 2002;27(11):1223-30.

Inaba R, Mirbod SM, Kurokawa J, Inoue M, Iwata H. Subjective symptoms among female workers and winter working conditions in a consumer cooperative. *Journal of Occupational Health*. 2005;47(5):454-65.

Johansson AC, Cornefjord M, Bergkvist L, Ohrvik J, Linton SJ. Psychosocial stress factors among patients with lumbar disc herniation, scheduled for disc surgery in comparison with patients scheduled for arthroscopic knee surgery. *Eur Spine J*. 2007;16(7):961-70.

Kierklo A, Kobus A, Jaworska M, Botulinski B. Work-related musculoskeletal disorders among dentists - a questionnaire survey. *Ann Agric Environ Med*. 2011;18(1):79-84.

Kjellberg K, Lagerstrom M, Hagberg M. Work technique of nurses in patient transfer tasks and associations with personal factors. *Scand J Work Environ Health*. 2003;29(6):468-77.

Knibbe JJ, Friele RD. Prevalence of back pain and characteristics of the physical workload of community nurses. *Ergonomics*. 1996;39(2):186-98.

- Kolstrup C, Lundqvist P, Pinzke S. Psychosocial work environment among employed Swedish dairy and pig farmworkers. *Journal of agromedicine*. 2008;13(1):23-36.
- Kostova V, Koleva M. Back disorders (low back pain, cervicobrachial and lumbosacral radicular syndromes) and some related risk factors. *J Neurol Sci*. 2001;192(1-2):17-25.
- Krajcarski S, Wells R. The time variation pattern of mechanical exposure and the reporting of low back pain. *Theoretical Issues in Ergonomics Science*. 2008;9(1):45-71.
- Krause N, Ragland DR, Greiner BA, Fisher JM, Holman BL, Selvin S. Physical workload and ergonomic factors associated with prevalence of back and neck pain in urban transit operators. *Spine (Phila Pa 1976)*. 1997;22(18):2117-26; discussion 27.
- Krause N, Ragland DR, Greiner BA, Syme SL, Fisher JM. Psychosocial job factors associated with back and neck pain in public transit operators. *Scand J Work Environ Health*. 1997;23(3):179-86.
- Kristensen TS. The occupational environment and health of slaughterhouse workers. V. Low back pain and absence on account of low back symptoms. *Ugeskrift for Laeger*. 1985;147(41):3276-83.
- Kromark K, Dulon M, Beck BB, Nienhaus A. Back disorders and lumbar load in nursing staff in geriatric care: a comparison of home-based care and nursing homes. *J Occup Med Toxicol*. 2009;4:33.
- Larsen PS, Strandberg-Larsen K, Juhl M, Svendsen SW, Bonde JP, Andersen AM. Occupational lifting and pelvic pain during pregnancy: a study within the Danish National Birth Cohort. *Scand J Work Environ Health*. 2013;39(1):88-95.
- Lavender SA, Marras WS, Ferguson SA, Splittstoesser RE, Yang G. Developing physical exposure-based back injury risk models applicable to manual handling jobs in distribution centers. *9(7):450-9*.
- Leijon O, Harenstam A, Waldenstrom K, Alderling M, Vingard E. Target groups for prevention of neck/shoulder and low back disorders: an exploratory cluster analysis of working and living conditions. *Work*. 2006;27(2):189-204.
- Linton SJ, Buer N. Working despite pain: factors associated with work attendance versus dysfunction. *International journal of behavioral medicine*. 1995;2(3):252-62.

Linton SJ, Hellsing A-L, Bryngelsson I-L. Psychosocial and functional risk factors in the early stages of back pain: Effects on moderate pain. *International journal of behavioral medicine*. 2000;7(4):291-304.

Linton SJ, Warg LE. Attributions (beliefs) and job satisfaction associated with back pain in an industrial setting. *Percept Mot Skills*. 1993;76(1):51-62.

Lundberg G, Gerdle B. Correlations between joint and spinal mobility, spinal sagittal configuration, segmental mobility, segmental pain, symptoms and disabilities in female homecare personnel. *Scand J Rehabil Med*. 2000;32(3):124-33.

Marras WS, Lavender SA, Leurgans SE, Fathallah FA, Ferguson SA, Allread WG, et al. Biomechanical risk factors for occupationally related low back disorders. *Ergonomics*. 1995;38(2):377-410.

Marras WS, Lavender SA, Leurgans SE, Rajulu SL, Allread WG, Fathallah FA, et al. The role of dynamic three-dimensional trunk motion in occupationally-related low back disorders. The effects of workplace factors, trunk position, and trunk motion characteristics on risk of injury. *Spine (Phila Pa 1976)*. 1993;18(5):617-28.

Milosavljevic S, Gregory DE, Pal P, Carman AB, Milburn PD, Callaghan JP. The interaction between skill, postures, forces and back pain in wool handling. *Appl Ergon*. 2011;42(6):801-6.

Mirbod SM, Inaba R, Iwata H. Low back pain among different groups of subjects exposed to hand-arm transmitted vibration. *Ind Health*. 1997;35(2):212-21.

Moens GF, Dohogne T, Jacques P, Van Helshoecht P. Back pain and its correlates among workers in family care. *Occupational Medicine*. 1993;43(2):78-84.

Mustard CA, Vermeulen M, Lavis JN. Is position in the occupational hierarchy a determinant of decline in perceived health status? *Soc Sci Med*. 2003;57(12):2291-303.

Netterstrom B, Juel K. Low back trouble among urban bus drivers in Denmark. *Scandinavian journal of social medicine*. 1989;17(2):203-6.

Nielsen K, Albertsen K, Brenner SO, Smith-Hansen L, Roepsdorff C. Comparing working conditions and physical and psychological health complaints in four occupational groups working in female-dominated workplaces. *Int Arch Occup Environ Health*. 2009;82(10):1229-39.

Nilsson A, Sjoden PO, Dahl J, Denison E. Factors related to long-duration pain and sick leave among Swedish staff working in the public health service. *Scand J Caring Sci.* 2005;19(4):419-26.

Orsello CA, Phillips AS, Rice GM. Height and in-flight low back pain association among military helicopter pilots. *Aviat Space Environ Med.* 2013;84(1):32-7.

Riihimaki H. Back pain and heavy physical work: a comparative study of concrete reinforcement workers and maintenance house painters. *British journal of industrial medicine.* 1985;42(4):226-32.

Tsai SP, Gilstrap EL, Cowles SR, Waddell LC, Jr., Ross CE. Personal and job characteristics of musculoskeletal injuries in an industrial population. *J Occup Med.* 1992;34(6):606-12.

Westgaard RH, Jansen T. Individual and work related factors associated with symptoms of musculoskeletal complaints. II. Different risk factors among sewing machine operators. *British journal of industrial medicine.* 1992;49(3):154-62.

Låg kvalitet. Studier av tvärsnittsdesign

Aasa U, Barnekow-Bergkvist M, Angquist KA, Brulin C. Relationships between work-related factors and disorders in the neck-shoulder and low-back region among female and male ambulance personnel. *J Occup Health*. 2005;47(6):481-9.

Abbas J, Hamoud K, May H, Peled N, Sarig R, Stein D, et al. Socioeconomic and physical characteristics of individuals with degenerative lumbar spinal stenosis. *Spine (Phila Pa 1976)*. 2013;38(9):E554-E61.

Ahlberg-Hulten GK, Theorell T, Sigala F. Social support, job strain and musculoskeletal pain among female health care personnel. *Scand J Work Environ Health*. 1995;21(6):435-9.

Ahlgren C, Malmgren Olsson EB, Brulin C. Gender analysis of musculoskeletal disorders and emotional exhaustion: interactive effects from physical and psychosocial work exposures and engagement in domestic work. *Ergonomics*. 2012;55(2):212-28.

Alcouffe J, Manillier P, Brehier M, Fabin C, Faupin F. Analysis by sex of low back pain among workers from small companies in the Paris area: severity and occupational consequences. *Occup Environ Med*. 1999;56(10):696-701.

Alexopoulos EC, Burdorf A, Kalokerinou A. Risk factors for musculoskeletal disorders among nursing personnel in Greek hospitals. *Int Arch Occup Environ Health*. 2003;76(4):289-94.

Alexopoulos EC, Stathi IC, Charizani F. Prevalence of musculoskeletal disorders in dentists. *BMC Musculoskelet Disord*. 2004;5:16.

Alexopoulos EC, Tanagra D, Konstantinou E, Burdorf A. Musculoskeletal disorders in shipyard industry: prevalence, health care use, and absenteeism. *BMC Musculoskelet Disord*. 2006;7:88.

Ando S, Ono Y, Shimaoka M, Hiruta S, Hattori Y, Hori F, et al. Associations of self estimated workloads with musculoskeletal symptoms among hospital nurses. *Occup Environ Med*. 2000;57(3):211-6.

Arlinghaus A, Caban-Martinez AJ, Marino M, Reme SE. The role of ergonomic and psychosocial workplace factors in the reporting of back injuries among U.S. home health aides. *Am J Ind Med*. 2013.

- Arlinghaus A, Caban-Martinez AJ, Marino M, Reme SE. The role of ergonomic and psychosocial workplace factors in the reporting of back injuries among U.S. home health aides. *Am J Ind Med.* 2013;56(10):1239-44.
- Barnekow-Bergkvist M, Hedberg GE, Janlert U, Jansson E. Determinants of self-reported neck-shoulder and low back symptoms in a general population. *Spine (Phila Pa 1976).* 1998;23(2):235-43.
- Bergenudd H, Nilsson B. The prevalence of locomotor complaints in middle age and their relationship to health and socioeconomic factors. *Clinical orthopaedics and related research.* 1994(308):264-70.
- Bernard C, Courouve L, Bouee S, Adjemian A, Chretien JC, Niedhammer I. Biomechanical and Psychosocial Work Exposures and Musculoskeletal Symptoms among Vineyard Workers. *J Occup Health.* 2011.
- Birnie D, Healey JS, Krahn AD, Ahmad K, Crystal E, Khaykin Y, et al. Prevalence and risk factors for cervical and lumbar spondylosis in interventional electrophysiologists. *Journal of cardiovascular electrophysiology.* 2011;22(9):957-60.
- Bos E, Krol B, van der Star L, Groothoff J. Risk factors and musculoskeletal complaints in non-specialized nurses, IC nurses, operation room nurses, and X-ray technologists. *Int Arch Occup Environ Health.* 2007;80(3):198-206.
- Bovenzi M, Zadini A. Self-reported low back symptoms in urban bus drivers exposed to whole-body vibration. *Spine (Phila Pa 1976).* 1992;17(9):1048-59.
- Bridger RS, Groom MR, Jones H, Pethybridge RJ, Pullinger N. Task and postural factors are related to back pain in helicopter pilots. *Aviat Space Environ Med.* 2002;73(8):805-11.
- Broniecki M, Esterman A, Grantham H. Risk factors for back, neck and shoulder musculoskeletal injuries and claims in ambulance officers. *Journal of Musculoskeletal Research.* 2012;15(1).
- Brulin C, Gerdle B, Granlund B, Hoog J, Knutson A, Sundelin G. Physical and psychosocial work-related risk factors associated with musculoskeletal symptoms among home care personnel. *Scand J Caring Sci.* 1998;12(2):104-10.
- Buckle PW, Kember PA, Wood AD, Wood SN. Factors influencing occupational back pain in Bedfordshire. *Spine (Phila Pa 1976).* 1980;5(3):254-8.

Burton AK, Tillotson KM, Symonds TL, Burke C, Mathewson T. Occupational risk factors for the first-onset and subsequent course of low back trouble. A study of serving police officers. *Spine (Phila Pa 1976)*. 1996;21(22):2612-20.

Byrns G, Reeder G, Jin G, Pachis K. Risk factors for work-related low back pain in registered nurses, and potential obstacles in using mechanical lifting devices. *J Occup Environ Hyg*. 2004;1(1):11-21.

Camerino D, Cesana GC, Molteni G, De Vito G, Evaristi C, Latocca R. Job strain and musculoskeletal disorders of Italian nurses. *Occupational Ergonomics*. 2001;2(4):215-23.

Cameron L, Lalich N, Bauer S, Booker V, Bogue HO, Samuels S, et al. Occupational health survey of farm workers by camp health aides. *J Agric Saf Health*. 2006;12(2):139-53.

Canjuga M, Läubli T, Bauer GF. Can the job demand control model explain back and neck pain? Cross-sectional study in a representative sample of Swiss working population. *Int J Ind Ergon*. 2010;40(6):663-8.

Carugno M, Pesatori AC, Ferrario MM, Ferrari AL, Silva FJ, Martins AC, et al. Physical and psychosocial risk factors for musculoskeletal disorders in Brazilian and Italian nurses. *Cadernos de saude publica / Ministerio da Saude, Fundacao Oswaldo Cruz, Escola Nacional de Saude Publica*. 2012;28(9):1632-42.

Coggon D, Ntani G, Palmer KT, Felli VE, Harari R, Barrero LH, et al. Disabling musculoskeletal pain in working populations: Is it the job, the person, or the culture? *Pain*. 2013;154(6):856-63.

Cole DC, Ibrahim SA, Shannon HS, Scott F, Eyles J. Work correlates of back problems and activity restriction due to musculoskeletal disorders in the Canadian national population health survey (NPHS) 1994-5 data. *Occup Environ Med*. 2001;58(11):728-34.

Courvoisier DS, Genevay S, Cedraschi C, Bessire N, Griesser-Delacretaz AC, Monnin D, et al. Job strain, work characteristics and back pain: a study in a university hospital. *Eur J Pain*. 2011;15(6):634-40.

Cunningham C, Flynn T, Blake C. Low back pain and occupation among Irish health service workers. *Occup Med (Lond)*. 2006;56(7):447-54.

- Damkot DK, Pope MH, Lord J, Frymoyer JW. The relationship between work history, work environment and low-back pain in men. *Spine (Phila Pa 1976)*. 1984;9(4):395-9.
- Daraiseh N, Genaidy AM, Karwowski W, Davis LS, Stambough J, Huston RI. Musculoskeletal outcomes in multiple body regions and work effects among nurses: the effects of stressful and stimulating working conditions. *Ergonomics*. 2003;46(12):1178-99.
- de Zwart BC, Broersen JP, Frings-Dresen MH, van Dijk FJ. Musculoskeletal complaints in The Netherlands in relation to age, gender and physically demanding work. *Int Arch Occup Environ Health*. 1997;70(5):352-60.
- Devereux JJ, Buckle PW, Vlachonikolis IG. Interactions between physical and psychosocial risk factors at work increase the risk of back disorders: an epidemiological approach. *Occup Environ Med*. 1999;56(5):343-53.
- Eatough EM, Way JD, Chang CH. Understanding the link between psychosocial work stressors and work-related musculoskeletal complaints. *Appl Ergon*. 2012;43(3):554-63.
- Elders LA, Burdorf A. Interrelations of risk factors and low back pain in scaffolders. *Occup Environ Med*. 2001;58(9):597-603.
- Engels JA, van der Gulden JW, Senden TF, van't Hof B. Work related risk factors for musculoskeletal complaints in the nursing profession: results of a questionnaire survey. *Occup Environ Med*. 1996;53(9):636-41.
- Engholm G, Holmstrom E. Dose-response associations between musculoskeletal disorders and physical and psychosocial factors among construction workers. *Scand J Work Environ Health*. 2005;31 Suppl 2:57-67.
- Engstrom T, Hanse JJ, Kadefors R. Musculoskeletal symptoms due to technical preconditions in long cycle time work in an automobile assembly plant: a study of prevalence and relation to psychosocial factors and physical exposure. *Appl Ergon*. 1999;30(5):443-53.
- Eriksen HR, Ihlebaek C, Jansen JP, Burdorf A. The relations between psychosocial factors at work and health status among workers in home care organizations. *International journal of behavioral medicine*. 2006;13(3):183-92.

Estryn-Behar M, Kaminski M, Peigne E, Maillard MF, Pelletier A, Berthier C, et al. Strenuous working conditions and musculo-skeletal disorders among female hospital workers. *Int Arch Occup Environ Health*. 1990;62(1):47-57.

Evans W, Jobe W, Seibert C. A cross-sectional prevalence study of lumbar disc degeneration in a working population. *Spine (Phila Pa 1976)*. 1989;14(1):60-4.

Farioli A, Mattioli S, Quagliari A, Curti S, Violante FS, Coggon D. Musculoskeletal pain in Europe: the role of personal, occupational, and social risk factors. *Scand J Work Environ Health*. 2014;40(1):36-46.

Feyer AM, Williamson A, Mandryk J, de Silva I, Healy S. Role of psychosocial risk factors in work-related low-back pain. *Scand J Work Environ Health*. 1992;18(6):368-75.

Firth H, Herbison P, McBride D, Feyer AM. Low back pain among farmers in Southland, NZ. *Journal of Occupational Health and Safety - Australia and New Zealand*. 2002;18(2):167-71.

Foppa I, Noack RH. The relation of self-reported back pain to psychosocial, behavioral, and health-related factors in a working population in Switzerland. *Soc Sci Med*. 1996;43(7):1119-26.

Frymoyer JW, Pope MH, Costanza MC. Epidemiologic studies of low-back pain. *Spine*. 1980;5(5):419-23.

Gamperiene M, Stigum H. Work related risk factors for musculoskeletal complaints in the spinning industry in Lithuania. *Occup Environ Med*. 1999;56(6):411-6.

Gheldof EL, Vinck J, Van den Bussche E, Vlaeyen JW, Hidding A, Crombez G. Pain and pain-related fear are associated with functional and social disability in an occupational setting: evidence of mediation by pain-related fear. *Eur J Pain*. 2006;10(6):513-25.

Gheldof EL, Vinck J, Vlaeyen JW, Hidding A, Crombez G. The differential role of pain, work characteristics and pain-related fear in explaining back pain and sick leave in occupational settings. *Pain*. 2005;113(1-2):71-81.

Goldsheyder D, Nordin M, Weiner SS, Hiebert R. Musculoskeletal symptom survey among mason tenders. *Am J Ind Med*. 2002;42(5):384-96.

- Greiner BA, Krause N. Observational stress factors and musculoskeletal disorders in urban transit operators. *Journal of occupational health psychology*. 2006;11(1):38-51.
- Grooten WJ, Wernstedt P, Campo M. Work-related musculoskeletal disorders in female Swedish physical therapists with more than 15 years of job experience: prevalence and associations with work exposures. *Physiother Theory Pract*. 2011;27(3):213-22.
- Grzywacz JG, Arcury TA, Mora D, Anderson AM, Chen H, Rosenbaum DA, et al. Work organization and musculoskeletal health: clinical findings from immigrant Latino poultry processing and other manual workers. *J Occup Environ Med*. 2012;54(8):995-1001.
- Guo H. Working hours spent on repeated activities and prevalence of back pain. *Occupational & Environmental Medicine*. 2002;59(10):680-8.
- Hagen KB, Magnus P, Vetlesen K. Neck/shoulder and low-back disorders in the forestry industry: relationship to work tasks and perceived psychosocial job stress. *Ergonomics*. 1998;41(10):1510-8.
- Hammig O, Gutzwiller F, Bauer G. Work-life conflict and associations with work- and nonwork-related factors and with physical and mental health outcomes: a nationally representative cross-sectional study in Switzerland. *BMC public health*. 2009;9:435.
- Hangai M, Kaneoka K, Kuno S, Hinotsu S, Sakane M, Mamizuka N, et al. Factors associated with lumbar intervertebral disc degeneration in the elderly. *Spine J*. 2008;8(5):732-40.
- Harber P, Billet E, Lew M, Horan M. Importance of non-patient transfer activities in nursing-related back pain: I. Questionnaire survey. *J Occup Med*. 1987;29(12):967-70.
- Harcombe H, McBride D, Derrett S, Gray A. Physical and psychosocial risk factors for musculoskeletal disorders in New Zealand nurses, postal workers and office workers. *Inj Prev*. 2010;16(2):96-100.
- Heliovaara M, Makela M, Knekt P, Impivaara O, Aromaa A. Determinants of sciatica and low-back pain. *Spine (Phila Pa 1976)*. 1991;16(6):608-14.
- Heneweer H, Picavet HSJ, Staes F, Kiers H, Vanhees L. Physical fitness, rather than self-reported physical activities, is more strongly associated with low back pain: Evidence from a working population. *European Spine Journal*. 2012;21(7):1265-72.

- Hermes ED, Webb TS, Wells TS. Aircraft type and other risk factors for spinal disorders: data from 19,673 military cockpit aircrew. *Aviat Space Environ Med.* 2010;81(9):850-6.
- Hollingdale R, Warin J. Back pain in nursing and associated factors: a study. *Nurs Stand.* 1997;11(39):35-8.
- Holmberg S, Thelin A, Stiernstrom E, Svardsudd K. The impact of physical work exposure on musculoskeletal symptoms among farmers and rural non-farmers. *Ann Agric Environ Med.* 2003;10(2):179-84.
- Inaba R, Mirbod SM. Subjective musculoskeletal symptoms in winter and summer among indoor working construction electricians. *Ind Health.* 2010;48(1):29-37.
- Inaba R, Okumura M, Mirbod SM. Subjective symptoms of female workers sorting goods in summer. *Ind Health.* 2011;49(4):464-74.
- Jablonska B, Soares JJF, Sundin O. Pain among women: Associations with socio-economic and work conditions. *European Journal of Pain.* 2006;10(5):435-47.
- Jacobsen S, Sonne-Holm S, Roving H, Monrad H, Gebuhr P. Degenerative lumbar spondylolisthesis: an epidemiological perspective: the Copenhagen Osteoarthritis Study. *Spine (Phila Pa 1976).* 2007;32(1):120-5.
- Jacobsson L, Lindgarde F, Manthorpe R, Ohlsson K. Effect of education, occupation and some lifestyle factors on common rheumatic complaints in a Swedish group aged 50-70 years. *Ann Rheum Dis.* 1992;51(7):835-43.
- Johansson JA. Psychosocial work factors, physical work load and associated musculoskeletal symptoms among home care workers. *Scandinavian journal of psychology.* 1995;36(2):113-29.
- Johansson JA, Rubenowitz S. Risk indicators in the psychosocial and physical work environment for work-related neck, shoulder and low back symptoms: a study among blue- and white-collar workers in eight companies. *Scand J Rehabil Med.* 1994;26(3):131-42.
- Jzelenberg WI, Burdorf A. Risk factors for musculoskeletal symptoms and ensuing health care use and sick leave. *Spine (Phila Pa 1976).* 2005;30(13):1550-6.

Jzelenberg WI, Molenaar D, Burdorf A. Different risk factors for musculoskeletal complaints and musculoskeletal sickness absence. *Scand J Work Environ Health*. 2004;30(1):56-63.

Karahan A, Kav S, Abbasoglu A, Dogan N. Low back pain: prevalence and associated risk factors among hospital staff. *J Adv Nurs*. 2009;65(3):516-24.

Karjalainen U, Paananen M, Okuloff A, Taimela S, Auvinen J, Mannikko M, et al. Role of environmental factors and history of low back pain in sciatica symptoms among finnish adolescents. *Spine (Phila Pa 1976)*. 2013;24.

Kim SS, Okechukwu CA, Dennerlein JT, Boden LI, Hopcia K, Hashimoto DM, et al. Association between perceived inadequate staffing and musculoskeletal pain among hospital patient care workers. *Int Arch Occup Environ Health*. 2013.

Krause N, Scherzer T, Rugulies R. Physical workload, work intensification, and prevalence of pain in low wage workers: results from a participatory research project with hotel room cleaners in Las Vegas. *Am J Ind Med*. 2005;48(5):326-37.

Kumar S. Cumulative load as a risk factor for back pain. *Spine (Phila Pa 1976)*. 1990;15(12):1311-6.

Lagerstrom M, Wenemark M, Hagberg M, Hjelm EW. Occupational and individual factors related to musculoskeletal symptoms in five body regions among Swedish nursing personnel. *Int Arch Occup Environ Health*. 1995;68(1):27-35.

Larsson LG, Mudholkar GS, Baum J, Srivastava DK. Benefits and liabilities of hypermobility in the back pain disorders of industrial workers. *J Intern Med*. 1995;238(5):461-7.

Latza U, Kohlmann T, Deck R, Raspe H. Influence of occupational factors on the relation between socioeconomic status and self-reported back pain in a population-based sample of German adults with back pain. *Spine (Phila Pa 1976)*. 2000;25(11):1390-7.

Leboeuf-Yde C, Kjaer P, Bendix T, Manniche C. Self-reported hard physical work combined with heavy smoking or overweight may result in so-called Modic changes. *BMC Musculoskelet Disord*. 2008;9:5.

Lee H, Wilbur J, Kim MJ, Miller AM. Psychosocial risk factors for work-related musculoskeletal disorders of the lower-back among long-haul international female flight attendants. *J Adv Nurs*. 2008;61(5):492-502.

Lee P, Helewa A, Goldsmith CH, Smythe HA, Stitt LW. Low back pain: prevalence and risk factors in an industrial setting. *J Rheumatol*. 2001;28(2):346-51.

Lefevre-Colau MM, Fayad F, Rannou F, Fermanian J, Coriat F, Mace Y, et al. Frequency and interrelations of risk factors for chronic low back pain in a primary care setting. *PLoS One*. 2009;4(3):e4874.

Lehto TU, Helenius HY, Alaranta HT. Musculoskeletal symptoms of dentists assessed by a multidisciplinary approach. *Community dentistry and oral epidemiology*. 1991;19(1):38-44.

Leino-Arjas P, Kaila-Kangas L, Kauppinen T, Notkola V, Keskimaki I, Mutanen P. Occupational exposures and inpatient hospital care for lumbar intervertebral disc disorders among Finns. *Am J Ind Med*. 2004;46(5):513-20.

Lemasters GK, Atterbury MR, Booth-Jones AD, Bhattacharya A, Ollila-Glenn N, Forrester C, et al. Prevalence of work related musculoskeletal disorders in active union carpenters. *Occup Environ Med*. 1998;55(6):421-7.

Leroux I, Dionne CE, Bourbonnais R, Brisson C. Prevalence of musculoskeletal pain and associated factors in the Quebec working population. *Int Arch Occup Environ Health*. 2005;78(5):379-86.

Levangie PK. Association of low back pain with self-reported risk factors among patients seeking physical therapy services. *Phys Ther*. 1999;79(8):757-66.

Liira JP, Shannon HS, Chambers LW, Haines TA. Long-term back problems and physical work exposures in the 1990 Ontario Health Survey. *Am J Public Health*. 1996;86(3):382-7.

Linton SJ. Risk factors for neck and back pain in a working population in Sweden. *Work & Stress*. 1990;4(1):41-9.

Lipscomb J, Trinkoff A, Brady B, Geiger-Brown J. Health care system changes and reported musculoskeletal disorders among registered nurses. *Am J Public Health*. 2004;94(8):1431-5.

- Lipscomb JA, Trinkoff AM, Geiger-Brown J, Brady B. Work-schedule characteristics and reported musculoskeletal disorders of registered nurses. *Scand J Work Environ Health*. 2002;28(6):394-401.
- Lorusso A, Bruno S, L'Abbate N. Musculoskeletal complaints among Italian X-ray technologists. *Ind Health*. 2007;45(5):705-8.
- Magnusson ML, Pope MH, Wilder DG, Areskoug B. Are occupational drivers at an increased risk for developing musculoskeletal disorders? *Spine (Phila Pa 1976)*. 1996;21(6):710-7.
- Mandel JH, Lohman W. Low back pain in nurses: the relative importance of medical history, work factors, exercise, and demographics. *Res Nurs Health*. 1987;10(3):165-70.
- Mariconda M, Galasso O, Imbimbo L, Lotti G, Milano C. Relationship between alterations of the lumbar spine, visualized with magnetic resonance imaging, and occupational variables. *Eur Spine J*. 2007;16(2):255-66.
- Masset D, Malchaire J. Low back pain. Epidemiologic aspects and work-related factors in the steel industry. *Spine (Phila Pa 1976)*. 1994;19(2):143-6.
- Matsudaira K, Palmer KT, Reading I, Hirai M, Yoshimura N, Coggon D. Prevalence and correlates of regional pain and associated disability in Japanese workers. *Occup Environ Med*. 2011;68(3):191-6.
- Matsui H, Maeda A, Tsuji H, Naruse Y. Risk indicators of low back pain among workers in Japan. Association of familial and physical factors with low back pain. *Spine (Phila Pa 1976)*. 1997;22(11):1242-7; discussion 8.
- McGill S, Grenier S, Bluhm M, Preuss R, Brown S, Russell C. Previous history of LBP with work loss is related to lingering deficits in biomechanical, physiological, personal, psychosocial and motor control characteristics. *Ergonomics*. 2003;46(7):731-46.
- Meeker W, Menke M, Waldorf T, Mootz R. Factors related to acute occupational low back injuries. *JNMS: Journal of the Neuromusculoskeletal System*. 1997;5(2):59-65.
- Mehlum IS, Kristensen P, Kjuus H, Wergeland E. Are occupational factors important determinants of socioeconomic inequalities in musculoskeletal pain? *Scand J Work Environ Health*. 2008;34(4):250-9.
- Meyer JP, Flenghi D, Deschamps JP. Effects of Manual Handling, Posture, and Whole Body Vibrations on Low-Back Pain. *Int J Occup Saf Ergon*. 1998;4(4):449-70.

Milutinovic D, Golubovic B, Brkic N, Prokes B. Professional stress and health among critical care nurses in Serbia. *Arhiv za higijenu rada i toksikologiju*. 2012;63(2):171-80.

Miranda H, Punnett L, Gore R, Boyer J. Violence at the workplace increases the risk of musculoskeletal pain among nursing home workers. *Occup Environ Med*. 2011;68(1):52-7.

Miranda H, Viikari-Juntura E, Martikainen R, Takala EP, Riihimaki H. Physical exercise and musculoskeletal pain among forest industry workers. *Scand J Med Sci Sports*. 2001;11(4):239-46.

Miyamoto M, Konno S, Gembun Y, Liu X, Minami K, Ito H. Epidemiological study of low back pain and occupational risk factors among taxi drivers. *Ind Health*. 2008;46(2):112-7.

Miyamoto M, Shirai Y, Nakayama Y, Gembun Y, Kaneda K. An epidemiologic study of occupational low back pain in truck drivers. *Journal of Nippon Medical School = Nippon Ika Daigaku zasshi*. 2000;67(3):186-90.

Morken T, Mageroy N, Moen BE. Physical activity is associated with a low prevalence of musculoskeletal disorders in the Royal Norwegian Navy: a cross sectional study. *BMC Musculoskelet Disord*. 2007;8:56.

Moscato U, Trinca D, Rega ML, Mannocci A, Chiaradia G, Grieco G, et al. Musculoskeletal injuries among operating room nurses: Results from a multicenter survey in Rome, Italy. *Journal of Public Health*. 2010;18(5):453-9.

Muraki S, Akune T, Oka H, Mabuchi A, En-Yo Y, Yoshida M, et al. Association of occupational activity with radiographic knee osteoarthritis and lumbar spondylosis in elderly patients of population-based cohorts: a large-scale population-based study. *Arthritis Rheum*. 2009;61(6):779-86.

Murtezani A, Ibraimi Z, Sllamniku S, Osmani T, Sherifi S. Prevalence and risk factors for low back pain in industrial workers. *Folia medica*. 2011;53(3):68-74.

Musson Y, Burdorf A, van Drimmelen D. Exposure to shock and vibration and symptoms in workers using impact power tools. *Ann Occup Hyg*. 1989;33(1):85-96.

Nag A, Vyas H, Shah P, Nag PK. Risk factors and musculoskeletal disorders among women workers performing fish processing. *Am J Ind Med*. 2012;55(9):833-43.

- Nagasu M, Sakai K, Ito A, Tomita S, Temmyo Y, Ueno M, et al. Prevalence and risk factors for low back pain among professional cooks working in school lunch services. *BMC public health*. 2007;7:171.
- Nahit ES, Macfarlane GJ, Pritchard CM, Cherry NM, Silman AJ. Short term influence of mechanical factors on regional musculoskeletal pain: a study of new workers from 12 occupational groups. *Occup Environ Med*. 2001;58(6):374-81.
- Nahit ES, Pritchard CM, Cherry NM, Silman AJ, MacFarlane GJ. The influence of work related psychosocial factors and psychological distress on regional musculoskeletal pain: a study of newly employed workers. *Journal of Rheumatology*. 2001;28(6):1378-84.
- Nayha S, Anttonen H, Hassi J. Snowmobile driving and symptoms of the locomotive organs. *Arctic medical research*. 1994;53 Suppl 3:41-4.
- Oksuz E. Prevalence, risk factors, and preference-based health states of low back pain in a Turkish population. *Spine (Phila Pa 1976)*. 2006;31(25):E968-E72.
- Okunribido OO, Shimbles SJ, Magnusson M, Pope M. City bus driving and low back pain: a study of the exposures to posture demands, manual materials handling and whole-body vibration. *Appl Ergon*. 2007;38(1):29-38.
- Ono Y, Shimaoka M, Hiruta S, Takeuchi Y. Low back pain among cooks in nursery schools. *Ind Health*. 1997;35(2):194-201.
- Osborne A, Finnegan G, Blake C, Meredith D, McNamara J, Phelan J, et al. An evaluation of low back pain among farmers in Ireland. *Occup Med (Lond)*. 2013;63(1):53-9.
- Ostgaard HC, Andersson GB, Karlsson K. Prevalence of back pain in pregnancy. *Spine (Phila Pa 1976)*. 1991;16(5):549-52.
- Ozguler A, Leclerc A, Landre MF, Pietri-Taleb F, Niedhammer I. Individual and occupational determinants of low back pain according to various definitions of low back pain. *J Epidemiol Community Health*. 2000;54(3):215-20.
- Palmer KT, Griffin MJ, Syddall HE, Pannett B, Cooper C, Coggon D. The relative importance of whole body vibration and occupational lifting as risk factors for low-back pain. *Occupational & Environmental Medicine*. 2003;60(10):715-21.

Picavet HS, Schouten JS. Physical load in daily life and low back problems in the general population-The MORGEN study. *Prev Med.* 2000;31(5):506-12.

Raanaas RK, Anderson D. A questionnaire survey of Norwegian taxi drivers' musculoskeletal health, and work-related risk factors. *Int J Ind Ergon.* 2008;38(3-4):280-90.

Riihimaki H, Tola S, Videman T, Hanninen K. Low-back pain and occupation. A cross-sectional questionnaire study of men in machine operating, dynamic physical work, and sedentary work. *Spine (Phila Pa 1976).* 1989;14(2):204-9.

Robstad Andersen G, Westgaard RH. Perceived occupational exposures of home care workers and the association to general tension, shoulder-neck and low back pain. *Work.* 2013.

Roelen CA, Schreuder KJ, Koopmans PC, Groothoff JW. Perceived job demands relate to self-reported health complaints. *Occup Med (Lond).* 2008;58(1):58-63.

Rolander B, Bellner A. Experience of musculo-skeletal disorders, intensity of pain, and general conditions in work -- the case of employees in non-private dental clinics in a county in southern Sweden. *Work.* 2001;17(1):65-73.

Roy TC, Ritland BM, Knapik JJ, Sharp MA. Lifting tasks are associated with injuries during the early portion of a deployment to Afghanistan. *Mil Med.* 2012;177(6):716-22.

Salyga J, Kusleikaite M. Factors influencing psychoemotional strain and fatigue, and relationship of these factors with health complaints at sea among Lithuanian seafarers. *Medicina (Kaunas, Lithuania).* 2011;47(12):675-81.

Sanders MJ, Turcotte CM. Occupational stress in dental hygienists. *Work: Journal of Prevention, Assessment & Rehabilitation.* 2010;35(4):455-65.

Saraste H, Hultman G. Life conditions of persons with and without low-back pain. *Scand J Rehabil Med.* 1987;19(3):109-13.

Schneider S, Lipinski S, Schiltewolf M. Occupations associated with a high risk of self-reported back pain: representative outcomes of a back pain prevalence study in the Federal Republic of Germany. *Eur Spine J.* 2006;15(6):821-33.

Schneider S, Schmitt H, Zoller S, Schiltewolf M. Workplace stress, lifestyle and social factors as correlates of back pain: a representative study of the German working population. *Int Arch Occup Environ Health.* 2005;78(4):253-69.

- Schreuder KJ, Roelen CA, Koopmans PC, Groothoff JW. Job demands and health complaints in white and blue collar workers. *Work*. 2008;31(4):425-32.
- Secer M, Nacar OA, Muradov MJ, Altintoprak F, Kabali B, Senol Z, et al. Nonspecific low back pain in a group of young adult men. *Turk Neurosurg*. 2011;21(2):135-9.
- Sjolander P, Daerga L, Edin-Liljegren A, Jacobsson L. Musculoskeletal symptoms and perceived work strain among reindeer herders in Sweden. *Occup Med (Lond)*. 2008;58(8):572-9.
- Skov T, Borg V, Orhede E. Psychosocial and physical risk factors for musculoskeletal disorders of the neck, shoulders, and lower back in salespeople. *Occup Environ Med*. 1996;53(5):351-6.
- Skovron ML, Szpalski M, Nordin M, Melot C, Cukier D. Sociocultural factors and back pain. A population-based study in Belgian adults. *Spine (Phila Pa 1976)*. 1994;19(2):129-37.
- Smedley J, Egger P, Cooper C, Coggon D. Manual handling activities and risk of low back pain in nurses. *Occup Environ Med*. 1995;52(3):160-3.
- Smith DR, Kondo N, Tanaka E, Tanaka H, Hirasawa K, Yamagata Z. Musculoskeletal disorders among hospital nurses in rural Japan. *Rural and remote health*. 2003;3(3):241.
- Smith DR, Leggat PA, Speare R. Musculoskeletal disorders and psychosocial risk factors among veterinarians in Queensland, Australia. *Australian veterinary journal*. 2009;87(7):260-5.
- Smith DR, Mihashi M, Adachi Y, Koga H, Ishitake T. A detailed analysis of musculoskeletal disorder risk factors among Japanese nurses. *Journal of Safety Research*. 2006;37(2):195-200.
- Sobti A, Cooper C, Inskip H, Searle S, Coggon D. Occupational physical activity and long-term risk of musculoskeletal symptoms: a national survey of post office pensioners. *Am J Ind Med*. 1997;32(1):76-83.
- Solanki M, Carr D, Martin M. Back pain among echocardiographers. *Heart (British Cardiac Society)*. 1997;78 Suppl 1:23-8.
- Sorensen G, Stoddard AM, Stoffel S, Buxton O, Sembajwe G, Hashimoto D, et al. The role of the work context in multiple wellness outcomes for hospital patient care workers. *J Occup Environ Med*. 2011;53(8):899-910.

Sormunen E, Remes J, Hassi J, Pienimäki T, Rintamäki H. Factors associated with self-estimated work ability and musculoskeletal symptoms among male and female workers in cooled food-processing facilities. *Ind Health*. 2009;47(3):271-82.

Sprigg CA, Stride CB, Wall TD, Holman DJ, Smith PR. Work characteristics, musculoskeletal disorders, and the mediating role of psychological strain: a study of call center employees. *The Journal of applied psychology*. 2007;92(5):1456-66.

Spyropoulos P, Papathanasiou G, Georgoudis G, Chronopoulos E, Koutis H, Koumoutsou F. Prevalence of low back pain in greek public office workers. *Pain Physician*. 2007;10(5):651-9.

Stergioulas A, Filippou DK, Triga A, Grigoriadis E, Shipkov CD. Low back pain in physical education teachers. *Folia medica*. 2004;46(3):51-5.

Studnek JR, Crawford JM, Wilkins JR, 3rd, Pennell ML. Back problems among emergency medical services professionals: the LEADS health and wellness follow-up study. *Am J Ind Med*. 2010;53(1):12-22.

Suadicani P, Hansen K, Fenger AM, Gyntelberg F. Low back pain in steelplant workers. *Occup Med (Lond)*. 1994;44(4):217-21.

Svensson HO, Andersson GB. Low-back pain in 40- to 47-year-old men: work history and work environment factors. *Spine (Phila Pa 1976)*. 1983;8(3):272-6.

Svensson HO, Andersson GB. The relationship of low-back pain, work history, work environment, and stress. A retrospective cross-sectional study of 38- to 64-year-old women. *Spine (Phila Pa 1976)*. 1989;14(5):517-22.

Takaki J, Taniguchi T, Hirokawa K. Associations of workplace bullying and harassment with pain. *Int J Environ Res Public Health*. 2013;10(10):4560-70.

Tissot F, Messing K. Perimenstrual symptoms and working conditions among hospital workers in Quebec. *Am J Ind Med*. 1995;27(4):511-22.

Tissot F, Messing K, Stock S. Studying the relationship between low back pain and working postures among those who stand and those who sit most of the working day. *Ergonomics*. 2009;52(11):1402-18.

Toomingas A, Theorell T, Michelsen H, Nordemar R. Associations between self-rated psychosocial work conditions and musculoskeletal symptoms and signs. Stockholm MUSIC I Study Group. *Scand J Work Environ Health*. 1997;23(2):130-9.

Toren A, Oberg K, Lembke B, Enlund K, Rask-Andersen A. Tractor-driving hours and their relation to self-reported low-back and hip symptoms. *Appl Ergon.* 2002;33(2):139-46.

Toroptsova NV, Benevolenskaya LI, Karyakin AN, Sergeev IL, Erdesz S. "Cross-sectional" study of low back pain among workers at an industrial enterprise in Russia. *Spine (Phila Pa 1976).* 1995;20(3):328-32.

Trinkoff AM, Lipscomb JA, Geiger-Brown J, Storr CL, Brady BA. Perceived physical demands and reported musculoskeletal problems in registered nurses. *Am J Prev Med.* 2003;24(3):270-5.

Truszczynska A, Lewkowicz R, Truszczynski O, Rapala K, Wojtkowiak M. Back pain in Polish military helicopter pilots. *International journal of occupational medicine and environmental health.* 2012;25(3):258-64.

Tsigonia A, Tanagra D, Linos A, Merakoulis G, Alexopoulos EC. Musculoskeletal disorders among cosmetologists. *Int J Environ Res Public Health.* 2009;6(12):2967-79.

Urquhart DM, Kelsall HL, Hoe VC, Cicuttini FM, Forbes AB, Sim MR. Are Psychosocial Factors Associated With Low Back Pain and Work Absence for Low Back Pain in an Occupational Cohort? *Clin J Pain.* 2013.

Urquhart DM, Phyomaung PP, Wluka AE, Sim MR, Forbes A, Jones G, et al. Is there a relationship between occupational activities and low back pain in obese, middle-aged women? *Climacteric.* 2013.

Walsh K, Cruddas M, Coggon D. Interaction of height and mechanical loading of the spine in the development of low-back pain. *Scand J Work Environ Health.* 1991;17(6):420-4.

Walsh K, Varnes N, Osmond C, Styles R, Coggon D. Occupational causes of low-back pain. *Scand J Work Environ Health.* 1989;15(1):54-9.

Van Nieuwenhuysse A, Fatkhutdinova L, Verbeke G, Pirenne D, Johannik K, Somville PR, et al. Risk factors for first-ever low back pain among workers in their first employment. *Occup Med (Lond).* 2004;54(8):513-9.

Vasiliadou A, Karvountzis G, Roumeliotis D, Soumilas A, Plati C, Nomikos I. Factors associated with back pain in nursing staff: a survey in Athens, Greece. *International journal of nursing practice.* 1997;3(1):15-20.

- Vasiliadou A, Karvountzis GG, Soumilas A, Roumeliotis D, Theodosopoulou E. Occupational low-back pain in nursing staff in a Greek hospital. *Journal of Advanced Nursing*. 1995;21(1):125-30.
- Widanarko B, Legg S, Stevenson M, Devereux J, Eng A, t Mannelje A, et al. Prevalence and work-related risk factors for reduced activities and absenteeism due to low back symptoms. *Appl Ergon*. 2012;43(4):727-37.
- Videman T, Levalahti E, Battie MC. The effects of anthropometrics, lifting strength, and physical activities in disc degeneration. *Spine (Phila Pa 1976)*. 2007;32(13):1406-13.
- Vieira ER, Kumar S, Coury HJ, Narayan Y. Low back problems and possible improvements in nursing jobs. *J Adv Nurs*. 2006;55(1):79-89.
- Violante FS, Fiori M, Fiorentini C, Risi A, Garagnani G, Bonfiglioli R, et al. Associations of psychosocial and individual factors with three different categories of back disorder among nursing staff. *J Occup Health*. 2004;46(2):100-8.
- Violante FS, Graziosi F, Bonfiglioli R, Curti S, Mattioli S. Relations between occupational, psychosocial and individual factors and three different categories of back disorder among supermarket workers. *Int Arch Occup Environ Health*. 2005;78(8):613-24.
- Yildirim Y, Gunay S, Karadibak D. Identifying factors associated with low back pain among employees working at a package producing industry. *J Back Musculoskelet Rehabil*. 2013.
- Zoidaki A, Riza E, Kastania A, Papadimitriou E, Linos A. Musculoskeletal disorders among dentists in the Greater Athens area, Greece: risk factors and correlations. *Journal of Public Health*.(2012):1-11.